



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:33 PM GMT

PDB ID : 4P8T
Title : Crystal structure of M. tuberculosis DprE1 in complex with the non-covalent inhibitor QN129
Authors : Neres, J.; Pojer, F.; Cole, S.T.
Deposited on : 2014-04-01
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4549 (2.58-2.50)
Clashscore	102246	5292 (2.58-2.50)
Ramachandran outliers	100387	5194 (2.58-2.50)
Sidechain outliers	100360	5196 (2.58-2.50)
RSRZ outliers	91569	4561 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	<div> <div>5%</div> <div> <div></div> <div>81%</div> <div>13%</div> <div>• 5%</div> </div> </div>
1	B	480	<div> <div>4%</div> <div> <div></div> <div>78%</div> <div>9%</div> <div>• 13%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	2J3	A	504	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7012 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable decaprenylphosphoryl-beta-D-ribose oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	1	0
			3511	2234	617	648	12			
1	B	419	Total	C	N	O	S	0	0	0
			3219	2042	571	596	10			

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	GLY	-	expression tag	UNP P72056
A	-17	SER	-	expression tag	UNP P72056
A	-16	SER	-	expression tag	UNP P72056
A	-15	HIS	-	expression tag	UNP P72056
A	-14	HIS	-	expression tag	UNP P72056
A	-13	HIS	-	expression tag	UNP P72056
A	-12	HIS	-	expression tag	UNP P72056
A	-11	HIS	-	expression tag	UNP P72056
A	-10	HIS	-	expression tag	UNP P72056
A	-9	SER	-	expression tag	UNP P72056
A	-8	SER	-	expression tag	UNP P72056
A	-7	GLY	-	expression tag	UNP P72056
A	-6	LEU	-	expression tag	UNP P72056
A	-5	VAL	-	expression tag	UNP P72056
A	-4	PRO	-	expression tag	UNP P72056
A	-3	ARG	-	expression tag	UNP P72056
A	-2	GLY	-	expression tag	UNP P72056
A	-1	SER	-	expression tag	UNP P72056
A	0	HIS	-	expression tag	UNP P72056
B	-18	GLY	-	expression tag	UNP P72056
B	-17	SER	-	expression tag	UNP P72056
B	-16	SER	-	expression tag	UNP P72056
B	-15	HIS	-	expression tag	UNP P72056
B	-14	HIS	-	expression tag	UNP P72056
B	-13	HIS	-	expression tag	UNP P72056

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	HIS	-	expression tag	UNP P72056
B	-11	HIS	-	expression tag	UNP P72056
B	-10	HIS	-	expression tag	UNP P72056
B	-9	SER	-	expression tag	UNP P72056
B	-8	SER	-	expression tag	UNP P72056
B	-7	GLY	-	expression tag	UNP P72056
B	-6	LEU	-	expression tag	UNP P72056
B	-5	VAL	-	expression tag	UNP P72056
B	-4	PRO	-	expression tag	UNP P72056
B	-3	ARG	-	expression tag	UNP P72056
B	-2	GLY	-	expression tag	UNP P72056
B	-1	SER	-	expression tag	UNP P72056
B	0	HIS	-	expression tag	UNP P72056

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is 3-[(4-cyanobenzyl)amino]-6-(trifluoromethyl)quinoxaline-2-carboxylic acid (three-letter code: R26) (formula: $C_{18}H_{11}F_3N_4O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			27	18	3	4	2		
3	B	1	Total	C	F	N	O	0	0
			27	18	3	4	2		


- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula: C₃H₅N₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C N	0	0
			5	3 2		

- Molecule 5 is (2R)-2-([(2R)-2-([(2R)-2-hydroxypropyl]oxy)propyl]oxy)propan-1-ol

(three-letter code: 2J3) (formula: C₉H₂₀O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			13	9	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	60	Total	O	0	0
			60	60		
6	B	44	Total	O	0	0
			44	44		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable decaprenylphosphoryl-beta-D-ribose oxidase



- Molecule 1: Probable decaprenylphosphoryl-beta-D-ribose oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.23 Å 83.04 Å 80.73 Å 90.00° 102.84° 90.00°	Depositor
Resolution (Å)	49.50 – 2.55 49.52 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.6 (49.50-2.55) 99.6 (49.52-2.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.54 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.184 , 0.237 0.184 , 0.237	Depositor DCC
R_{free} test set	1625 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.7	EDS
Estimated twinning fraction	0.025 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 32500 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7012	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: R26, 2J3, IMD, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	4/3600 (0.1%)	0.59	0/4893
1	B	0.51	4/3292 (0.1%)	0.57	0/4466
All	All	0.52	8/6892 (0.1%)	0.58	0/9359

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	16	TRP	CD2-CE2	5.35	1.47	1.41
1	A	384	TRP	CD2-CE2	5.32	1.47	1.41
1	B	230	TRP	CD2-CE2	5.29	1.47	1.41
1	A	172	TRP	CD2-CE2	5.24	1.47	1.41
1	A	230	TRP	CD2-CE2	5.24	1.47	1.41
1	A	16	TRP	CD2-CE2	5.12	1.47	1.41
1	B	172	TRP	CD2-CE2	5.06	1.47	1.41
1	B	296	TRP	CD2-CE2	5.04	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3511	0	3479	41	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3219	0	3198	24	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	27	0	10	0	0
3	B	27	0	10	0	0
4	A	5	0	5	0	0
5	A	13	0	20	4	0
6	A	60	0	0	2	0
6	B	44	0	0	1	0
All	All	7012	0	6784	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:ARG:HH11	1:B:159:ARG:HG2	1.00	1.12
1:A:325:ARG:N	1:A:326:ALA:HA	1.67	1.09
1:A:325:ARG:H	1:A:326:ALA:HA	1.02	1.09
1:A:442:ARG:HG3	1:A:442:ARG:HH11	1.21	1.02
1:B:159:ARG:NH1	1:B:159:ARG:HG2	1.80	0.90
1:B:347:PHE:CE1	1:B:386:ILE:HD11	2.08	0.88
1:A:325:ARG:H	1:A:326:ALA:CA	1.88	0.87
1:A:442:ARG:CG	1:A:442:ARG:HH11	1.89	0.86
1:A:41:ARG:O	1:A:44:GLU:HG2	1.80	0.81
1:B:159:ARG:CG	1:B:159:ARG:HH11	1.92	0.78
1:B:347:PHE:HE1	1:B:386:ILE:HD11	1.49	0.76
1:A:221:GLU:HG2	6:A:659:HOH:O	1.88	0.74
1:B:93:ASP:O	1:B:96:VAL:HG13	1.88	0.73
1:A:119:ARG:HH11	1:A:119:ARG:HG3	1.54	0.72
1:A:83:ILE:HG12	1:A:90:VAL:HG12	1.77	0.67
1:A:442:ARG:NH1	1:A:442:ARG:HG3	2.01	0.66
1:A:119:ARG:HH11	1:A:119:ARG:CG	2.08	0.65
1:A:240:LEU:HD22	1:A:355:GLN:HG2	1.80	0.63
1:B:83:ILE:HG12	1:B:90:VAL:HG12	1.81	0.61
1:A:325:ARG:N	1:A:326:ALA:CA	2.51	0.59
1:B:347:PHE:HE1	1:B:386:ILE:CD1	2.15	0.57
1:A:60:TYR:CE1	1:A:334:GLN:HG3	2.40	0.57
1:A:442:ARG:CG	1:A:442:ARG:NH1	2.58	0.56
1:A:206:THR:HB	1:A:211:GLU:HB3	1.88	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:102:MET:HG3	6:A:647:HOH:O	2.05	0.55
1:A:316:PRO:HG2	5:A:504:2J3:H4	1.89	0.54
1:A:116:PRO:HG2	1:A:118:THR:O	2.10	0.52
1:A:61:GLY:HA3	1:A:419:ASP:O	2.09	0.52
1:A:119:ARG:NH1	1:A:119:ARG:HG3	2.26	0.51
1:B:153:THR:HG21	1:B:159:ARG:HD3	1.93	0.51
1:A:200:ILE:HD11	1:A:255:GLN:OE1	2.10	0.51
1:A:256:LEU:O	1:A:261:ARG:NH2	2.43	0.50
1:B:206:THR:O	1:B:242:ARG:HB2	2.10	0.50
1:A:75:THR:N	1:A:76:PRO:HD2	2.27	0.50
1:A:209:LEU:HB2	1:A:241:GLY:HA3	1.95	0.49
1:B:99:ASP:OD2	1:B:119:ARG:NH1	2.46	0.48
1:B:270:PRO:HG3	1:B:311:THR:HG21	1.96	0.48
1:B:461:LEU:HD12	1:B:461:LEU:H	1.79	0.47
1:B:231:PHE:CE2	1:B:364:ASN:HB3	2.49	0.47
1:B:75:THR:N	1:B:76:PRO:HD2	2.30	0.47
1:A:375:ALA:HB3	1:A:378:SER:HB2	1.96	0.47
1:B:240:LEU:HD22	1:B:355:GLN:HG2	1.95	0.47
1:A:441:ARG:HH11	1:A:441:ARG:CG	2.29	0.46
1:B:421:ARG:CZ	1:B:421:ARG:HB2	2.45	0.45
1:B:206:THR:HB	1:B:211:GLU:HB3	2.00	0.44
1:B:221:GLU:HG2	6:B:639:HOH:O	2.18	0.44
1:A:103:LYS:NZ	1:A:267:PHE:CD2	2.86	0.44
1:B:209:LEU:HB2	1:B:241:GLY:HA3	2.01	0.43
1:A:228:SER:OG	5:A:504:2J3:H14	2.18	0.43
1:B:26:VAL:HA	1:B:71:VAL:O	2.19	0.43
1:A:61:GLY:HA2	1:A:420:SER:HB3	2.01	0.43
1:A:367:LYS:HG3	1:A:368:LEU:N	2.34	0.43
1:A:46:GLY:HA2	1:A:447:LEU:HD13	2.01	0.43
1:B:147:ARG:HA	1:B:147:ARG:HD3	1.74	0.43
1:A:119:ARG:HB2	1:A:119:ARG:HH11	1.84	0.42
1:A:119:ARG:CB	1:A:119:ARG:HH11	2.32	0.42
1:A:441:ARG:HH11	1:A:441:ARG:HG3	1.85	0.41
1:A:246:SER:HB2	5:A:504:2J3:H7	2.03	0.41
5:A:504:2J3:H12	5:A:504:2J3:H4	1.91	0.41
1:A:92:ILE:HB	1:A:96:VAL:HG11	2.02	0.41
1:B:338:VAL:HA	1:B:384:TRP:O	2.20	0.41
1:B:88:LYS:HE3	1:B:88:LYS:HB2	1.93	0.41
1:A:166:GLU:O	1:A:167:ASP:HB2	2.20	0.41
1:A:330:ALA:HA	1:A:331:GLY:HA2	1.79	0.40
1:A:200:ILE:HG13	1:A:251:ALA:HB2	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:THR:HG21	1:A:159:ARG:HD2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/480 (95%)	442 (97%)	12 (3%)	1 (0%)	52	73
1	B	413/480 (86%)	405 (98%)	8 (2%)	0	100	100
All	All	868/960 (90%)	847 (98%)	20 (2%)	1 (0%)	56	76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	323	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/384 (95%)	345 (94%)	20 (6%)	27	46
1	B	335/384 (87%)	321 (96%)	14 (4%)	36	60
All	All	700/768 (91%)	666 (95%)	34 (5%)	31	52

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	MET
1	A	77	LEU
1	A	88	LYS
1	A	92	ILE
1	A	96	VAL
1	A	119	ARG
1	A	221	GLU
1	A	253	VAL
1	A	272	LEU
1	A	314	TYR
1	A	324	ASN
1	A	344	VAL
1	A	355	GLN
1	A	396	LEU
1	A	406	ARG
1	A	413	ARG
1	A	421	ARG
1	A	441	ARG
1	A	442	ARG
1	A	448	ARG
1	B	37	LYS
1	B	44	GLU
1	B	88	LYS
1	B	92	ILE
1	B	103	LYS
1	B	159	ARG
1	B	205	VAL
1	B	295	LEU
1	B	355	GLN
1	B	396	LEU
1	B	413	ARG
1	B	421	ARG
1	B	422	THR
1	B	425	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	324	ASN
1	B	25	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FAD	A	501	-	48,58,58	1.45	7 (14%)	54,89,89	2.22	11 (20%)
3	R26	A	502	-	25,29,29	1.64	2 (8%)	37,42,42	1.15	3 (8%)
4	IMD	A	503	-	3,5,5	0.37	0	4,5,5	0.66	0
5	2J3	A	504	-	12,12,12	0.75	0	11,14,14	1.20	1 (9%)
2	FAD	B	501	-	48,58,58	1.50	8 (16%)	54,89,89	2.27	10 (18%)
3	R26	B	502	-	25,29,29	1.66	3 (12%)	37,42,42	1.19	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	501	-	-	0/30/50/50	0/6/6/6
3	R26	A	502	-	-	0/13/17/17	0/3/3/3
4	IMD	A	503	-	-	0/0/0/0	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	2J3	A	504	-	-	0/12/12/12	0/0/0/0
2	FAD	B	501	-	-	0/30/50/50	0/6/6/6
3	R26	B	502	-	-	0/13/17/17	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	R26	CAV-CAW	-6.23	1.28	1.44
3	A	502	R26	CAV-CAW	-6.06	1.29	1.44
2	B	501	FAD	C4X-N5	2.03	1.36	1.33
2	A	501	FAD	O4B-C1B	2.15	1.43	1.41
3	B	502	R26	CAC-NAD	2.37	1.35	1.32
2	A	501	FAD	C9A-N10	2.45	1.42	1.38
2	B	501	FAD	C9A-N10	2.48	1.42	1.38
2	B	501	FAD	O4B-C1B	2.56	1.44	1.41
2	A	501	FAD	C8-C7	3.07	1.49	1.41
2	A	501	FAD	C5A-C4A	3.18	1.47	1.40
3	B	502	R26	CAP-NAO	3.22	1.36	1.31
2	B	501	FAD	C5A-C4A	3.34	1.48	1.40
2	B	501	FAD	C8-C7	3.37	1.50	1.41
3	A	502	R26	CAP-NAO	3.59	1.36	1.31
2	B	501	FAD	C9A-C5X	3.67	1.50	1.42
2	B	501	FAD	C4-C4X	3.77	1.48	1.41
2	A	501	FAD	C4-C4X	3.79	1.48	1.41
2	A	501	FAD	C4X-C10	3.83	1.48	1.41
2	A	501	FAD	C9A-C5X	3.98	1.50	1.42
2	B	501	FAD	C4X-C10	4.00	1.48	1.41

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	FAD	N3A-C2A-N1A	-8.01	122.76	128.89
2	A	501	FAD	N3A-C2A-N1A	-7.27	123.33	128.89
2	B	501	FAD	C4-C4X-C10	-6.55	115.75	119.94
2	A	501	FAD	C4-C4X-C10	-6.24	115.95	119.94
2	B	501	FAD	C4X-C4-N3	-3.46	118.85	123.59
2	B	501	FAD	P-O3P-PA	-3.29	123.48	132.73
2	A	501	FAD	C4X-C4-N3	-3.26	119.13	123.59
3	B	502	R26	CAC-CAP-NAO	-2.98	119.57	122.02
2	B	501	FAD	C4A-C5A-N7A	-2.81	106.90	109.48
3	A	502	R26	CAP-CAC-NAD	-2.58	118.73	121.02
2	A	501	FAD	P-O3P-PA	-2.58	125.49	132.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	C1B-N9A-C4A	-2.46	123.23	126.94
2	A	501	FAD	C4A-C5A-N7A	-2.30	107.36	109.48
2	A	501	FAD	C2B-C1B-N9A	-2.27	110.83	114.29
3	A	502	R26	CAC-CAP-NAO	-2.24	120.18	122.02
2	B	501	FAD	O4B-C1B-N9A	2.09	112.48	108.10
3	A	502	R26	CAC-NAD-CAE	2.17	121.37	117.08
5	A	504	2J3	O2-C4-C3	2.47	113.67	108.84
2	B	501	FAD	C4X-N5-C5X	3.14	120.37	116.76
2	A	501	FAD	C4-C4X-N5	3.44	122.90	118.72
2	A	501	FAD	C4X-N5-C5X	3.55	120.84	116.76
2	A	501	FAD	C1'-N10-C9A	3.61	122.91	118.86
2	B	501	FAD	C4-C4X-N5	3.65	123.16	118.72
2	B	501	FAD	C1'-N10-C9A	3.82	123.14	118.86
2	B	501	FAD	C4-N3-C2	8.27	122.39	115.25
2	A	501	FAD	C4-N3-C2	8.33	122.45	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	504	2J3	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	456/480 (95%)	0.15	26 (5%) 27 32	21, 34, 83, 115	0
1	B	419/480 (87%)	0.19	21 (5%) 32 38	25, 43, 82, 102	0
All	All	875/960 (91%)	0.17	47 (5%) 29 34	21, 39, 83, 115	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	47	GLY	7.0
1	A	327	TYR	6.8
1	A	323	TRP	5.8
1	A	324	ASN	5.6
1	A	325	ARG	5.3
1	A	46	GLY	4.8
1	B	46	GLY	4.6
1	A	320	PHE	4.1
1	A	321	GLY	4.0
1	A	329	PRO	3.7
1	A	319	MET	3.7
1	B	8	THR	3.6
1	B	291	PRO	3.6
1	B	268	ASP	3.4
1	A	273	LEU	3.4
1	A	283	LEU	3.3
1	B	292	ILE	3.3
1	B	44	GLU	3.0
1	A	328	GLY	2.9
1	B	18	ARG	2.9
1	A	280	PRO	2.9
1	A	272	LEU	2.9
1	A	322	GLU	2.8
1	B	16	TRP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	270	PRO	2.7
1	B	315	HIS	2.6
1	B	447	LEU	2.6
1	B	34	MET	2.6
1	A	19	THR	2.5
1	A	317	LEU	2.5
1	A	330	ALA	2.5
1	B	37	LYS	2.4
1	A	14	THR	2.4
1	A	278	VAL	2.3
1	B	33	GLU	2.2
1	B	47	GLY	2.2
1	B	269	ALA	2.1
1	A	12	ARG	2.1
1	A	315	HIS	2.1
1	B	394	ASP	2.1
1	A	362	PHE	2.1
1	B	15	GLY	2.0
1	B	9	THR	2.0
1	A	18	ARG	2.0
1	A	326	ALA	2.0
1	B	10	ALA	2.0
1	B	41	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	2J3	A	504	13/13	0.85	0.22	3.57	62,67,69,69	0
3	R26	B	502	27/27	0.97	0.19	0.65	44,46,54,58	0
3	R26	A	502	27/27	0.96	0.19	0.00	35,37,50,53	0
2	FAD	A	501	53/53	0.98	0.14	-0.13	24,26,29,30	0
2	FAD	B	501	53/53	0.98	0.13	-0.50	26,30,31,33	0
4	IMD	A	503	5/5	0.98	0.13	-	40,40,42,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.