



# Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 07:34 PM GMT

PDB ID : 4PAA  
Title : Crystal structure of the mature form of rat DMGDH complexed with tetrahydrofolate  
Authors : Luka, Z.; Pakhomova, S.; Loukachevitch, L.V.; Newcomer, M.E.; Wagner, C.  
Deposited on : 2014-04-07  
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1640 (2.28-2.24)
Clashscore	102246	1095 (2.26-2.26)
Ramachandran outliers	100387	1063 (2.26-2.26)
Sidechain outliers	100360	1063 (2.26-2.26)
RSRZ outliers	91569	1647 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	848	%	89%	6%	..
1	B	848	2%	89%	7%	.

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 13358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dimethylglycine dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	816	Total	C	N	O	S	0	0	0
			6476	4143	1111	1203	19			
1	B	816	Total	C	N	O	S	0	0	0
			6476	4143	1111	1203	19			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	expression tag	UNP Q5RKL4
A	21	ALA	-	expression tag	UNP Q5RKL4
A	858	ALA	-	expression tag	UNP Q5RKL4
A	859	ALA	-	expression tag	UNP Q5RKL4
A	860	LEU	-	expression tag	UNP Q5RKL4
A	861	GLU	-	expression tag	UNP Q5RKL4
A	862	HIS	-	expression tag	UNP Q5RKL4
A	863	HIS	-	expression tag	UNP Q5RKL4
A	864	HIS	-	expression tag	UNP Q5RKL4
A	865	HIS	-	expression tag	UNP Q5RKL4
A	866	HIS	-	expression tag	UNP Q5RKL4
A	867	HIS	-	expression tag	UNP Q5RKL4
B	20	MET	-	expression tag	UNP Q5RKL4
B	21	ALA	-	expression tag	UNP Q5RKL4
B	858	ALA	-	expression tag	UNP Q5RKL4
B	859	ALA	-	expression tag	UNP Q5RKL4
B	860	LEU	-	expression tag	UNP Q5RKL4
B	861	GLU	-	expression tag	UNP Q5RKL4
B	862	HIS	-	expression tag	UNP Q5RKL4
B	863	HIS	-	expression tag	UNP Q5RKL4
B	864	HIS	-	expression tag	UNP Q5RKL4
B	865	HIS	-	expression tag	UNP Q5RKL4
B	866	HIS	-	expression tag	UNP Q5RKL4
B	867	HIS	-	expression tag	UNP Q5RKL4

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0
2	B	1	53	27	9	15	2	0	0

- Molecule 3 is (6S)-5,6,7,8-TETRAHYDROFOLATE (three-letter code: THG) (formula: C<sub>19</sub>H<sub>23</sub>N<sub>7</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 32 19 7 6	0	0
3	B	1	Total C N O 32 19 7 6	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	116	Total O 116 116	0	0
4	B	120	Total O 120 120	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dimethylglycine dehydrogenase



- Molecule 1: Dimethylglycine dehydrogenase



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.93 Å    130.66 Å    171.86 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	40.00 – 2.26 49.58 – 2.26	Depositor EDS
% Data completeness (in resolution range)	96.7 (40.00-2.26) 96.7 (49.58-2.26)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.38 (at 2.27 Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
$R$ , $R_{free}$	0.241 , 0.267 0.244 , 0.268	Depositor DCC
$R_{free}$ test set	1743 reflections (2.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.3	Xtriage
Anisotropy	0.520	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 24.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$<  L  > = 0.45$ , $< L^2 > = 0.28$	Xtriage
Outliers	1 of 87180 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	13358	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: THG, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/6641	0.59	1/9015 (0.0%)
1	B	0.38	0/6641	0.58	0/9015
All	All	0.38	0/13282	0.59	1/18030 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	850	LEU	CA-CB-CG	5.95	128.99	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6476	0	6409	26	0
1	B	6476	0	6409	23	0
2	A	53	0	29	1	0
2	B	53	0	30	1	0
3	A	32	0	21	0	0
3	B	32	0	21	0	0
4	A	116	0	0	2	0
4	B	120	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	13358	0	12919	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:ILE:HG21	1:A:351:VAL:CG1	2.21	0.71
1:A:271:ILE:HG21	1:A:351:VAL:HG11	1.77	0.67
1:A:271:ILE:HD13	1:A:351:VAL:CG1	2.30	0.61
1:A:349:VAL:HB	1:A:352:LEU:HD23	1.88	0.56
1:A:497:GLN:NE2	4:A:1075:HOH:O	2.38	0.55
1:B:131:ALA:HB2	1:B:140:PHE:CE2	2.43	0.54
1:B:262:GLN:HE21	1:B:314:GLN:NE2	2.06	0.53
1:A:271:ILE:HG21	1:A:351:VAL:HG13	1.92	0.52
1:B:46:VAL:HG23	1:B:234:ALA:HB2	1.92	0.52
1:A:131:ALA:HB2	1:A:140:PHE:CE2	2.44	0.51
1:B:488:TRP:CE2	1:B:507:TRP:HB3	2.46	0.51
1:A:46:VAL:HG23	1:A:234:ALA:HB2	1.92	0.51
1:A:285:ARG:NH2	4:A:1031:HOH:O	2.45	0.49
1:A:488:TRP:CE2	1:A:507:TRP:HB3	2.47	0.49
1:B:797:SER:HB2	4:B:1057:HOH:O	2.13	0.49
1:A:98:LEU:HD21	1:A:420:LEU:HD11	1.95	0.49
1:B:839:PRO:HG2	1:B:842:LEU:HD23	1.95	0.48
1:B:247:GLU:HG3	4:B:1090:HOH:O	2.12	0.48
1:B:98:LEU:HD21	1:B:420:LEU:HD11	1.96	0.47
1:B:263:HIS:HD2	1:B:442:TYR:OH	1.98	0.46
1:A:839:PRO:HG2	1:A:842:LEU:HD23	1.96	0.46
1:B:264:GLN:HA	1:B:361:VAL:O	2.16	0.46
1:A:496:THR:HG21	1:B:134:PRO:HB2	1.97	0.46
1:A:349:VAL:HB	1:A:352:LEU:CD2	2.46	0.45
1:A:264:GLN:HA	1:A:361:VAL:O	2.17	0.45
1:A:271:ILE:HD13	1:A:351:VAL:HG12	1.99	0.45
1:B:74:LYS:HG3	1:B:211:PRO:HB3	1.99	0.45
1:B:262:GLN:HE21	1:B:314:GLN:HE22	1.64	0.44
1:B:187:ASP:HB3	1:B:190:SER:OG	2.17	0.44
1:A:74:LYS:HG3	1:A:211:PRO:HB3	1.99	0.43
1:B:275:LYS:HD3	1:B:298:ASP:CG	2.38	0.43
1:A:291:TYR:OH	1:A:345:ALA:HA	2.19	0.43
1:A:265:TYR:CE2	2:A:901:FAD:HM72	2.54	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:VAL:O	1:A:364:PRO:HA	2.19	0.43
1:B:291:TYR:OH	1:B:345:ALA:HA	2.19	0.43
1:B:595:HIS:HD2	1:B:596:ASP:OD1	2.01	0.43
1:A:595:HIS:HD2	1:A:596:ASP:OD1	2.03	0.42
1:B:258:LEU:HD11	1:B:388:ILE:HD12	2.01	0.42
1:B:261:VAL:O	1:B:364:PRO:HA	2.19	0.42
1:A:475:CYS:HB3	1:A:489:PHE:CD1	2.55	0.42
1:A:258:LEU:HD11	1:A:388:ILE:HD12	2.01	0.41
1:A:297:ARG:HD3	1:A:297:ARG:HA	1.90	0.41
1:A:268:THR:HG21	1:A:271:ILE:HD11	2.01	0.41
1:A:270:THR:HG23	1:A:298:ASP:HB2	2.01	0.41
1:B:297:ARG:N	1:B:297:ARG:HD3	2.36	0.41
1:B:270:THR:HG23	1:B:298:ASP:HB2	2.02	0.41
1:B:93:HIS:CE1	1:B:98:LEU:HD13	2.56	0.40
1:B:265:TYR:CE2	2:B:901:FAD:HM72	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	814/848 (96%)	789 (97%)	23 (3%)	2 (0%)	52 61
1	B	814/848 (96%)	789 (97%)	23 (3%)	2 (0%)	52 61
All	All	1628/1696 (96%)	1578 (97%)	46 (3%)	4 (0%)	52 61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	371	ILE
1	B	371	ILE
1	A	593	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	593	GLU

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	689/714 (96%)	664 (96%)	25 (4%)	42 51
1	B	689/714 (96%)	665 (96%)	24 (4%)	43 53
All	All	1378/1428 (96%)	1329 (96%)	49 (4%)	42 51

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	ARG
1	A	268	THR
1	A	285	ARG
1	A	313	LEU
1	A	316	SER
1	A	328	LYS
1	A	329	GLU
1	A	339	THR
1	A	352	LEU
1	A	371	ILE
1	A	404	LEU
1	A	497	GLN
1	A	567	ARG
1	A	614	ARG
1	A	647	PHE
1	A	658	SER
1	A	713	LEU
1	A	724	MET
1	A	731	LEU
1	A	770	LEU
1	A	773	ASP
1	A	803	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	813	VAL
1	A	817	GLU
1	A	850	LEU
1	B	38	GLU
1	B	41	ASP
1	B	158	GLU
1	B	216	LYS
1	B	275	LYS
1	B	297	ARG
1	B	313	LEU
1	B	328	LYS
1	B	336	ASP
1	B	347	GLU
1	B	356	ASP
1	B	368	SER
1	B	371	ILE
1	B	550	ASN
1	B	647	PHE
1	B	690	ARG
1	B	713	LEU
1	B	724	MET
1	B	731	LEU
1	B	736	ASP
1	B	746	ASN
1	B	770	LEU
1	B	773	ASP
1	B	813	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	118	GLN
1	A	497	GLN
1	A	579	GLN
1	A	803	GLN
1	A	821	GLN
1	B	263	HIS
1	B	314	GLN
1	B	550	ASN
1	B	579	GLN
1	B	821	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	FAD	A	901	1	48,58,58	1.31	7 (14%)	54,89,89	2.20	8 (14%)
3	THG	A	902	-	26,34,34	1.11	3 (11%)	28,47,47	2.30	11 (39%)
2	FAD	B	901	1	48,58,58	1.28	7 (14%)	54,89,89	2.09	9 (16%)
3	THG	B	902	-	26,34,34	1.05	1 (3%)	28,47,47	1.58	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	901	1	-	0/30/50/50	0/6/6/6
3	THG	A	902	-	-	0/16/31/31	0/3/3/3
2	FAD	B	901	1	-	0/30/50/50	0/6/6/6
3	THG	B	902	-	-	0/16/31/31	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	FAD	C1'-N10	-2.33	1.46	1.48
3	A	902	THG	C4A-N5	-2.16	1.33	1.38
2	B	901	FAD	C9A-N10	2.15	1.41	1.38
2	B	901	FAD	C2A-N3A	2.31	1.36	1.32
2	A	901	FAD	C4-C4X	2.67	1.46	1.41
3	A	902	THG	C4A-C8A	2.70	1.47	1.41
2	B	901	FAD	C5A-C4A	2.76	1.46	1.40
2	A	901	FAD	C8-C7	2.76	1.48	1.41
3	A	902	THG	C7-C6	2.77	1.55	1.52
2	B	901	FAD	C9A-C5X	2.86	1.48	1.42
2	A	901	FAD	C9A-N10	2.86	1.42	1.38
2	A	901	FAD	C5A-C4A	2.94	1.47	1.40
2	A	901	FAD	C9A-C5X	3.21	1.49	1.42
2	B	901	FAD	C4-C4X	3.24	1.47	1.41
2	B	901	FAD	C8-C7	3.29	1.49	1.41
2	B	901	FAD	C4X-C10	3.52	1.47	1.41
2	A	901	FAD	C4X-C10	3.93	1.48	1.41
3	B	902	THG	C4A-C8A	3.94	1.49	1.41

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	FAD	N3A-C2A-N1A	-8.37	122.49	128.89
2	B	901	FAD	N3A-C2A-N1A	-7.75	122.96	128.89
2	A	901	FAD	C4-C4X-C10	-5.45	116.45	119.94
2	B	901	FAD	C4-C4X-C10	-5.03	116.72	119.94
2	A	901	FAD	C4X-C10-N10	-4.65	117.78	120.52
3	A	902	THG	C9-C6-N5	-4.12	100.82	109.34
2	B	901	FAD	C4X-C4-N3	-3.13	119.31	123.59
2	A	901	FAD	C4X-C4-N3	-3.00	119.48	123.59
3	A	902	THG	C5'-C4'-N10	-2.96	115.40	121.06
3	B	902	THG	C5'-C4'-N10	-2.87	115.57	121.06
2	B	901	FAD	O2'-C2'-C1'	-2.61	103.53	109.94
2	B	901	FAD	C4A-C5A-N7A	-2.47	107.20	109.48
3	B	902	THG	C3'-C4'-N10	2.26	125.39	121.06
3	A	902	THG	C9-N10-C4'	2.26	127.16	121.46
3	A	902	THG	C1'-C11-N	2.33	121.08	116.93
3	B	902	THG	C2-N1-C8A	2.60	120.38	114.54
3	A	902	THG	C8A-C4A-N5	2.62	122.16	118.85
2	A	901	FAD	C4X-N5-C5X	2.93	120.13	116.76
3	A	902	THG	C7-C6-N5	2.93	116.55	110.51
3	B	902	THG	C4-C4A-C8A	3.04	117.31	114.56

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	A	902	THG	C3'-C4'-N10	3.07	126.94	121.06
2	B	901	FAD	C4-C4X-N5	3.16	122.56	118.72
3	A	902	THG	C2-N1-C8A	3.20	121.72	114.54
2	A	901	FAD	C1'-N10-C9A	3.32	122.59	118.86
2	A	901	FAD	C5X-C9A-N10	3.61	120.36	117.62
2	B	901	FAD	C1'-N10-C9A	3.72	123.04	118.86
3	A	902	THG	C4-N3-C2	3.91	121.36	115.94
3	A	902	THG	C6-C9-N10	4.09	125.01	113.30
2	B	901	FAD	C4X-N5-C5X	4.09	121.47	116.76
3	B	902	THG	C4-N3-C2	5.05	122.94	115.94
3	A	902	THG	C4-C4A-C8A	5.37	119.42	114.56
2	B	901	FAD	C4-N3-C2	6.82	121.14	115.25
2	A	901	FAD	C4-N3-C2	7.27	121.53	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	FAD	1	0
2	B	901	FAD	1	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	816/848 (96%)	0.05	7 (0%) 85 87	14, 23, 38, 89	0
1	B	816/848 (96%)	0.08	17 (2%) 67 71	12, 25, 45, 82	0
All	All	1632/1696 (96%)	0.07	24 (1%) 76 79	12, 24, 42, 89	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	853	ASP	5.2
1	B	853	ASP	4.9
1	B	279	ARG	3.7
1	A	851	GLN	3.3
1	B	850	LEU	3.3
1	A	696	GLN	3.2
1	B	851	GLN	3.2
1	B	696	GLN	3.0
1	B	852	LYS	2.8
1	B	357	ILE	2.8
1	A	849	ARG	2.7
1	B	38	GLU	2.7
1	B	646	VAL	2.6
1	B	645	ASP	2.6
1	B	697	GLU	2.5
1	B	659	ASP	2.5
1	B	208	TYR	2.4
1	B	155	TYR	2.3
1	A	850	LEU	2.3
1	B	557	PHE	2.2
1	A	320	HIS	2.2
1	B	325	GLY	2.2
1	A	354	LYS	2.1
1	B	271	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	THG	B	902	32/32	0.85	0.17	1.67	26,29,35,36	0
3	THG	A	902	32/32	0.86	0.15	0.94	27,30,33,34	0
2	FAD	A	901	53/53	0.95	0.12	-0.18	15,19,21,21	0
2	FAD	B	901	53/53	0.95	0.10	-0.60	12,13,16,18	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.