



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 04:10 PM BST

PDB ID : 1PEH
Title : NMR STRUCTURE OF THE MEMBRANE-BINDING DOMAIN OF CTP
PHOSPHOCHOLINE CYTIDYLYLTRANSFERASE, 10 STRUCTURES
Authors : Dunne, S.J.; Cornell, R.B.; Johnson, J.E.; Glover, N.R.; Tracey, A.S.
Deposited on : 1996-06-10

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	35	<div> <div>9%</div> <div>29%</div> <div>14%</div> <div>49%</div> </div>

2 Ensemble composition and analysis

This entry contains 10 models. Model 6 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:251-A:268 (18)	0.23	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 5, 6, 7
2	4, 8, 9
Single-model clusters	3; 10

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 604 atoms, of which 311 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PEPNH1.

Mol	Chain	Residues	Atoms					Trace
1	A	35	Total	C	H	N	O	1
			604	185	311	53	55	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	263	TRP	PHE	ENGINEERED	UNP P19836

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: PEPNH1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: PEPNH1




4.2.2 Score per residue for model 2

- Molecule 1: PEPNH1




4.2.3 Score per residue for model 3

- Molecule 1: PEPNH1




4.2.4 Score per residue for model 4

- Molecule 1: PEPNH1




4.2.5 Score per residue for model 5

- Molecule 1: PEPNH1




4.2.6 Score per residue for model 6 (medoid)

- Molecule 1: PEPNH1




4.2.7 Score per residue for model 7

- Molecule 1: PEPNH1




4.2.8 Score per residue for model 8

- Molecule 1: PEPNH1




4.2.9 Score per residue for model 9

- Molecule 1: PEPNH1



4.2.10 Score per residue for model 10

- Molecule 1: PEPNH1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *RESTRAINED DISTANCE GEOMETRY*.

Of the 10 calculated structures, 10 were deposited, based on the following criterion: *ALL SUBMITTED*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DGII	refinement	
BIOSYM MOLECULAR SIMULATIONS/FELIX	structure solution	SIMULATIONS/FELIX

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	2.06±0.00	6±1/156 (3.5±0.3%)	1.65±0.01	3±0/205 (1.6±0.2%)
All	All	2.06	55/1560 (3.5%)	1.65	32/2050 (1.6%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	268	GLU	CD-OE2	10.66	1.37	1.25	3	7
1	A	268	GLU	CD-OE1	10.55	1.37	1.25	1	3
1	A	258	GLU	CD-OE2	10.14	1.36	1.25	4	7
1	A	257	GLU	CD-OE1	10.05	1.36	1.25	3	6
1	A	257	GLU	CD-OE2	10.05	1.36	1.25	1	4
1	A	262	GLU	CD-OE2	10.04	1.36	1.25	2	6
1	A	262	GLU	CD-OE1	9.99	1.36	1.25	10	4
1	A	258	GLU	CD-OE1	9.99	1.36	1.25	9	3
1	A	255	ASP	CG-OD2	5.26	1.37	1.25	4	7
1	A	268	GLU	CB-CG	5.25	1.62	1.52	5	5
1	A	255	ASP	CG-OD1	5.24	1.37	1.25	9	3

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	255	ASP	CB-CG-OD2	-5.96	112.93	118.30	4	10
1	A	255	ASP	CB-CG-OD1	-5.96	112.94	118.30	9	10
1	A	263	TRP	CD1-NE1-CE2	-5.82	103.76	109.00	2	10
1	A	267	VAL	CA-CB-CG1	5.67	119.41	110.90	7	2

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	155	165	165	3±2
All	All	1550	1650	1650	32

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:253:VAL:O	1:A:257:GLU:HB3	0.73	1.83	6	5
1:A:267:VAL:HG12	1:A:268:GLU:HG3	0.73	1.61	7	4
1:A:257:GLU:O	1:A:257:GLU:HG3	0.65	1.92	7	2
1:A:264:VAL:HG12	1:A:268:GLU:OE2	0.64	1.92	6	1
1:A:257:GLU:HG3	1:A:257:GLU:O	0.64	1.93	10	3
1:A:251:LYS:C	1:A:252:LYS:HD3	0.61	2.15	9	1
1:A:261:LYS:O	1:A:264:VAL:HG23	0.60	1.97	5	3
1:A:267:VAL:HG12	1:A:268:GLU:CG	0.54	2.33	5	2
1:A:251:LYS:O	1:A:251:LYS:HG3	0.54	2.02	8	1
1:A:261:LYS:CG	1:A:262:GLU:N	0.50	2.75	10	1
1:A:264:VAL:HG22	1:A:268:GLU:OE2	0.47	2.10	9	1
1:A:267:VAL:CG1	1:A:268:GLU:N	0.47	2.77	7	2
1:A:251:LYS:HG2	1:A:251:LYS:O	0.47	2.10	4	1
1:A:266:LYS:CG	1:A:267:VAL:N	0.46	2.78	6	1
1:A:252:LYS:HD3	1:A:252:LYS:N	0.44	2.28	9	1
1:A:253:VAL:O	1:A:257:GLU:HB2	0.42	2.14	1	1
1:A:252:LYS:HB2	1:A:252:LYS:HE3	0.42	1.65	2	1
1:A:267:VAL:CG1	1:A:268:GLU:HG3	0.40	2.42	7	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	18/35 (51%)	16±1 (91±4%)	2±1 (9±4%)	0±0 (0±0%)	100	100
All	All	180/350 (51%)	164 (91%)	16 (9%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	18/33 (55%)	10±2 (53±10%)	8±2 (47±10%)	0	2
All	All	180/330 (55%)	96 (53%)	84 (47%)	0	2

All 15 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	253	VAL	10
1	A	264	VAL	9
1	A	258	GLU	7
1	A	257	GLU	7
1	A	251	LYS	7
1	A	256	VAL	7
1	A	252	LYS	6
1	A	266	LYS	6
1	A	259	LYS	5
1	A	255	ASP	5
1	A	254	LYS	4
1	A	267	VAL	3
1	A	262	GLU	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	261	LYS	3
1	A	260	SER	2

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided