



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:59 PM GMT

PDB ID : 4PFD  
Title : Crystal structure of M. tuberculosis in complex with a cBT - non-covalent adduct  
Authors : Neres, J.; Panda, M.; Cole, S.  
Deposited on : 2014-04-28  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	 4% 82% 6% • 11%
1	B	480	 3% 75% 9% • 15%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IMD	B	503	-	-	-	X
4	2R2	B	502	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6779 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable decaprenylphosphoryl-beta-D-ribose oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	427	Total	C	N	O	S	0	2	0
			3290	2088	585	606	11			
1	B	410	Total	C	N	O	S	0	1	0
			3141	1989	558	583	11			

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	GLY	-	expression tag	UNP P9WJF1
A	-17	SER	-	expression tag	UNP P9WJF1
A	-16	SER	-	expression tag	UNP P9WJF1
A	-15	HIS	-	expression tag	UNP P9WJF1
A	-14	HIS	-	expression tag	UNP P9WJF1
A	-13	HIS	-	expression tag	UNP P9WJF1
A	-12	HIS	-	expression tag	UNP P9WJF1
A	-11	HIS	-	expression tag	UNP P9WJF1
A	-10	HIS	-	expression tag	UNP P9WJF1
A	-9	SER	-	expression tag	UNP P9WJF1
A	-8	SER	-	expression tag	UNP P9WJF1
A	-7	GLY	-	expression tag	UNP P9WJF1
A	-6	LEU	-	expression tag	UNP P9WJF1
A	-5	VAL	-	expression tag	UNP P9WJF1
A	-4	PRO	-	expression tag	UNP P9WJF1
A	-3	ARG	-	expression tag	UNP P9WJF1
A	-2	GLY	-	expression tag	UNP P9WJF1
A	-1	SER	-	expression tag	UNP P9WJF1
A	0	HIS	-	expression tag	UNP P9WJF1
B	-18	GLY	-	expression tag	UNP P9WJF1
B	-17	SER	-	expression tag	UNP P9WJF1
B	-16	SER	-	expression tag	UNP P9WJF1
B	-15	HIS	-	expression tag	UNP P9WJF1
B	-14	HIS	-	expression tag	UNP P9WJF1
B	-13	HIS	-	expression tag	UNP P9WJF1

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	HIS	-	expression tag	UNP P9WJF1
B	-11	HIS	-	expression tag	UNP P9WJF1
B	-10	HIS	-	expression tag	UNP P9WJF1
B	-9	SER	-	expression tag	UNP P9WJF1
B	-8	SER	-	expression tag	UNP P9WJF1
B	-7	GLY	-	expression tag	UNP P9WJF1
B	-6	LEU	-	expression tag	UNP P9WJF1
B	-5	VAL	-	expression tag	UNP P9WJF1
B	-4	PRO	-	expression tag	UNP P9WJF1
B	-3	ARG	-	expression tag	UNP P9WJF1
B	-2	GLY	-	expression tag	UNP P9WJF1
B	-1	SER	-	expression tag	UNP P9WJF1
B	0	HIS	-	expression tag	UNP P9WJF1

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula:  $C_3H_5N_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			5	3	2		
3	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 4 is [4-(2-methoxyethyl)piperazin-1-yl][6-methyl-7-nitro-5-(trifluoromethyl)-1,3-benzothiazol-2-yl]methanone (three-letter code: 2R2) (formula: C<sub>17</sub>H<sub>19</sub>F<sub>3</sub>N<sub>4</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	S	0	0
			29	17	3	4	4	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	F	N	O	S	0	0
			29	17	3	4	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	97	Total	O	0	0
			97	97		
5	B	77	Total	O	0	0
			77	77		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable decaprenylphosphoryl-beta-D-ribose oxidase



- Molecule 1: Probable decaprenylphosphoryl-beta-D-ribose oxidase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.52Å 84.13Å 90.15Å 90.00° 100.73° 90.00°	Depositor
Resolution (Å)	48.02 – 2.30 48.02 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.4 (48.02-2.30) 98.4 (48.02-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.43 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.209 , 0.236 0.209 , 0.236	Depositor DCC
$R_{free}$ test set	1928 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.9	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 31.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 38415 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6779	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IMD, FAD, 2R2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/3371	0.51	1/4573 (0.0%)
1	B	0.44	2/3212 (0.1%)	0.51	0/4356
All	All	0.44	2/6583 (0.0%)	0.51	1/8929 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	172	TRP	CD2-CE2	5.09	1.47	1.41
1	B	230	TRP	CD2-CE2	5.04	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	414	LEU	CA-CB-CG	5.83	128.70	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3290	0	3277	16	0
1	B	3141	0	3128	30	0
2	A	53	0	31	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	53	0	31	0	0
3	A	5	0	5	0	0
3	B	5	0	5	0	0
4	A	29	0	19	4	0
4	B	29	0	19	2	0
5	A	97	0	0	0	0
5	B	77	0	0	0	0
All	All	6779	0	6515	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:ARG:HH11	1:B:159:ARG:HG2	1.17	1.08
1:A:159:ARG:HH11	1:A:159:ARG:HG2	1.33	0.90
1:A:159:ARG:CG	1:A:159:ARG:HH11	1.97	0.77
1:B:442:ARG:HG2	1:B:442:ARG:HH11	1.51	0.75
1:A:159:ARG:NH1	1:A:159:ARG:HG2	1.97	0.72
1:A:93:ASP:O	1:A:96:VAL:HG13	1.98	0.63
2:A:501:FAD:HM72	4:A:503:2R2:H11	1.82	0.60
1:B:314:TYR:O	1:B:315:HIS:HB3	2.01	0.60
1:A:392:ILE:HA	1:A:396:LEU:HD12	1.84	0.59
4:A:503:2R2:OBB	4:A:503:2R2:H2	2.01	0.59
1:B:363:LEU:HD13	4:B:502:2R2:H15	1.84	0.59
1:B:392:ILE:HA	1:B:396:LEU:HD12	1.85	0.58
1:B:442:ARG:CG	1:B:442:ARG:HH11	2.18	0.56
1:B:159:ARG:CG	1:B:159:ARG:HH11	2.03	0.55
1:A:240:LEU:HD22	1:A:355:GLN:HG2	1.89	0.54
1:A:99:ASP:OD1	1:A:119:ARG:NH1	2.41	0.54
1:B:83:ILE:HG12	1:B:90:VAL:HG12	1.91	0.53
1:A:92:ILE:HG13	1:A:123:VAL:HG21	1.91	0.53
1:B:159:ARG:NH1	1:B:159:ARG:HG2	1.97	0.53
1:B:231:PHE:HB3	1:B:364:ASN:HB2	1.92	0.52
1:A:206:THR:HB	1:A:211:GLU:HG3	1.91	0.52
1:A:83:ILE:HG12	1:A:90:VAL:HG12	1.92	0.52
1:B:103:LYS:HE2	1:B:267:PHE:CE2	2.47	0.49
1:B:209:LEU:HG	1:B:241:GLY:HA3	1.93	0.49
1:B:240:LEU:HD22	1:B:355:GLN:HG2	1.94	0.49
1:A:375:ALA:HB3	1:A:378:SER:HB2	1.95	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:254:GLU:HG2	1:B:255:GLN:N	2.29	0.48
1:B:99:ASP:HB2	1:B:119:ARG:HB3	1.97	0.47
1:B:346:GLU:HG2	1:B:410:PHE:CD1	2.50	0.45
1:B:116:PRO:HG2	1:B:118:THR:O	2.17	0.45
1:A:363:LEU:HD13	4:A:503:2R2:H16	1.99	0.45
1:B:134:LYS:HG3	1:B:228:SER:HB3	1.99	0.45
1:A:51:ALA:O	1:A:450:PHE:HA	2.17	0.44
1:B:88:LYS:HD2	1:B:193:PRO:HD3	2.00	0.44
1:B:375:ALA:HB3	1:B:378:SER:HB2	2.00	0.44
1:A:317:LEU:HD21	4:A:503:2R2:H1	2.00	0.44
1:B:442:ARG:NH1	1:B:442:ARG:CG	2.80	0.43
1:B:137:HIS:HA	1:B:370:GLY:HA3	2.00	0.43
1:B:29:THR:HG21	1:B:34:MET:HB3	2.00	0.43
1:B:159:ARG:CG	1:B:159:ARG:NH1	2.71	0.42
1:B:448:ARG:HG2	1:B:461:LEU:HD23	2.02	0.42
1:B:99:ASP:OD1	1:B:119:ARG:NH1	2.53	0.42
1:B:93:ASP:O	1:B:96:VAL:HG13	2.19	0.42
1:B:49:ARG:HA	1:B:49:ARG:HD2	1.73	0.42
1:B:47:GLY:HA3	1:B:48:GLY:HA2	1.65	0.41
1:A:456:ARG:HG2	1:A:461:LEU:HD22	2.02	0.41
4:B:502:2R2:H16	4:B:502:2R2:H8	1.79	0.41
1:B:371:PRO:HA	1:B:382:PRO:HG3	2.03	0.40
1:A:441[B]:ARG:NH2	1:A:450:PHE:O	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/480 (88%)	415 (98%)	8 (2%)	0	100	100
1	B	405/480 (84%)	401 (99%)	4 (1%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	828/960 (86%)	816 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/384 (89%)	327 (96%)	15 (4%)	35	46
1	B	328/384 (85%)	317 (97%)	11 (3%)	44	59
All	All	670/768 (87%)	644 (96%)	26 (4%)	39	53

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	THR
1	A	92	ILE
1	A	159	ARG
1	A	292	ILE
1	A	296	TRP
1	A	298	ARG
1	A	317	LEU
1	A	342	GLU
1	A	355	GLN
1	A	367	LYS
1	A	396	LEU
1	A	406	ARG
1	A	413	ARG
1	A	414	LEU
1	A	421	ARG
1	B	44	GLU
1	B	92	ILE
1	B	96	VAL
1	B	103	LYS
1	B	159	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	209	LEU
1	B	253	VAL
1	B	254	GLU
1	B	396	LEU
1	B	413	ARG
1	B	448	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	A	501	-	52,58,58	1.41	7 (13%)	52,89,89	2.20	10 (19%)
3	IMD	A	502	-	3,5,5	0.40	0	4,5,5	0.55	0
4	2R2	A	503	-	28,31,31	2.54	5 (17%)	31,46,46	1.62	5 (16%)
2	FAD	B	501	-	52,58,58	1.40	7 (13%)	52,89,89	2.19	10 (19%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	2R2	B	502	-	28,31,31	2.50	4 (14%)	31,46,46	1.55	4 (12%)
3	IMD	B	503	-	3,5,5	0.39	0	4,5,5	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	501	-	-	0/30/50/50	0/6/6/6
3	IMD	A	502	-	-	0/0/0/0	0/1/1/1
4	2R2	A	503	-	-	0/17/32/32	0/3/3/3
2	FAD	B	501	-	-	0/30/50/50	0/6/6/6
4	2R2	B	502	-	-	0/17/32/32	1/3/3/3
3	IMD	B	503	-	-	0/0/0/0	0/1/1/1

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	2R2	CAZ-CAY	2.02	1.49	1.42
2	A	501	FAD	C10-N10	2.04	1.41	1.39
2	B	501	FAD	C10-N10	2.12	1.41	1.39
4	B	502	2R2	CAI-CAY	2.14	1.49	1.42
4	A	503	2R2	CAI-CAY	2.19	1.49	1.42
2	A	501	FAD	C9A-N10	2.33	1.42	1.38
2	B	501	FAD	C9A-N10	2.40	1.42	1.38
2	B	501	FAD	C5A-C4A	3.08	1.47	1.40
2	A	501	FAD	C5A-C4A	3.13	1.47	1.40
2	B	501	FAD	C8-C7	3.31	1.49	1.41
2	A	501	FAD	C8-C7	3.32	1.49	1.41
2	B	501	FAD	C9A-C5X	3.62	1.50	1.42
2	A	501	FAD	C9A-C5X	3.64	1.50	1.42
4	A	503	2R2	CAK-NAJ	3.71	1.36	1.31
2	B	501	FAD	C4-C4X	3.80	1.49	1.41
2	A	501	FAD	C4-C4X	3.92	1.49	1.41
4	B	502	2R2	CAK-NAJ	3.97	1.37	1.31
2	A	501	FAD	C4X-C10	4.27	1.48	1.40
2	B	501	FAD	C4X-C10	4.30	1.48	1.40
4	A	503	2R2	CAC-CAB	5.65	1.50	1.40
4	B	502	2R2	CAC-CAB	5.70	1.50	1.40
4	B	502	2R2	CAZ-CAB	10.49	1.50	1.39
4	A	503	2R2	CAZ-CAB	10.82	1.50	1.39

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	FAD	N3A-C2A-N1A	-7.49	122.98	128.87
2	A	501	FAD	N3A-C2A-N1A	-7.33	123.11	128.87
2	A	501	FAD	C4-C4X-C10	-5.94	116.14	119.94
2	B	501	FAD	C4-C4X-C10	-5.73	116.27	119.94
2	B	501	FAD	C4X-C4-N3	-3.68	118.71	123.52
2	A	501	FAD	C4X-C4-N3	-3.65	118.75	123.52
2	A	501	FAD	N3-C2-N1	-3.19	122.33	127.69
2	B	501	FAD	N3-C2-N1	-3.09	122.49	127.69
4	A	503	2R2	CAU-CAV-NAM	-2.79	104.57	110.44
2	A	501	FAD	C1B-N9A-C4A	-2.44	124.08	126.81
4	A	503	2R2	CAO-CAN-NAM	-2.42	105.34	110.44
2	B	501	FAD	C1B-N9A-C4A	-2.39	124.14	126.81
4	B	502	2R2	FAE-CAD-CAC	-2.20	108.93	112.67
2	A	501	FAD	C5X-C9A-N10	2.10	119.16	117.58
2	B	501	FAD	C5X-C9A-N10	2.18	119.21	117.58
2	B	501	FAD	C1'-N10-C9A	2.38	121.59	118.83
4	A	503	2R2	CAC-CAH-CAI	2.40	123.27	119.66
2	A	501	FAD	C1'-N10-C9A	2.41	121.62	118.83
4	B	502	2R2	CAC-CAH-CAI	2.51	123.43	119.66
4	B	502	2R2	CAV-NAM-CAN	2.54	117.21	112.57
4	A	503	2R2	CAU-NAP-CAO	2.56	114.59	108.87
2	B	501	FAD	C4-C4X-N5	3.16	122.54	118.70
2	A	501	FAD	C4-C4X-N5	3.36	122.78	118.70
2	A	501	FAD	C4X-N5-C5X	3.66	121.04	116.72
2	B	501	FAD	C4X-N5-C5X	3.70	121.08	116.72
4	A	503	2R2	CAK-NAJ-CAI	5.48	114.58	103.83
4	B	502	2R2	CAK-NAJ-CAI	5.60	114.81	103.83
2	B	501	FAD	C4-N3-C2	8.54	122.28	115.16
2	A	501	FAD	C4-N3-C2	8.66	122.38	115.16

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	502	2R2	CAN-CAO-CAU-CAV-NAM-NAP

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FAD	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	2R2	4	0
4	B	502	2R2	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	427/480 (88%)	0.18	17 (3%) 42 51	20, 31, 63, 77	0
1	B	410/480 (85%)	0.16	14 (3%) 49 58	23, 32, 51, 65	0
All	All	837/960 (87%)	0.17	31 (3%) 45 54	20, 31, 56, 77	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	326	ALA	7.5
1	B	46	GLY	5.2
1	B	267	PHE	4.8
1	B	47	GLY	4.7
1	A	296	TRP	4.4
1	A	46	GLY	4.3
1	A	327	TYR	3.9
1	B	315	HIS	3.7
1	A	297	TYR	3.7
1	A	47	GLY	3.6
1	B	230	TRP	3.5
1	A	298	ARG	3.3
1	A	461	LEU	3.2
1	B	394	ASP	3.0
1	B	260	LEU	2.9
1	B	363	LEU	2.7
1	A	318	ASP	2.7
1	A	329	PRO	2.6
1	B	37	LYS	2.5
1	B	258	ALA	2.4
1	B	360	TYR	2.4
1	B	300	SER	2.3
1	A	362	PHE	2.3
1	B	362	PHE	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	12	ARG	2.2
1	A	316	PRO	2.2
1	A	7	THR	2.2
1	A	292	ILE	2.2
1	A	317	LEU	2.1
1	A	319	MET	2.1
1	B	7	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	IMD	B	503	5/5	0.94	0.33	3.87	43,44,44,44	0
4	2R2	B	502	29/29	0.85	0.26	2.28	48,50,54,55	0
4	2R2	A	503	29/29	0.87	0.21	1.13	45,46,49,50	0
2	FAD	A	501	53/53	0.96	0.14	0.10	23,24,26,27	0
2	FAD	B	501	53/53	0.97	0.13	-0.02	24,25,26,26	0
3	IMD	A	502	5/5	0.98	0.18	-	37,37,37,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.