



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:57 PM GMT

PDB ID : 4PFE  
Title : Crystal structure of vsfGFP-0  
Authors : Jauch, R.; Chen, S.L.  
Deposited on : 2014-04-29  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	 82% 16% •
1	B	343	 86% 12% ••

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5555 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	343	Total	C	N	O	S	0	0	0
			2715	1707	476	522	10			
1	B	341	Total	C	N	O	S	0	0	0
			2699	1698	472	519	10			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	ARG	SER	engineered mutation	UNP P42212
A	39	ASN	TYR	engineered mutation	UNP P42212
A	64	LEU	PHE	engineered mutation	UNP P42212
A	66	CRO	SER	chromophore	UNP P42212
A	66	CRO	TYR	chromophore	UNP P42212
A	66	CRO	GLY	chromophore	UNP P42212
A	80	ARG	GLN	engineered mutation	UNP P42212
A	99	SER	PHE	engineered mutation	UNP P42212
A	105	THR	ASN	engineered mutation	UNP P42212
A	145	PHE	TYR	engineered mutation	UNP P42212
A	153	THR	MET	engineered mutation	UNP P42212
A	163	ALA	VAL	engineered mutation	UNP P42212
A	171	VAL	ILE	engineered mutation	UNP P42212
A	206	VAL	ALA	engineered mutation	UNP P42212
B	30	ARG	SER	engineered mutation	UNP P42212
B	39	ASN	TYR	engineered mutation	UNP P42212
B	64	LEU	PHE	engineered mutation	UNP P42212
B	66	CRO	SER	chromophore	UNP P42212
B	66	CRO	TYR	chromophore	UNP P42212
B	66	CRO	GLY	chromophore	UNP P42212
B	80	ARG	GLN	engineered mutation	UNP P42212
B	99	SER	PHE	engineered mutation	UNP P42212
B	105	THR	ASN	engineered mutation	UNP P42212
B	145	PHE	TYR	engineered mutation	UNP P42212
B	153	THR	MET	engineered mutation	UNP P42212

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	163	ALA	VAL	engineered mutation	UNP P42212
B	171	VAL	ILE	engineered mutation	UNP P42212
B	206	VAL	ALA	engineered mutation	UNP P42212


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	72	Total O 72 72	0	0
2	B	69	Total O 69 69	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein



- Molecule 1: Green fluorescent protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.43Å 83.43Å 228.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.16 – 2.60 47.16 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.16-2.60) 94.5 (47.16-2.60)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.01 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.199 , 0.255 0.208 , 0.256	Depositor DCC
$R_{free}$ test set	1308 reflections (5.68%)	DCC
Wilson B-factor (Å <sup>2</sup> )	58.0	Xtriage
Anisotropy	0.394	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 42.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 25648 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5555	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/2753	0.49	0/3723
1	B	0.25	0/2736	0.48	0/3700
All	All	0.25	0/5489	0.49	0/7423

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2715	0	2622	34	0
1	B	2699	0	2610	29	0
2	A	72	0	0	2	0
2	B	69	0	0	2	0
All	All	5555	0	5232	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:SER:HA	1:A:5:GLU:HG2	1.60	0.82
1:B:77:HIS:O	1:B:80:ARG:NH1	2.16	0.79
1:B:232:VAL:HG21	1:B:330:VAL:HG21	1.72	0.71
1:B:163:ALA:HB3	1:B:183:GLN:HB3	1.73	0.71
1:B:321:THR:HG23	1:B:342:THR:HA	1.75	0.68
1:A:231:GLN:N	1:A:231:GLN:OE1	2.28	0.66
1:A:306:ARG:HB2	1:A:308:THR:HG22	1.79	0.65
1:A:323:VAL:HG22	1:A:340:GLN:HG2	1.81	0.62
1:A:163:ALA:HB3	1:A:183:GLN:HB3	1.82	0.60
1:A:313:MET:HB3	1:A:316:LEU:HD21	1.82	0.60
1:B:330:VAL:C	1:B:332:PHE:H	2.07	0.58
1:A:203:THR:HG22	1:A:224:VAL:HG13	1.86	0.56
1:A:213:GLU:OE1	2:A:401:HOH:O	2.18	0.55
1:A:17:GLU:OE1	1:A:122:ARG:NH1	2.41	0.54
1:B:71:PHE:HE2	1:B:119:LEU:HD22	1.74	0.52
1:B:143:TYR:CZ	1:B:209:LYS:HE3	2.44	0.52
1:B:19:ASP:OD1	2:B:464:HOH:O	2.18	0.52
1:A:98:ILE:HB	1:A:106:TYR:HB2	1.91	0.52
1:A:42:LEU:HB2	1:A:222:GLU:HB3	1.93	0.51
1:B:74:TYR:O	1:B:79:LYS:NZ	2.43	0.51
1:B:262:TYR:CE1	1:B:330:VAL:HG12	2.45	0.51
1:A:155:ASP:HB2	1:A:162:LYS:HG3	1.93	0.50
1:A:69:GLN:O	1:A:72:SER:OG	2.21	0.50
1:B:313:MET:HB3	1:B:316:LEU:HD21	1.94	0.49
1:A:22:VAL:HG12	1:A:27:PHE:HE1	1.76	0.49
1:A:73:ARG:NE	2:A:440:HOH:O	2.39	0.49
1:B:71:PHE:CE2	1:B:119:LEU:HD22	2.48	0.48
1:A:321:THR:HG23	1:A:342:THR:HA	1.95	0.48
1:B:190:ASP:OD1	1:B:190:ASP:N	2.46	0.48
1:A:143:TYR:CZ	1:A:209:LYS:HE3	2.49	0.47
1:B:250:LEU:HG	1:B:313:MET:HE1	1.96	0.47
1:A:206:VAL:HG13	1:A:221:LEU:HB3	1.95	0.47
1:B:209:LYS:HE2	1:B:217:HIS:O	2.14	0.47
1:A:209:LYS:HE2	1:A:217:HIS:O	2.14	0.47
1:A:330:VAL:C	1:A:332:PHE:H	2.18	0.47
1:B:155:ASP:HB2	1:B:162:LYS:HG3	1.96	0.47
1:A:190:ASP:OD1	1:A:190:ASP:N	2.47	0.46
1:B:259:VAL:HG13	1:B:302:ARG:HD2	1.97	0.46
1:A:277:TRP:CZ2	1:B:175:SER:HA	2.50	0.46
1:A:204:GLN:HG3	1:A:223:PHE:HB2	1.97	0.46
1:B:203:THR:HG22	1:B:224:VAL:HG13	1.98	0.46
1:A:81:HIS:O	1:A:196:PRO:HB3	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:302:ARG:NH1	2:B:450:HOH:O	2.48	0.46
1:B:262:TYR:CZ	1:B:330:VAL:HG12	2.51	0.45
1:B:306:ARG:HB2	1:B:308:THR:HG22	1.98	0.45
1:B:242:VAL:HG11	1:B:316:LEU:HD12	1.99	0.45
1:A:259:VAL:HG13	1:A:302:ARG:HD2	1.99	0.45
1:A:242:VAL:HG11	1:A:316:LEU:HD12	1.99	0.45
1:A:294:VAL:HB	1:A:298:PHE:CG	2.52	0.44
1:B:69:GLN:O	1:B:72:SER:OG	2.29	0.44
1:A:66:CRO:HD1	1:A:66:CRO:N2	2.33	0.44
1:A:231:GLN:NE2	1:A:334:TYR:OH	2.52	0.43
1:B:155:ASP:N	1:B:160:GLY:O	2.40	0.43
1:B:330:VAL:HG23	1:B:330:VAL:O	2.19	0.42
1:A:175:SER:HA	1:B:277:TRP:CZ2	2.54	0.42
1:B:77:HIS:HB2	1:B:231:GLN:NE2	2.33	0.42
1:A:71:PHE:HE2	1:A:119:LEU:HD22	1.84	0.42
1:A:71:PHE:CE2	1:A:119:LEU:HD22	2.55	0.42
1:B:98:ILE:HB	1:B:106:TYR:HB2	2.01	0.41
1:A:12:VAL:HG13	1:A:71:PHE:HE1	1.86	0.41
1:A:313:MET:HE2	1:A:316:LEU:HD21	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/343 (98%)	333 (98%)	5 (2%)	0	100	100
1	B	336/343 (98%)	330 (98%)	6 (2%)	0	100	100
All	All	674/686 (98%)	663 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/293 (100%)	275 (94%)	18 (6%)	23	46
1	B	291/293 (99%)	279 (96%)	12 (4%)	37	66
All	All	584/586 (100%)	554 (95%)	30 (5%)	29	55

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	THR
1	A	12	VAL
1	A	22	VAL
1	A	72	SER
1	A	80	ARG
1	A	99	SER
1	A	118	THR
1	A	119	LEU
1	A	164	ASN
1	A	190	ASP
1	A	204	GLN
1	A	225	THR
1	A	248	LEU
1	A	259	VAL
1	A	278	VAL
1	A	288	SER
1	A	306	ARG
1	A	341	VAL
1	B	72	SER
1	B	119	LEU
1	B	126	LYS
1	B	164	ASN
1	B	190	ASP
1	B	204	GLN
1	B	225	THR
1	B	248	LEU
1	B	259	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	293	SER
1	B	306	ARG
1	B	341	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	77	HIS
1	B	231	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CRO	A	66	1	23,23,24	4.14	5 (21%)	30,32,34	4.95	8 (26%)
1	CRO	B	66	1	23,23,24	4.11	5 (21%)	30,32,34	5.01	8 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	66	1	-	0/12/31/32	0/2/2/2
1	CRO	B	66	1	-	0/12/31/32	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRO	C2-N3	-2.96	1.32	1.39
1	B	66	CRO	C2-N3	-2.91	1.32	1.39
1	A	66	CRO	CG2-CB2	2.71	1.52	1.46
1	A	66	CRO	O2-C2	2.72	1.28	1.23
1	B	66	CRO	O2-C2	2.72	1.28	1.23
1	B	66	CRO	CG2-CB2	2.75	1.52	1.46
1	B	66	CRO	C1-N2	3.81	1.38	1.32
1	A	66	CRO	C1-N2	3.90	1.38	1.32
1	B	66	CRO	CB2-CA2	18.57	1.52	1.35
1	A	66	CRO	CB2-CA2	18.63	1.52	1.35

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CRO	O2-C2-CA2	-12.79	123.62	130.97
1	B	66	CRO	O2-C2-CA2	-12.60	123.73	130.97
1	B	66	CRO	C2-CA2-N2	-10.16	101.29	109.03
1	A	66	CRO	C2-CA2-N2	-10.04	101.39	109.03
1	B	66	CRO	CG2-CB2-CA2	-9.42	118.34	130.27
1	A	66	CRO	CG2-CB2-CA2	-8.38	119.67	130.27
1	A	66	CRO	CA1-C1-N3	-2.20	122.31	124.98
1	B	66	CRO	CA1-C1-N2	2.24	126.92	123.71
1	A	66	CRO	CA1-C1-N2	2.25	126.93	123.71
1	B	66	CRO	CB2-CA2-N2	2.40	132.97	128.71
1	B	66	CRO	CB2-CA2-C2	2.48	125.73	122.24
1	A	66	CRO	CB2-CA2-N2	2.87	133.81	128.71
1	A	66	CRO	CA2-N2-C1	5.22	109.93	105.54
1	B	66	CRO	CA2-N2-C1	5.63	110.27	105.54
1	B	66	CRO	CA2-C2-N3	18.35	112.45	103.37
1	A	66	CRO	CA2-C2-N3	18.41	112.48	103.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66	CRO	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	342/343 (99%)	-0.26	0 100 100	24, 50, 81, 107	0
1	B	340/343 (99%)	-0.25	0 100 100	30, 51, 84, 112	0
All	All	682/686 (99%)	-0.25	0 100 100	24, 51, 82, 112	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CRO	B	66	22/23	0.91	0.20	-	36,52,63,65	0
1	CRO	A	66	22/23	0.95	0.15	-	22,33,48,63	0

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.