



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 22, 2016 – 07:38 AM EDT

PDB ID : 4PFP  
Title : Myosin VI motor domain in the Pi release state (with Pi) space group P21  
Authors : Isabet, T.; Benisty, H.; Llinas, P.; Sweeney, H.L.; Houdusse, A.  
Deposited on : 2014-04-30  
Resolution : 2.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0122  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4425 (2.34-2.30)
Clashscore	102246	5057 (2.34-2.30)
Ramachandran outliers	100387	5008 (2.34-2.30)
Sidechain outliers	100360	5007 (2.34-2.30)
RSRZ outliers	91569	4432 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	788	<div> <div>10%</div> <div>88%</div> <div>8% . .</div> </div>
1	C	788	<div> <div>5%</div> <div>91%</div> <div>6% .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PGO	A	803	-	-	-	X
5	PO4	A	804	-	-	-	X
5	PO4	A	805	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12526 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Unconventional myosin-VI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	760	Total	C	N	O	S	0	6	0
			5988	3822	1032	1105	29			
1	C	766	Total	C	N	O	S	0	4	0
			6051	3854	1035	1132	30			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is S-1,2-PROPANEDIOL (three-letter code: PGO) (formula:  $C_3H_8O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			5	3	2		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula:  $O_4P$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O P 5 4 1	0	0
5	A	1	Total O P 5 4 1	0	0
5	C	1	Total O P 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	170	Total O 170 170	0	0
6	C	241	Total O 241 241	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Unconventional myosin-VI



#### • Molecule 1: Unconventional myosin-VI



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.65Å 94.89Å 97.30Å 90.00° 99.55° 90.00°	Depositor
Resolution (Å)	23.39 – 2.32 43.98 – 2.32	Depositor EDS
% Data completeness (in resolution range)	99.8 (23.39-2.32) 99.8 (43.98-2.32)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 2.32Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, $R_{free}$	0.189 , 0.229 0.199 , 0.239	Depositor DCC
$R_{free}$ test set	3902 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.2	Xtriage
Anisotropy	0.526	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 42.5	EDS
Estimated twinning fraction	0.016 for l,-k,h	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 78099 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12526	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.74 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.6209e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PGO, PO4, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	3/6121 (0.0%)	0.67	4/8270 (0.0%)
1	C	0.52	0/6175	0.66	1/8340 (0.0%)
All	All	0.54	3/12296 (0.0%)	0.67	5/16610 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	392	THR	C-N	-17.98	0.92	1.34
1	A	393	ARG	C-N	8.50	1.53	1.34
1	A	5	LYS	C-N	-7.92	1.19	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	LYS	O-C-N	-8.81	104.36	121.10
1	A	5	LYS	CA-C-N	8.22	140.12	117.10
1	A	25	GLY	C-N-CD	5.99	140.97	128.40
1	A	90	VAL	N-CA-C	-5.11	97.20	111.00
1	C	90	VAL	N-CA-C	-5.08	97.30	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	5	LYS	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5988	0	5795	43	0
1	C	6051	0	5862	25	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
3	A	27	0	12	0	0
3	C	27	0	12	0	0
4	A	5	0	8	0	0
5	A	10	0	0	0	0
5	C	5	0	0	0	0
6	A	170	0	0	2	0
6	C	241	0	0	3	0
All	All	12526	0	11689	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:GLY:O	1:A:358:THR:HG23	1.78	0.83
1:C:68:LEU:HD12	1:C:702:GLN:HG2	1.64	0.79
1:C:723:PRO:HD2	1:C:726:LEU:HD12	1.67	0.77
1:A:299:GLU:CD	6:A:1019:HOH:O	2.26	0.73
1:C:457:ILE:HD13	1:C:480:LEU:HD12	1.69	0.72
1:A:356:GLY:C	1:A:358:THR:HG23	2.09	0.71
1:A:357:SER:HA	1:A:358:THR:C	2.10	0.71
1:A:357:SER:HA	1:A:358:THR:HG23	1.77	0.65
1:A:358:THR:OG1	1:A:359:SER:N	2.33	0.60
1:A:357:SER:CA	1:A:358:THR:HG23	2.33	0.58
1:A:449:SER:C	1:A:450:TYR:HA	2.23	0.58
1:C:592:GLN:HA	6:C:926:HOH:O	2.04	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:ASP:O	1:A:181:ARG:HG3	2.04	0.57
1:A:357:SER:CA	1:A:358:THR:CG2	2.85	0.54
1:A:739:PHE:CE2	1:A:756:VAL:HG21	2.45	0.52
1:C:512:GLN:HA	1:C:515:ILE:HD12	1.91	0.52
1:A:73:LEU:HD22	1:A:698:LEU:HD21	1.92	0.52
1:A:464:GLU:HG2	1:A:465:HIS:CD2	2.45	0.51
1:C:515:ILE:HD11	6:C:1093:HOH:O	2.10	0.51
1:A:357:SER:HA	1:A:358:THR:CG2	2.41	0.50
1:A:394:VAL:HG12	1:A:604:SER:HB3	1.92	0.50
1:C:739:PHE:CE2	1:C:756:VAL:HG21	2.47	0.50
1:A:489:LEU:HD21	1:A:697:VAL:HG12	1.94	0.49
1:C:127:VAL:HG11	1:C:160:ASN:OD1	2.12	0.49
1:A:87:TYR:HB3	1:A:94:LEU:HD11	1.95	0.48
1:C:716:ASN:O	1:C:719:LYS:HG2	2.13	0.48
1:A:551:HIS:HB2	1:A:557:LEU:HD22	1.95	0.48
1:A:393:ARG:HB2	1:A:411:LEU:HD21	1.96	0.48
1:A:705:PHE:HB3	1:A:757:PHE:HB3	1.95	0.48
1:C:87:TYR:HB3	1:C:94:LEU:HD11	1.95	0.48
1:C:516:ASP:O	1:C:520:ALA:HB2	2.16	0.46
1:C:739:PHE:HE2	1:C:756:VAL:HG21	1.80	0.46
1:C:11:HIS:HD2	1:C:14:ASP:H	1.64	0.46
1:A:134:ALA:O	1:A:145:GLN:HG3	2.15	0.46
1:C:134:ALA:O	1:C:145:GLN:HG3	2.16	0.46
1:C:268:PRO:HB2	1:C:279:THR:HB	1.98	0.45
1:C:68:LEU:CD1	1:C:702:GLN:HG2	2.40	0.45
1:C:166:ARG:O	1:C:170:GLU:HB2	2.18	0.44
1:A:166:ARG:O	1:A:170:GLU:HB2	2.18	0.44
1:A:449:SER:C	1:A:450:TYR:CA	2.86	0.44
1:A:357:SER:CA	1:A:358:THR:C	2.85	0.44
1:A:517:LEU:O	1:A:525:ILE:HG13	2.17	0.44
1:A:268:PRO:HB2	1:A:279:THR:HB	2.00	0.44
1:A:357:SER:N	1:A:358:THR:HG23	2.32	0.43
1:A:520:ALA:O	1:A:524:GLY:HA3	2.18	0.43
1:A:521:ARG:O	1:A:523:VAL:N	2.48	0.43
1:C:647:PHE:CE2	1:C:651:LEU:HD11	2.53	0.43
1:A:723:PRO:HD2	1:A:726:LEU:HD12	2.00	0.43
1:A:771:LYS:HB3	1:A:776:HIS:CD2	2.54	0.43
1:C:705:PHE:HB3	1:C:757:PHE:HB3	1.99	0.43
1:C:233:SER:HA	1:C:425:LYS:HE2	2.00	0.43
1:A:490:LYS:O	1:A:494:GLU:HG3	2.20	0.42
1:A:742:LEU:HD22	1:A:783:ARG:HH11	1.83	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:742:LEU:HD22	1:C:783:ARG:HH11	1.84	0.42
1:A:229:LEU:HD11	1:A:647:PHE:CE2	2.54	0.42
1:A:742:LEU:HD12	1:A:769:ILE:HD11	2.01	0.42
1:A:181:ARG:HG2	1:A:327:ILE:HG22	2.02	0.42
1:C:519:GLU:HB2	1:C:648:LYS:HE3	2.01	0.42
1:C:29:LEU:HG	1:C:45:ILE:HG12	2.00	0.42
1:A:462:TYR:CD2	1:A:586:VAL:HG22	2.54	0.42
1:A:759:ARG:HD3	6:A:1009:HOH:O	2.18	0.42
1:A:475:TYR:OH	1:A:648:LYS:HG3	2.21	0.41
1:A:357:SER:C	1:A:358:THR:HG22	2.41	0.41
1:A:233:SER:HA	1:A:425:LYS:HE2	2.02	0.41
1:A:739:PHE:HE2	1:A:756:VAL:HG21	1.84	0.41
1:C:719:LYS:HA	1:C:722:MET:HG2	2.01	0.41
1:C:479:LYS:HA	6:C:1093:HOH:O	2.21	0.41
1:A:356:GLY:C	1:A:358:THR:CG2	2.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	758/788 (96%)	745 (98%)	13 (2%)	0	100	100
1	C	764/788 (97%)	751 (98%)	13 (2%)	0	100	100
All	All	1522/1576 (97%)	1496 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	630/696 (90%)	618 (98%)	12 (2%)	65	80
1	C	644/696 (92%)	635 (99%)	9 (1%)	74	86
All	All	1274/1392 (92%)	1253 (98%)	21 (2%)	68	84

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	SER
1	A	330	ASP
1	A	357	SER
1	A	359	SER
1	A	462	TYR
1	A	464	GLU
1	A	523	VAL
1	A	599	ASP
1	A	639	SER
1	A	708	ARG
1	A	732	ARG
1	A	733	LEU
1	C	179	ASP
1	C	180	ASP
1	C	359	SER
1	C	406	VAL
1	C	543	PHE
1	C	599	ASP
1	C	614	ASP
1	C	708	ARG
1	C	733	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	186	ASN
1	A	507	HIS
1	A	716	ASN
1	C	11	HIS
1	C	482	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	ADP	A	802	2	24,29,29	0.67	0	23,45,45	0.91	1 (4%)
4	PGO	A	803	-	4,4,4	0.70	0	2,4,4	2.16	1 (50%)
5	PO4	A	804	2	4,4,4	1.93	2 (50%)	6,6,6	0.26	0
5	PO4	A	805	-	4,4,4	1.80	2 (50%)	6,6,6	0.25	0
3	ADP	C	801	2	24,29,29	0.66	0	23,45,45	0.64	0
5	PO4	C	802	2	4,4,4	1.82	3 (75%)	6,6,6	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	802	2	-	0/12/32/32	0/3/3/3
4	PGO	A	803	-	-	0/2/2/2	0/0/0/0
5	PO4	A	804	2	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PO4	A	805	-	-	0/0/0/0	0/0/0/0
3	ADP	C	801	2	-	0/12/32/32	0/3/3/3
5	PO4	C	802	2	-	0/0/0/0	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	802	PO4	P-O1	2.02	1.60	1.53
5	C	802	PO4	P-O2	2.05	1.60	1.53
5	A	805	PO4	P-O4	2.12	1.60	1.53
5	A	805	PO4	P-O3	2.15	1.60	1.53
5	C	802	PO4	P-O4	2.21	1.60	1.53
5	A	804	PO4	P-O4	2.28	1.60	1.53
5	A	804	PO4	P-O2	2.28	1.60	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	ADP	O3'-C3'-C4'	2.08	117.21	111.01
4	A	803	PGO	O1-C1-C2	3.04	117.77	110.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	3

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	449:SER	C	450:TYR	N	2.84
1	A	5:LYS	C	6:PRO	N	1.19
1	A	392:THR	C	393:ARG	N	0.92



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	760/788 (96%)	0.58	82 (10%) 8 12	29, 57, 86, 114	1 (0%)
1	C	766/788 (97%)	0.36	38 (4%) 32 41	26, 48, 75, 110	0
All	All	1526/1576 (96%)	0.47	120 (7%) 15 22	26, 53, 83, 114	1 (0%)

All (120) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	177	ASP	7.9
1	C	178	ILE	7.6
1	A	517	LEU	7.3
1	A	177	ASP	6.3
1	A	757	PHE	5.1
1	A	522	LEU	4.9
1	A	572	ARG	4.8
1	C	789	ILE	4.8
1	A	788	LEU	4.8
1	C	38	GLY	4.7
1	A	610	CYS	4.6
1	A	729	LEU	4.4
1	A	545	SER	4.3
1	A	756	VAL	4.2
1	A	301	LEU	4.2
1	A	579	ILE	4.2
1	A	786	HIS	4.1
1	A	576	GLY	4.1
1	A	577	PHE	4.1
1	A	781	VAL	4.1
1	C	753	LEU	4.0
1	C	667[A]	ARG	4.0
1	A	557	LEU	3.9
1	A	560	PRO	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	310	LEU	3.9
1	A	751	PHE	3.9
1	A	753	LEU	3.9
1	A	733	LEU	3.8
1	C	24	ILE	3.7
1	A	558	SER	3.7
1	A	300	TYR	3.6
1	A	639	SER	3.5
1	A	393	ARG	3.5
1	C	757	PHE	3.4
1	C	547	VAL	3.4
1	C	509	VAL	3.3
1	C	405	THR	3.3
1	A	281	TYR	3.3
1	A	561	ARG	3.3
1	A	547	VAL	3.2
1	C	179	ASP	3.2
1	C	691[A]	CYS	3.2
1	A	29	LEU	3.2
1	A	726	LEU	3.2
1	A	5	LYS	3.2
1	C	571	ILE	3.1
1	A	35	ASN	3.1
1	A	304	GLY	3.1
1	A	292	LEU	3.0
1	C	35	ASN	3.0
1	A	559	ILE	3.0
1	A	176	GLN	3.0
1	A	580	ARG	3.0
1	A	291	ILE	2.9
1	C	557	LEU	2.9
1	A	39	LYS	2.9
1	A	714	LEU	2.9
1	C	786	HIS	2.8
1	A	306	LEU	2.8
1	A	523	VAL	2.7
1	A	571	ILE	2.7
1	C	480	LEU	2.7
1	A	718	TYR	2.7
1	A	520	ALA	2.6
1	C	517	LEU	2.6
1	C	726	LEU	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	528	ILE	2.6
1	C	462	TYR	2.6
1	A	568	HIS	2.6
1	A	22	VAL	2.6
1	C	570	ASN	2.6
1	C	22	VAL	2.6
1	C	692	SER	2.5
1	C	176	GLN	2.5
1	A	735	CYS	2.5
1	A	305	SER	2.5
1	A	732	ARG	2.5
1	A	311	LEU	2.4
1	A	178	ILE	2.4
1	A	149	VAL	2.4
1	A	525	ILE	2.4
1	A	541	GLN	2.4
1	A	581	HIS	2.4
1	C	300	TYR	2.4
1	A	182	ILE	2.3
1	A	282	PHE	2.3
1	A	549	GLN	2.3
1	C	543	PHE	2.3
1	A	45	ILE	2.3
1	C	639	SER	2.3
1	A	394	VAL	2.2
1	A	566	ALA	2.2
1	A	38	GLY	2.2
1	A	666	ILE	2.2
1	A	578	ILE	2.2
1	A	268	PRO	2.2
1	A	602	HIS	2.1
1	A	780	LEU	2.1
1	A	395	MET	2.1
1	C	636	GLY	2.1
1	C	31	ILE	2.1
1	A	731	PRO	2.1
1	A	48	VAL	2.1
1	A	784	VAL	2.1
1	A	789	ILE	2.1
1	C	21	ILE	2.1
1	C	149	VAL	2.1
1	A	518	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	335	LEU	2.1
1	A	697	VAL	2.0
1	A	462	TYR	2.0
1	C	289	LYS	2.0
1	A	570	ASN	2.0
1	C	558	SER	2.0
1	C	607	SER	2.0
1	A	548	HIS	2.0
1	C	551	HIS	2.0
1	C	756	VAL	2.0
1	A	711	PHE	2.0
1	A	569	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	PO4	A	804	5/5	0.90	0.27	4.39	18,29,33,36	5
5	PO4	A	805	5/5	0.90	0.32	3.79	79,80,84,86	0
4	PGO	A	803	5/5	0.85	0.27	2.49	68,69,71,71	0
3	ADP	A	802	27/27	0.98	0.14	-0.36	24,34,47,49	0
2	MG	A	801	1/1	0.97	0.15	-0.39	29,29,29,29	0
3	ADP	C	801	27/27	0.98	0.14	-0.45	25,29,36,37	0
2	MG	C	803	1/1	0.97	0.15	-0.71	31,31,31,31	0
5	PO4	C	802	5/5	0.99	0.13	-1.34	32,36,38,41	0

## 6.5 Other polymers

There are no such residues in this entry.