



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:37 PM GMT

PDB ID : 4PGL  
Title : Crystal structure of engineered D-tagatose 3-epimerase PcDTE-ILS6  
Authors : Hee, C.S.; Bosshart, A.; Schirmer, T.  
Deposited on : 2014-05-02  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	<div> <div>4%</div> <div>95%</div> <div>5%</div> </div>
1	B	298	<div> <div>%</div> <div>93%</div> <div>• •</div> </div>
1	C	298	<div> <div>2%</div> <div>95%</div> <div>5%</div> </div>
1	D	298	<div> <div>2%</div> <div>94%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SOL	B	504[A]	-	-	-	X
4	SOE	A	503	-	-	-	X
4	SOE	B	502	-	-	-	X
4	SOE	D	502	-	-	-	X
5	LTG	A	505[B]	-	-	-	X
5	LTG	B	503[B]	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-tagatose 3-epimerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	4	0
			2402	1521	423	438	20			
1	B	290	Total	C	N	O	S	0	1	0
			2304	1460	399	427	18			
1	C	298	Total	C	N	O	S	0	1	0
			2375	1504	416	437	18			
1	D	290	Total	C	N	O	S	0	1	0
			2301	1459	396	427	19			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	SER	THR	engineered mutation	UNP O50580
A	39	SER	GLY	engineered mutation	UNP O50580
A	109	ASN	THR	engineered mutation	UNP O50580
A	116	HIS	SER	engineered mutation	UNP O50580
A	122	VAL	LYS	engineered mutation	UNP O50580
A	153	ALA	VAL	engineered mutation	UNP O50580
A	157	TYR	PHE	engineered mutation	UNP O50580
A	183	HIS	GLN	engineered mutation	UNP O50580
A	194	ASN	THR	engineered mutation	UNP O50580
A	215	GLN	ALA	engineered mutation	UNP O50580
A	245	ILE	MET	engineered mutation	UNP O50580
A	251	THR	LYS	engineered mutation	UNP O50580
A	260	CYS	GLY	engineered mutation	UNP O50580
A	265	LEU	MET	engineered mutation	UNP O50580
A	291	LEU	-	expression tag	UNP O50580
A	292	GLU	-	expression tag	UNP O50580
A	293	HIS	-	expression tag	UNP O50580
A	294	HIS	-	expression tag	UNP O50580
A	295	HIS	-	expression tag	UNP O50580
A	296	HIS	-	expression tag	UNP O50580
A	297	HIS	-	expression tag	UNP O50580

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	298	HIS	-	expression tag	UNP O50580
B	9	SER	THR	engineered mutation	UNP O50580
B	39	SER	GLY	engineered mutation	UNP O50580
B	109	ASN	THR	engineered mutation	UNP O50580
B	116	HIS	SER	engineered mutation	UNP O50580
B	122	VAL	LYS	engineered mutation	UNP O50580
B	153	ALA	VAL	engineered mutation	UNP O50580
B	157	TYR	PHE	engineered mutation	UNP O50580
B	183	HIS	GLN	engineered mutation	UNP O50580
B	194	ASN	THR	engineered mutation	UNP O50580
B	215	GLN	ALA	engineered mutation	UNP O50580
B	245	ILE	MET	engineered mutation	UNP O50580
B	251	THR	LYS	engineered mutation	UNP O50580
B	260	CYS	GLY	engineered mutation	UNP O50580
B	265	LEU	MET	engineered mutation	UNP O50580
B	291	LEU	-	expression tag	UNP O50580
B	292	GLU	-	expression tag	UNP O50580
B	293	HIS	-	expression tag	UNP O50580
B	294	HIS	-	expression tag	UNP O50580
B	295	HIS	-	expression tag	UNP O50580
B	296	HIS	-	expression tag	UNP O50580
B	297	HIS	-	expression tag	UNP O50580
B	298	HIS	-	expression tag	UNP O50580
C	9	SER	THR	engineered mutation	UNP O50580
C	39	SER	GLY	engineered mutation	UNP O50580
C	109	ASN	THR	engineered mutation	UNP O50580
C	116	HIS	SER	engineered mutation	UNP O50580
C	122	VAL	LYS	engineered mutation	UNP O50580
C	153	ALA	VAL	engineered mutation	UNP O50580
C	157	TYR	PHE	engineered mutation	UNP O50580
C	183	HIS	GLN	engineered mutation	UNP O50580
C	194	ASN	THR	engineered mutation	UNP O50580
C	215	GLN	ALA	engineered mutation	UNP O50580
C	245	ILE	MET	engineered mutation	UNP O50580
C	251	THR	LYS	engineered mutation	UNP O50580
C	260	CYS	GLY	engineered mutation	UNP O50580
C	265	LEU	MET	engineered mutation	UNP O50580
C	291	LEU	-	expression tag	UNP O50580
C	292	GLU	-	expression tag	UNP O50580
C	293	HIS	-	expression tag	UNP O50580
C	294	HIS	-	expression tag	UNP O50580
C	295	HIS	-	expression tag	UNP O50580

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	296	HIS	-	expression tag	UNP O50580
C	297	HIS	-	expression tag	UNP O50580
C	298	HIS	-	expression tag	UNP O50580
D	9	SER	THR	engineered mutation	UNP O50580
D	39	SER	GLY	engineered mutation	UNP O50580
D	109	ASN	THR	engineered mutation	UNP O50580
D	116	HIS	SER	engineered mutation	UNP O50580
D	122	VAL	LYS	engineered mutation	UNP O50580
D	153	ALA	VAL	engineered mutation	UNP O50580
D	157	TYR	PHE	engineered mutation	UNP O50580
D	183	HIS	GLN	engineered mutation	UNP O50580
D	194	ASN	THR	engineered mutation	UNP O50580
D	215	GLN	ALA	engineered mutation	UNP O50580
D	245	ILE	MET	engineered mutation	UNP O50580
D	251	THR	LYS	engineered mutation	UNP O50580
D	260	CYS	GLY	engineered mutation	UNP O50580
D	265	LEU	MET	engineered mutation	UNP O50580
D	291	LEU	-	expression tag	UNP O50580
D	292	GLU	-	expression tag	UNP O50580
D	293	HIS	-	expression tag	UNP O50580
D	294	HIS	-	expression tag	UNP O50580
D	295	HIS	-	expression tag	UNP O50580
D	296	HIS	-	expression tag	UNP O50580
D	297	HIS	-	expression tag	UNP O50580
D	298	HIS	-	expression tag	UNP O50580

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mn 1 1	0	0
2	A	2	Total Mn 2 2	0	0
2	D	1	Total Mn 1 1	0	0
2	C	1	Total Mn 1 1	0	0

- Molecule 3 is SUGAR (L-sorbose) (three-letter code: SOL) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	1
			12	6	6		
3	B	1	Total	C	O	0	1
			12	6	6		
3	C	1	Total	C	O	0	1
			12	6	6		
3	D	1	Total	C	O	0	1
			12	6	6		

- Molecule 4 is SUGAR (alpha-L-sorbypyranose) (three-letter code: SOE) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 12 6 6	0	0
4	A	1	Total C O 12 6 6	0	0
4	B	1	Total C O 12 6 6	0	0
4	C	1	Total C O 12 6 6	0	0
4	D	1	Total C O 12 6 6	0	0

- Molecule 5 is L-tagatose (three-letter code: LTG) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 12 6 6	0	1
5	B	1	Total C O 12 6 6	0	1
5	C	1	Total C O 12 6 6	0	1
5	D	1	Total C O 12 6 6	0	1

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	82	Total O 85 85	0	3

*Continued on next page...*




*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	91	Total 95	O 95	0	4
6	C	96	Total 100	O 100	0	4
6	D	98	Total 101	O 101	0	3

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-tagatose 3-epimerase




- Molecule 1: D-tagatose 3-epimerase



- Molecule 1: D-tagatose 3-epimerase



- Molecule 1: D-tagatose 3-epimerase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.81Å 47.44Å 126.38Å 90.00° 102.49° 90.00°	Depositor
Resolution (Å)	36.85 – 2.10 36.85 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.7 (36.85-2.10) 98.8 (36.85-2.10)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.83 (at 2.10Å)	Xtriage
Refinement program	COOT	Depositor
R, $R_{free}$	0.152 , 0.181 0.163 , 0.187	Depositor DCC
$R_{free}$ test set	3506 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.5	Xtriage
Anisotropy	0.144	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 53.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 69447 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9924	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 27.30 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.2562e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LTG, MN, SOE, SOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/2470	0.71	1/3336 (0.0%)
1	B	0.59	0/2360	0.71	0/3189
1	C	0.59	0/2437	0.71	2/3295 (0.1%)
1	D	0.58	0/2357	0.71	2/3185 (0.1%)
All	All	0.59	0/9624	0.71	5/13005 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	50	ARG	NE-CZ-NH2	5.52	123.06	120.30
1	A	164	ASP	CB-CG-OD1	5.51	123.26	118.30
1	C	92	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	D	224	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	D	164	ASP	CB-CG-OD1	5.06	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2402	0	2341	11	0
1	B	2304	0	2252	12	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2375	0	2300	8	0
1	D	2301	0	2248	7	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	12	0	0	1	0
3	B	12	0	0	0	0
3	C	12	0	0	0	0
3	D	12	0	0	0	0
4	A	24	0	24	1	0
4	B	12	0	12	1	0
4	C	12	0	12	1	0
4	D	12	0	12	1	0
5	A	12	0	11	1	0
5	B	12	0	12	1	0
5	C	12	0	12	2	0
5	D	12	0	12	3	0
6	A	85	0	0	0	0
6	B	95	0	0	1	0
6	C	100	0	0	3	0
6	D	101	0	0	1	0
All	All	9924	0	9248	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:504:SOE:H6	1:B:121:MET:HE1	1.69	0.74
1:B:115:GLN:HE21	1:B:160:TRP:HE1	1.36	0.74
1:B:263[B]:ARG:NH1	6:B:640:HOH:O	2.19	0.74
4:D:502:SOE:HO1	4:D:502:SOE:HO3	1.32	0.73
5:C:303[B]:LTG:H3	5:C:303[B]:LTG:H12	1.36	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/298 (101%)	295 (98%)	5 (2%)	0	100	100
1	B	289/298 (97%)	286 (99%)	3 (1%)	0	100	100
1	C	297/298 (100%)	293 (99%)	4 (1%)	0	100	100
1	D	289/298 (97%)	286 (99%)	3 (1%)	0	100	100
All	All	1175/1192 (99%)	1160 (99%)	15 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/250 (102%)	249 (98%)	5 (2%)	63	68
1	B	243/250 (97%)	242 (100%)	1 (0%)	93	96
1	C	250/250 (100%)	247 (99%)	3 (1%)	78	84
1	D	243/250 (97%)	243 (100%)	0	100	100
All	All	990/1000 (99%)	981 (99%)	9 (1%)	84	89

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	276	GLU
1	C	267	ASN
1	C	122	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	122	VAL
1	B	99	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	115	GLN
1	B	116	HIS
1	C	283	GLN
1	B	2	ASN
1	C	298	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 5 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SOL	A	502[A]	2	11,11,11	0.44	0	10,14,14	1.69	2 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SOE	A	503	-	12,12,12	1.48	1 (8%)	16,18,18	3.12	8 (50%)
4	SOE	A	504	-	12,12,12	0.75	0	16,18,18	1.77	2 (12%)
5	LTG	A	505[B]	2	11,11,11	0.58	0	10,14,14	1.59	2 (20%)
4	SOE	B	502	-	12,12,12	1.11	0	16,18,18	3.17	10 (62%)
5	LTG	B	503[B]	2	11,11,11	0.50	0	10,14,14	1.23	1 (10%)
3	SOL	B	504[A]	2	11,11,11	0.39	0	10,14,14	1.23	1 (10%)
4	SOE	C	302	-	12,12,12	1.50	3 (25%)	16,18,18	1.75	5 (31%)
5	LTG	C	303[B]	2	11,11,11	0.46	0	10,14,14	1.48	3 (30%)
3	SOL	C	304[A]	2	11,11,11	0.46	0	10,14,14	1.53	2 (20%)
4	SOE	D	502	-	12,12,12	1.29	1 (8%)	16,18,18	3.72	10 (62%)
5	LTG	D	503[B]	2	11,11,11	0.56	0	10,14,14	1.01	0
3	SOL	D	504[A]	2	11,11,11	0.39	0	10,14,14	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SOL	A	502[A]	2	-	0/16/16/16	0/0/0/0
4	SOE	A	503	-	-	0/3/23/23	0/1/1/1
4	SOE	A	504	-	-	0/3/23/23	0/1/1/1
5	LTG	A	505[B]	2	-	0/16/16/16	0/0/0/0
4	SOE	B	502	-	-	0/3/23/23	0/1/1/1
5	LTG	B	503[B]	2	-	0/16/16/16	0/0/0/0
3	SOL	B	504[A]	2	-	0/16/16/16	0/0/0/0
4	SOE	C	302	-	-	0/3/23/23	0/1/1/1
5	LTG	C	303[B]	2	-	0/16/16/16	0/0/0/0
3	SOL	C	304[A]	2	-	0/16/16/16	0/0/0/0
4	SOE	D	502	-	-	0/3/23/23	0/1/1/1
5	LTG	D	503[B]	2	-	0/16/16/16	0/0/0/0
3	SOL	D	504[A]	2	-	0/16/16/16	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	302	SOE	C2-C3	-2.25	1.51	1.54
4	C	302	SOE	O2-C2	2.15	1.44	1.41
4	D	502	SOE	O2-C2	3.03	1.46	1.41
4	C	302	SOE	O6-C2	3.44	1.45	1.42

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	SOE	O6-C2	3.96	1.46	1.42

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	502	SOE	C5-C4-C3	-10.09	99.15	110.50
4	A	503	SOE	C5-C4-C3	-7.03	102.59	110.50
4	D	502	SOE	O6-C2-C3	-5.80	103.06	109.87
4	B	502	SOE	O3-C3-C2	-5.76	101.53	110.11
4	C	302	SOE	O2-C2-C1	-3.71	105.32	111.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502[A]	SOL	1	0
4	A	504	SOE	1	0
5	A	505[B]	LTG	1	0
4	B	502	SOE	1	0
5	B	503[B]	LTG	1	0
4	C	302	SOE	1	0
5	C	303[B]	LTG	2	0
4	D	502	SOE	1	0
5	D	503[B]	LTG	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	298/298 (100%)	-0.07	13 (4%) 38 47	12, 21, 40, 69	0
1	B	290/298 (97%)	-0.07	4 (1%) 78 82	13, 22, 39, 64	0
1	C	298/298 (100%)	-0.13	7 (2%) 64 70	11, 19, 37, 60	0
1	D	290/298 (97%)	-0.11	5 (1%) 73 78	13, 22, 37, 64	0
All	All	1176/1192 (98%)	-0.10	29 (2%) 61 67	11, 21, 38, 69	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	122	VAL	7.2
1	D	1	MET	7.0
1	C	298	HIS	5.2
1	A	1	MET	4.6
1	B	1	MET	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	SOE	A	503	12/12	0.79	0.27	16.13	38,57,66,68	0
4	SOE	B	502	12/12	0.79	0.26	10.52	32,47,49,59	0
4	SOE	D	502	12/12	0.88	0.17	10.50	27,38,48,48	0
5	LTG	B	503[B]	12/12	0.94	0.23	3.70	27,33,34,37	12
3	SOL	B	504[A]	12/12	0.94	0.22	3.03	37,50,61,70	12
5	LTG	A	505[B]	12/12	0.92	0.20	2.96	16,21,27,27	12
3	SOL	A	502[A]	12/12	0.95	0.17	1.99	29,49,61,63	12
5	LTG	C	303[B]	12/12	0.94	0.16	1.58	20,26,31,32	12
3	SOL	C	304[A]	12/12	0.94	0.16	1.53	25,35,49,50	12
5	LTG	D	503[B]	12/12	0.95	0.18	1.52	25,32,36,37	12
3	SOL	D	504[A]	12/12	0.94	0.18	1.41	28,44,58,64	12
4	SOE	C	302	12/12	0.92	0.18	0.62	24,32,39,41	0
4	SOE	A	504	12/12	0.92	0.15	0.44	24,29,34,35	0
2	MN	A	506	1/1	0.99	0.09	-0.27	13,13,13,13	0
2	MN	B	501	1/1	0.99	0.07	-3.08	27,27,27,27	0
2	MN	D	501	1/1	1.00	0.07	-3.18	26,26,26,26	0
2	MN	C	301	1/1	1.00	0.07	-4.76	21,21,21,21	0
2	MN	A	501	1/1	1.00	0.07	-4.85	22,22,22,22	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.