



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:35 AM GMT

PDB ID : 3PHO
Title : Crystal structure of S64-4 in complex with PSBP
Authors : Evans, D.W.; Evans, S.V.
Deposited on : 2010-11-04
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	<div> <div></div> <div>84%</div> <div>13%</div> <div>•</div> </div>
2	B	222	<div> <div>3%</div> <div>81%</div> <div>16%</div> <div>•</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called S64-4 Fab (IgG1) light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	S	0	1	0
			1673	1038	289	339	7			

- Molecule 2 is a protein called S64-4 Fab (IgG1) heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	216	Total	C	N	O	S	0	0	0
			1636	1042	269	317	8			

- Molecule 3 is 3-DEOXY-ALPHA-D-MANNO-OCT-2-ULOPYRANONOSYL-(2->8)-3-DEOXY-ALPHA-D-MANNO-OCT-2-ULOPYRANONOSYL-(2->4)-3-DEOXY-ALPHA-D-MANNO-OCT-2-ULOPYRANONOSYL-(2->6)-2-AMINO-2-DEOXY-4-O-PHOSPHONO-BETA-D-GLUCOPYRANOSE (three-letter code: KD1) (formula: C₃₀H₅₀NO₂₉P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			61	30	1	29	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	90	Total 90	O 90	0	0
4	B	56	Total 56	O 56	0	0

- Molecule 1: S64-4 Fab (IgG1) light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	163.36 Å 163.36 Å 43.04 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.96 – 2.60 19.96 – 2.60	Depositor EDS
% Data completeness (in resolution range)	94.7 (19.96-2.60) 94.7 (19.96-2.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 2.59 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.219 , 0.275 0.217 , 0.268	Depositor DCC
R_{free} test set	993 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	39.8	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.2	EDS
Estimated twinning fraction	0.012 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Outliers	1 of 19490 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3516	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KD1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	1/1713 (0.1%)	0.68	1/2324 (0.0%)
2	B	0.54	0/1680	0.64	0/2296
All	All	0.54	1/3393 (0.0%)	0.66	1/4620 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	137	CYS	CB-SG	-5.43	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	184	LEU	CA-CB-CG	5.35	127.60	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1673	0	1611	30	0
2	B	1636	0	1598	33	0
3	A	61	0	45	3	0
4	A	90	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	56	0	0	4	0
All	All	3516	0	3254	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:SER:OG	1:A:8:PRO:HD2	1.37	1.22
1:A:7:SER:HB3	1:A:22:SER:HB2	1.56	0.88
1:A:7:SER:CB	1:A:8:PRO:CD	2.51	0.87
1:A:7:SER:OG	1:A:8:PRO:CD	2.22	0.84
2:B:52:ARG:HH11	2:B:56:THR:HG23	1.45	0.81
2:B:71:ARG:NH1	4:B:270:HOH:O	2.01	0.79
1:A:7:SER:CB	1:A:8:PRO:HD2	2.13	0.76
2:B:82:MET:HE3	2:B:90:TYR:CE2	2.22	0.75
2:B:52:ARG:HD2	2:B:56:THR:CG2	2.18	0.73
2:B:151:GLU:HG3	2:B:152:PRO:HA	1.69	0.72
1:A:7:SER:HB2	1:A:8:PRO:CD	2.19	0.71
2:B:33:TYR:HA	4:B:270:HOH:O	1.94	0.68
2:B:52:ARG:HD2	2:B:56:THR:HG23	1.77	0.67
1:A:6:GLN:HE22	1:A:91:TYR:HA	1.59	0.67
3:A:218:KD1:O1	2:B:33:TYR:OH	2.08	0.66
1:A:109:ILE:H	1:A:169:GLN:HE22	1.43	0.66
1:A:7:SER:HB2	1:A:8:PRO:HD3	1.77	0.66
2:B:71:ARG:HE	2:B:73:ASN:HD21	1.42	0.66
1:A:8:PRO:O	1:A:9:ALA:HB3	1.98	0.64
2:B:151:GLU:CG	2:B:152:PRO:HA	2.27	0.64
1:A:2:ILE:N	4:A:276:HOH:O	2.30	0.63
2:B:17:SER:HB2	2:B:82(A):ASN:HD22	1.64	0.63
1:A:185:THR:OG1	1:A:188:GLU:HG2	2.01	0.61
1:A:161:GLY:O	1:A:183:THR:HG22	2.00	0.61
2:B:71:ARG:NE	2:B:73:ASN:HD21	1.98	0.60
2:B:71:ARG:HE	2:B:73:ASN:ND2	2.02	0.57
2:B:1:GLU:HG3	4:B:245:HOH:O	2.05	0.57
1:A:94:HIS:HD2	4:A:291:HOH:O	1.88	0.56
1:A:193:ASN:HD21	1:A:215:ASN:H	1.52	0.55
2:B:82:MET:CE	2:B:90:TYR:CE2	2.89	0.55
2:B:33:TYR:HB2	2:B:95:ASP:HB3	1.89	0.55
1:A:6:GLN:NE2	1:A:104:GLY:H	2.07	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:150:PRO:O	2:B:202:HIS:HE1	1.93	0.52
1:A:163:LEU:HD21	2:B:174:GLN:HG3	1.92	0.52
1:A:94:HIS:O	1:A:99:ARG:HA	2.10	0.52
2:B:151:GLU:HG3	2:B:152:PRO:CA	2.38	0.51
1:A:183:THR:HG21	2:B:174:GLN:HE22	1.76	0.51
1:A:94:HIS:HE1	4:A:269:HOH:O	1.95	0.50
1:A:7:SER:HG	1:A:8:PRO:HD2	1.67	0.49
2:B:151:GLU:OE1	2:B:152:PRO:HA	2.12	0.48
1:A:193:ASN:ND2	1:A:215:ASN:H	2.11	0.48
1:A:188:GLU:O	1:A:192:HIS:HD2	1.96	0.48
1:A:8:PRO:O	1:A:9:ALA:CB	2.62	0.47
2:B:38:ARG:HB3	2:B:48:LEU:HD11	1.96	0.46
2:B:186:VAL:HB	2:B:187:PRO:HD2	1.98	0.46
1:A:108:GLU:OE2	1:A:145:LYS:NZ	2.45	0.45
2:B:29:PHE:CD2	2:B:76:SER:HA	2.52	0.45
2:B:33:TYR:CB	2:B:95:ASP:HB3	2.46	0.44
2:B:126:PRO:HD3	2:B:211:LYS:HD2	1.99	0.44
2:B:155:VAL:HG22	2:B:200:VAL:HG22	1.99	0.44
2:B:32:TYR:CD1	2:B:97:GLY:HA3	2.54	0.43
1:A:44:PRO:O	4:A:307:HOH:O	2.20	0.43
2:B:151:GLU:CG	4:B:240:HOH:O	2.68	0.42
3:A:218:KD1:O24	3:A:218:KD1:O28	2.30	0.42
1:A:139:LEU:CD1	1:A:199:ALA:HB2	2.49	0.42
2:B:17:SER:CB	2:B:82(A):ASN:HD22	2.33	0.42
1:A:119:SER:O	1:A:137:CYS:HB2	2.20	0.42
1:A:41:GLN:HB2	1:A:51:LEU:HD11	2.02	0.41
2:B:17:SER:HB2	2:B:82(A):ASN:ND2	2.33	0.41
2:B:55:TYR:CD1	2:B:71:ARG:HD3	2.56	0.40
1:A:18:ARG:NH1	4:A:232:HOH:O	2.54	0.40
3:A:218:KD1:O1	3:A:218:KD1:H16	2.20	0.40
2:B:157:TRP:CZ3	2:B:198:CYS:HB3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/217 (99%)	206 (96%)	8 (4%)	1 (0%)	34	60
2	B	212/222 (96%)	201 (95%)	10 (5%)	1 (0%)	34	60
All	All	427/439 (97%)	407 (95%)	18 (4%)	2 (0%)	34	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	7	SER
2	B	129	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/191 (100%)	184 (96%)	7 (4%)	41	69
2	B	184/188 (98%)	179 (97%)	5 (3%)	52	79
All	All	375/379 (99%)	363 (97%)	12 (3%)	46	74

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ILE
1	A	84	GLU
1	A	119	SER
1	A	139	LEU
1	A	145	LYS
1	A	158	ARG
1	A	211	SER
2	B	35	SER
2	B	82	MET
2	B	82(C)	LEU
2	B	139	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	183	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	46	GLN
1	A	78	ASN
1	A	94	HIS
1	A	127	GLN
1	A	140	ASN
1	A	164	ASN
1	A	169	GLN
1	A	192	HIS
1	A	193	ASN
2	B	73	ASN
2	B	82(A)	ASN
2	B	174	GLN
2	B	202	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	KD1	A	218	-	51,64,64	1.15	6 (11%)	63,99,99	1.70	10 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KD1	A	218	-	-	0/36/130/130	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	218	KD1	C27-C26	-2.87	1.49	1.53
3	A	218	KD1	O21-C18	-2.71	1.38	1.42
3	A	218	KD1	O20-C18	-2.36	1.36	1.40
3	A	218	KD1	O6-C16	-2.27	1.37	1.42
3	A	218	KD1	P1-O28	-2.19	1.46	1.54
3	A	218	KD1	O6-C2	-2.08	1.36	1.40

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	218	KD1	C11-C12-C13	-6.87	104.52	110.84
3	A	218	KD1	C30-O20-C18	-4.07	108.33	116.31
3	A	218	KD1	O6-C16-C15	-3.55	99.82	107.78
3	A	218	KD1	C19-C20-C21	-3.16	106.10	111.37
3	A	218	KD1	O24-C27-C26	-2.79	105.59	110.31
3	A	218	KD1	C27-C26-N1	-2.45	106.31	110.86
3	A	218	KD1	O20-C30-C29	-2.15	102.79	107.77
3	A	218	KD1	P1-O25-C28	2.21	126.86	121.56
3	A	218	KD1	C16-C15-C14	2.34	116.67	111.41
3	A	218	KD1	O21-C22-C21	3.45	113.06	108.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	218	KD1	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	216/217 (99%)	-0.21	3 (1%) 78 74	17, 27, 40, 55	0
2	B	216/222 (97%)	0.16	7 (3%) 51 44	20, 37, 49, 53	0
All	All	432/439 (98%)	-0.02	10 (2%) 64 57	17, 31, 47, 55	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	217	CYS	3.6
2	B	130	GLY	3.4
1	A	215	ASN	3.3
2	B	1	GLU	2.8
2	B	206	SER	2.7
2	B	65	GLY	2.5
2	B	175	SER	2.3
2	B	163	SER	2.3
2	B	3	MET	2.2
1	A	160	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	KD1	A	218	61/61	0.92	0.17	0.30	23,28,44,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.