



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:32 PM GMT

PDB ID : 1PIO  
Title : AN ENGINEERED STAPHYLOCOCCUS AUREUS PC1 BETA-LACTAMASE THAT HYDROLYSES THIRD GENERATION CEPHALOSPORINS  
Authors : Zawadzke, L.E.; Herzberg, O.  
Deposited on : 1995-10-11  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	257	
1	B	257	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4153 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-LACTAMASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	0	0
			2022	1287	340	392	3			
1	B	256	Total	C	N	O	S	0	0	0
			2022	1287	340	392	3			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	238	SER	ALA	ENGINEERED	UNP P00807
A	?	-	ILE	ENGINEERED DELETION	UNP P00807
B	238	SER	ALA	ENGINEERED	UNP P00807
B	?	-	ILE	ENGINEERED DELETION	UNP P00807

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	53	Total	O	0	0
			53	53		
2	B	56	Total	O	0	0
			56	56		



Note EDS was not executed.

Chain B:

Amino Acid	Category
MET	Grey
A125	Green
K31	Green
E32	Green
L33	Green
I32	Green
D35	Green
Y40	Green
N41	Green
A42	Green
H43	Green
I44	Green
G45	Green
V46	Green
Y47	Green
A48	Green
L49	Green
D50	Green
F60	Green
H61	Green
S62	Green
F66	Green
A67	Green
Y68	Green
A69	Green
K73	Green
L80	Green
H84	Green
P87	Green
L89	Green
Y88	Green
H89	Green
K90	Green
K93	Green
K94	Green
V95	Green
D101	Green
A103	Green
L109	Green
Y105	Green
S106	Green
P107	Green
E110	Green
K111	Green
Y112	Green
V113	Green
G114	Green
K115	Green
D116	Green
L119	Green
I123	Green
E124	Green
L125	Green
K126	Green
L127	Green
K128	Green
S129	Green
S130	Green
D131	Green
K132	Green
T133	Green
A134	Green
H135	Green
H136	Green
K137	Green
I138	Green
T139	Green
A140	Green
L142	Green
K146	Green
K147	Green
V148	Green
K149	Green
L155	Green
K158	Green
P162	Green
V163	Green
I167	Green
E168	Green
L169	Green
P187	Green
N170	Green
K177	Green
K178	Green
T182	Green
P183	Green
G187	Green
A188	Green
T189	Green
L190	Green
N191	Green
K192	Green
A195	Green
N196	Green
G197	Green
K198	Green
A199	Green
K204	Green
K205	Green
D206	Green
L207	Green
L208	Green
D209	Green
E210	Green
L212	Green
K215	Green
S216	Green
G217	Green
K222	Green
V225	Green
P226	Green
K227	Green
D228	Green
V231	Green
A232	Green
D233	Green
Q237	Green
S238	Green
T240	Green
Y241	Green
A242	Green
S243	Green
K244	Green
A248	Green
Y251	Green
P252	Green
K253	Green
G254	Green
Q255	Green
S256	Green
E257	Green
P258	Green
L259	Green
L261	Green
L262	Green
V262	Green
T265	Green
K266	Green
K267	Green
S271	Green
D272	Green
N275	Green
D276	Green
K277	Green
L278	Green
L279	Green
A283	Green
N286	Green
V287	Green
L290	Green
E290	Green

## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.00 Å 67.60 Å 141.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.80	Depositor
% Data completeness (in resolution range)	86.5 (8.00-2.80)	Depositor
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.189 , 0.189	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4153	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/2052	0.82	2/2754 (0.1%)
1	B	0.63	0/2052	0.84	2/2754 (0.1%)
All	All	0.60	0/4104	0.83	4/5508 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	LEU	CA-CB-CG	5.96	129.02	115.30
1	B	198	LYS	N-CA-C	5.24	125.15	111.00
1	A	166	GLU	N-CA-C	-5.24	96.86	111.00
1	B	45	GLY	N-CA-C	-5.22	100.04	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2022	0	2104	99	0
1	B	2022	0	2104	87	0
2	A	53	0	0	1	0
2	B	56	0	0	4	0
All	All	4153	0	4208	182	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 182 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:ALA:HB3	1:A:267:LYS:HB2	1.47	0.94
1:A:146:LYS:HD2	1:A:146:LYS:H	1.37	0.88
1:A:124:GLU:HG2	1:A:210:LEU:HD21	1.54	0.86
1:B:146:LYS:HD2	1:B:146:LYS:H	1.43	0.81
1:A:80:LEU:HD23	1:A:123:ILE:HD11	1.66	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/257 (99%)	226 (89%)	27 (11%)	1 (0%)	39	74
1	B	254/257 (99%)	230 (91%)	21 (8%)	3 (1%)	16	47
All	All	508/514 (99%)	456 (90%)	48 (9%)	4 (1%)	24	58

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	168	GLU
1	B	198	LYS
1	B	113	VAL
1	A	107	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/228 (100%)	212 (93%)	15 (7%)	21	51
1	B	227/228 (100%)	213 (94%)	14 (6%)	23	54
All	All	454/456 (100%)	425 (94%)	29 (6%)	22	52

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	256	SER
1	B	33	LEU
1	B	191	ASN
1	A	284	LYS
1	B	62	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	161	ASN
1	B	41	ASN
1	B	132	ASN
1	A	135	ASN
1	B	89	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.