



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:40 PM GMT

PDB ID : 4PLC
Title : Crystal structure of ancestral apicomplexan lactate dehydrogenase with malate.
Authors : Boucher, J.I.; Jacobowitz, J.R.; Beckett, B.C.; Classen, S.; Theobald, D.L.
Deposited on : 2014-05-16
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>7%</div> <div>93%</div> <div>.</div> <div>.</div> </div>
1	B	334	<div> <div>18%</div> <div>93%</div> <div>.</div> <div>.</div> </div>
1	C	334	<div> <div>9%</div> <div>94%</div> <div>.</div> <div>.</div> </div>
1	D	334	<div> <div>17%</div> <div>93%</div> <div>.</div> <div>.</div> </div>

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 22070 atoms, of which 10501 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called lactate dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	324	Total	C	H	N	O	S	0	9	0
			5154	1603	2624	430	480	17			
1	B	324	Total	C	H	N	O	S	0	3	0
			5077	1580	2587	424	470	16			
1	C	324	Total	C	H	N	O	S	0	4	0
			5085	1583	2590	425	471	16			
1	D	324	Total	C	H	N	O	S	0	3	0
			5070	1579	2582	423	470	16			

- Molecule 2 is PYRUVIC ACID (three-letter code: PYR) (formula: C₃H₄O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			9	3	3	3		
2	B	1	Total	C	H	O	0	0
			9	3	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	H	O	0	0
			9	3	3	3		
2	D	1	Total	C	H	O	0	0
			9	3	3	3		

- Molecule 3 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula: $C_{21}H_{29}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	0	0
			71	21	27	7	14	2		
3	C	1	Total	C	H	N	O	P	0	0
			71	21	27	7	14	2		

- Molecule 4 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	B	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		
4	D	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	387	Total	O	0	0
			387	387		
5	B	303	Total	O	0	0
			303	303		
5	C	361	Total	O	0	0
			361	361		
5	D	315	Total	O	0	0
			315	315		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: lactate dehydrogenase




- Molecule 1: lactate dehydrogenase




- Molecule 1: lactate dehydrogenase



- Molecule 1: lactate dehydrogenase





4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	134.37Å 134.37Å 159.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.95 – 1.50 46.95 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.95-1.50) 99.7 (46.95-1.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 1.50Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.124 , 0.152 0.127 , 0.155	Depositor DCC
R_{free} test set	2016 reflections (0.79%)	DCC
Wilson B-factor (Å ²)	17.3	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 53.8	EDS
Estimated twinning fraction	0.500 for h,-h-k,-l 0.487 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.500 for h,-h-k,-l	Depositor
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.21$	Xtriage
Outliers	0 of 258762 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	22070	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PYR, NAI, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/2575	0.45	0/3467
1	B	0.27	0/2535	0.44	0/3411
1	C	0.28	0/2540	0.44	0/3418
1	D	0.27	0/2533	0.45	0/3410
All	All	0.28	0/10183	0.45	0/13706

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2530	2624	2613	10	0
1	B	2490	2587	2584	10	0
1	C	2495	2590	2585	7	0
1	D	2488	2582	2580	9	0
2	A	6	3	3	0	0
2	B	6	3	3	0	0
2	C	6	3	3	1	0
2	D	6	3	3	0	0
3	A	44	27	27	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	44	27	27	1	0
4	B	44	26	26	0	0
4	D	44	26	26	0	0
5	A	387	0	0	6	2
5	B	303	0	0	9	1
5	C	361	0	0	3	0
5	D	315	0	0	5	3
All	All	11569	10501	10480	35	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:MET:SD	1:C:290:ASN:ND2	2.49	0.85
1:C:64[A]:LYS:NZ	5:C:801:HOH:O	2.23	0.69
1:D:313:GLU:OE2	1:D:316:ARG:NH2	2.25	0.68
1:B:85:GLY:O	5:B:706:HOH:O	2.13	0.65
1:D:104:PRO:O	5:D:577:HOH:O	2.13	0.65
1:B:223:GLU:OE1	5:B:643:HOH:O	2.14	0.64
1:C:238:GLN:NE2	5:C:807:HOH:O	2.30	0.64
1:B:328:GLU:OE2	5:B:591:HOH:O	2.16	0.60
1:D:29:GLN:NE2	5:D:596:HOH:O	2.31	0.59
1:B:313:GLU:OE2	5:B:501:HOH:O	2.16	0.58
1:C:317:LYS:NZ	5:C:501:HOH:O	2.36	0.58
1:A:320:LYS:NZ	5:A:831:HOH:O	2.37	0.58
1:C:303:GLU:N	1:C:303:GLU:OE2	2.35	0.58
1:A:119:LYS:NZ	5:A:795:HOH:O	2.36	0.58
1:B:254:GLU:OE2	5:B:624:HOH:O	2.17	0.56
1:B:64:LYS:NZ	5:B:542:HOH:O	2.37	0.56
1:D:223:GLU:OE1	5:D:598:HOH:O	2.18	0.55
1:B:320:LYS:NZ	5:B:662:HOH:O	2.39	0.55
1:A:242:TYR:OH	5:A:754:HOH:O	2.20	0.49
1:D:88:LYS:O	5:D:720:HOH:O	2.20	0.49
1:A:220:GLU:OE2	5:A:713:HOH:O	2.20	0.49
1:D:143:HIS:CE1	1:D:328:GLU:HG2	2.48	0.48
2:C:401:PYR:H32	3:C:402:NAI:C5N	2.44	0.48
1:D:212:ARG:O	5:D:761:HOH:O	2.20	0.47
1:C:209:LYS:NZ	1:D:299:GLU:OE1	2.43	0.47
1:A:95:LYS:NZ	5:A:850:HOH:O	2.47	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:GLU:OE2	5:B:756:HOH:O	2.21	0.45
1:B:313:GLU:OE2	5:B:575:HOH:O	2.20	0.45
1:B:305:LYS:NZ	1:B:309:ASP:OD1	2.51	0.44
1:A:234:ASN:ND2	5:A:783:HOH:O	2.41	0.43
1:A:133:LEU:O	1:A:137[B]:VAL:HG23	2.18	0.43
1:C:41:VAL:O	1:C:69:ASN:ND2	2.51	0.42
1:A:88:LYS:NZ	1:A:94:ASP:OD1	2.40	0.42
1:A:128:VAL:HB	1:A:137[B]:VAL:HG22	2.02	0.42
1:D:208:ILE:HD11	1:D:218:ILE:CD1	2.51	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:518:HOH:O	5:D:515:HOH:O[3_555]	2.12	0.08
5:A:594:HOH:O	5:D:519:HOH:O[3_555]	2.15	0.05
5:A:543:HOH:O	5:D:511:HOH:O[2_665]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/334 (99%)	322 (97%)	9 (3%)	0	100	100
1	B	325/334 (97%)	316 (97%)	8 (2%)	1 (0%)	46	19
1	C	326/334 (98%)	321 (98%)	5 (2%)	0	100	100
1	D	325/334 (97%)	312 (96%)	13 (4%)	0	100	100
All	All	1307/1336 (98%)	1271 (97%)	35 (3%)	1 (0%)	56	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	155	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/281 (100%)	277 (99%)	3 (1%)	80	58
1	B	275/281 (98%)	273 (99%)	2 (1%)	88	73
1	C	275/281 (98%)	274 (100%)	1 (0%)	93	84
1	D	275/281 (98%)	274 (100%)	1 (0%)	93	84
All	All	1105/1124 (98%)	1098 (99%)	7 (1%)	90	78

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	93[A]	SER
1	A	93[B]	SER
1	A	327	LEU
1	B	163	PHE
1	B	279	MET
1	C	163	PHE
1	D	163	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	165	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PYR	A	401	-	2,5,5	0.23	0	2,6,6	0.31	0
3	NAI	A	402	-	38,48,48	0.94	2 (5%)	48,73,73	1.73	8 (16%)
2	PYR	B	401	-	2,5,5	0.21	0	2,6,6	0.45	0
4	NAD	B	402	-	38,48,48	0.78	1 (2%)	47,73,73	1.71	6 (12%)
2	PYR	C	401	-	2,5,5	0.27	0	2,6,6	0.19	0
3	NAI	C	402	-	38,48,48	0.97	2 (5%)	48,73,73	1.67	7 (14%)
2	PYR	D	401	-	2,5,5	0.26	0	2,6,6	0.32	0
4	NAD	D	402	-	38,48,48	0.79	1 (2%)	47,73,73	1.77	9 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PYR	A	401	-	-	0/0/4/4	0/0/0/0
3	NAI	A	402	-	-	0/25/72/72	0/5/5/5
2	PYR	B	401	-	-	0/0/4/4	0/0/0/0
4	NAD	B	402	-	-	0/22/62/62	0/5/5/5
2	PYR	C	401	-	-	0/0/4/4	0/0/0/0
3	NAI	C	402	-	-	0/25/72/72	0/5/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PYR	D	401	-	-	0/0/4/4	0/0/0/0
4	NAD	D	402	-	-	0/22/62/62	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	NAI	C5A-C4A	2.62	1.46	1.40
3	C	402	NAI	C5A-C4A	2.80	1.46	1.40
4	D	402	NAD	C5A-C4A	2.80	1.46	1.40
3	A	402	NAI	C6N-C5N	2.84	1.38	1.33
4	B	402	NAD	C5A-C4A	2.84	1.46	1.40
3	C	402	NAI	C6N-C5N	2.87	1.38	1.33

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	NAI	N3A-C2A-N1A	-8.51	122.38	128.89
4	D	402	NAD	N3A-C2A-N1A	-8.24	122.58	128.89
3	C	402	NAI	N3A-C2A-N1A	-7.95	122.81	128.89
4	B	402	NAD	N3A-C2A-N1A	-7.75	122.96	128.89
4	B	402	NAD	C1B-N9A-C4A	-4.05	120.83	126.94
4	B	402	NAD	C4A-C5A-N7A	-3.67	106.11	109.48
4	D	402	NAD	C1B-N9A-C4A	-3.65	121.43	126.94
3	A	402	NAI	C4B-O4B-C1B	-3.46	105.92	109.72
4	D	402	NAD	C4A-C5A-N7A	-3.42	106.33	109.48
3	C	402	NAI	C4A-C5A-N7A	-3.34	106.41	109.48
3	C	402	NAI	C4B-O4B-C1B	-2.92	106.51	109.72
3	A	402	NAI	C4A-C5A-N7A	-2.88	106.83	109.48
4	D	402	NAD	PN-O3-PA	-2.81	124.83	132.73
4	B	402	NAD	C4B-O4B-C1B	-2.67	106.79	109.72
3	A	402	NAI	C1B-N9A-C4A	-2.61	123.00	126.94
3	C	402	NAI	C1B-N9A-C4A	-2.57	123.06	126.94
3	C	402	NAI	PN-O3-PA	-2.46	125.83	132.73
4	D	402	NAD	C4B-O4B-C1B	-2.32	107.17	109.72
3	A	402	NAI	O3-PN-O5D	-2.24	96.98	102.94
3	A	402	NAI	C1D-N1N-C2N	-2.23	117.02	120.91
3	C	402	NAI	C1D-N1N-C2N	-2.13	117.19	120.91
4	D	402	NAD	O3-PN-O5D	-2.07	97.45	102.94
4	D	402	NAD	C3N-C7N-N7N	2.04	120.05	117.82
3	A	402	NAI	C2A-N1A-C6A	2.08	122.48	118.77
4	D	402	NAD	C2A-N1A-C6A	2.11	122.54	118.77
4	B	402	NAD	O4B-C1B-N9A	2.14	112.59	108.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	NAI	O4B-C1B-N9A	2.15	112.61	108.10
4	B	402	NAD	C3N-C7N-N7N	2.20	120.22	117.82
3	C	402	NAI	O4B-C1B-N9A	2.39	113.11	108.10
4	D	402	NAD	O4B-C1B-N9A	2.47	113.26	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	401	PYR	1	0
3	C	402	NAI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/334 (97%)	0.77	22 (6%) 20 20	11, 16, 30, 72	0
1	B	324/334 (97%)	1.41	59 (18%) 2 2	11, 19, 48, 101	0
1	C	324/334 (97%)	0.94	31 (9%) 10 10	12, 17, 32, 113	0
1	D	324/334 (97%)	1.53	58 (17%) 2 2	11, 18, 62, 173	0
All	All	1296/1336 (97%)	1.16	170 (13%) 5 4	11, 17, 43, 173	0

All (170) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	327	LEU	34.4
1	D	105	PHE	15.2
1	D	97	TRP	13.8
1	D	89	ALA	10.8
1	D	102	LEU	10.7
1	B	92	LYS	9.9
1	B	89	ALA	9.8
1	B	103	LEU	9.8
1	D	91	GLY	9.6
1	D	93	SER	9.5
1	D	96	GLU	8.4
1	D	90	PRO	8.2
1	B	105	PHE	8.1
1	C	327	LEU	8.1
1	B	327	LEU	8.0
1	B	93	SER	8.0
1	C	95	LYS	7.2
1	B	86	LEU	7.2
1	B	95	LYS	6.8
1	B	325	LEU	6.3
1	B	90	PRO	5.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	91	GLY	5.7
1	D	94	ASP	5.7
1	B	326	GLY	5.6
1	B	97	TRP	5.6
1	D	324	ALA	5.5
1	D	100	ASP	5.4
1	D	87	THR	5.3
1	C	146	LEU	5.3
1	B	96	GLU	5.2
1	D	318	LEU	5.2
1	D	146	LEU	5.1
1	B	102	LEU	4.9
1	B	316	ARG	4.9
1	D	325	LEU	4.8
1	D	92	LYS	4.8
1	D	316	ARG	4.7
1	D	307	LEU	4.6
1	B	108	LYS	4.4
1	A	94	ASP	4.3
1	D	238	GLN	4.3
1	D	104	PRO	4.2
1	D	5	MET	4.2
1	D	86	LEU	4.1
1	B	98	SER	4.1
1	B	87	THR	4.0
1	D	95	LYS	3.9
1	B	318	LEU	3.9
1	D	322	ILE	3.9
1	A	119	LYS	3.8
1	A	71	TYR	3.7
1	D	120	TYR	3.7
1	B	328	GLU	3.7
1	B	324	ALA	3.7
1	C	325	LEU	3.7
1	B	320	LYS	3.7
1	C	5	MET	3.6
1	B	101	ASP	3.6
1	D	328	GLU	3.6
1	D	326	GLY	3.6
1	D	239	GLY	3.5
1	D	98	SER	3.5
1	B	94	ASP	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	317	LYS	3.4
1	B	100	ASP	3.3
1	A	327	LEU	3.3
1	D	103	LEU	3.3
1	A	250	ILE	3.3
1	C	93	SER	3.3
1	B	119	LYS	3.3
1	D	321	ALA	3.1
1	A	74	ILE	3.1
1	B	117	ILE	3.1
1	D	312	VAL	3.1
1	A	5	MET	3.1
1	A	95	LYS	3.1
1	C	120	TYR	3.1
1	C	124	ALA	3.1
1	D	59	ALA	3.1
1	D	320	LYS	3.0
1	D	140	LEU	3.0
1	D	109	ILE	3.0
1	B	139	VAL	2.9
1	D	189	LYS	2.9
1	B	312	VAL	2.8
1	B	104	PRO	2.8
1	C	269	TYR	2.8
1	C	295	VAL	2.8
1	B	238	GLN	2.8
1	B	28	ALA	2.8
1	A	7	LYS	2.7
1	A	97	TRP	2.7
1	B	247	ALA	2.7
1	A	146	LEU	2.7
1	C	318	LEU	2.7
1	D	61	THR	2.7
1	A	218	ILE	2.7
1	A	113	VAL	2.7
1	B	25	TYR	2.6
1	C	166	PHE	2.6
1	D	306	GLU	2.6
1	C	59	ALA	2.6
1	C	117	ILE	2.6
1	C	119	LYS	2.6
1	B	313	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	146	LEU	2.6
1	C	7	LYS	2.6
1	D	58	ILE	2.6
1	B	135	VAL	2.6
1	A	25	TYR	2.6
1	A	90	PRO	2.5
1	D	204	LEU	2.5
1	D	310	LYS	2.5
1	B	209	LYS	2.5
1	B	32	LEU	2.5
1	C	140	LEU	2.5
1	C	316	ARG	2.5
1	B	322	ILE	2.5
1	B	239	GLY	2.4
1	D	71	TYR	2.4
1	B	99	ARG	2.4
1	B	323	LYS	2.4
1	D	247	ALA	2.4
1	C	94	ASP	2.4
1	B	115	GLU	2.4
1	B	6	THR	2.4
1	D	66	THR	2.4
1	C	27	CYS	2.4
1	D	27	CYS	2.4
1	C	65	VAL	2.4
1	B	319	GLN	2.3
1	D	99	ARG	2.3
1	B	27	CYS	2.3
1	B	142	GLU	2.3
1	A	65	VAL	2.3
1	C	123	ASN	2.3
1	B	242	TYR	2.2
1	C	25	TYR	2.2
1	D	250	ILE	2.2
1	B	58	ILE	2.2
1	B	220	GLU	2.2
1	D	72	GLU	2.2
1	B	88	LYS	2.2
1	B	120	TYR	2.2
1	C	196	TYR	2.2
1	C	328	GLU	2.2
1	A	148	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	275	GLY	2.2
1	C	58	ILE	2.1
1	D	243	PHE	2.1
1	D	26	LEU	2.1
1	C	121	CYS	2.1
1	C	115	GLU	2.1
1	A	6	THR	2.1
1	B	66	THR	2.1
1	A	269	TYR	2.1
1	D	156	GLY	2.1
1	A	93[A]	SER	2.1
1	B	208	ILE	2.1
1	C	28	ALA	2.1
1	C	212	ARG	2.1
1	D	139	VAL	2.1
1	A	328	GLU	2.1
1	B	246	ALA	2.0
1	B	204	LEU	2.0
1	D	211	GLY	2.0
1	D	246	ALA	2.0
1	D	101	ASP	2.0
1	A	61	THR	2.0
1	B	57	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	PYR	C	401	6/6	0.92	0.14	1.49	16,23,39,39	0
2	PYR	B	401	6/6	0.83	0.19	1.22	27,38,48,48	0
2	PYR	D	401	6/6	0.87	0.18	0.21	29,39,50,50	0
3	NAI	C	402	44/44	0.93	0.11	0.05	13,21,29,34	0
4	NAD	D	402	44/44	0.89	0.15	-0.08	22,33,41,47	0
3	NAI	A	402	44/44	0.95	0.12	-0.17	12,19,28,38	0
2	PYR	A	401	6/6	0.93	0.11	-0.18	16,20,36,36	0
4	NAD	B	402	44/44	0.90	0.13	-0.21	18,31,40,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.