



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:40 AM GMT

PDB ID : 3PMH
Title : Mechanism of Sulfotyrosine-Mediated Glycoprotein Ib Interaction with Two Distinct alpha-Thrombin Sites
Authors : Varughese, K.I.; Celikel, R.
Deposited on : 2010-11-16
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	36	<div> <div>11%</div> <div> <div>47%</div> <div>47%</div> <div>6%</div> </div> </div>
2	B	259	<div> <div>57%</div> <div>40%</div> <div>.</div> </div>
3	G	290	<div> <div>4%</div> <div>42%</div> <div>47%</div> <div>9%</div> <div>.</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4723 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THROMBIN ALPHA-CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	36	Total	C	N	O	S	0	0	0
			287	177	48	61	1			

- Molecule 2 is a protein called THROMBIN BETA-CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	259	Total	C	N	O	S	0	0	0
			2093	1334	370	375	14			


- Molecule 3 is a protein called Platelet glycoprotein Ib alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	284	Total	C	N	O	S	0	0	0
			2237	1427	363	436	11			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	65	ALA	CYS	ENGINEERED MUTATION	UNP P07359

- Molecule 4 is D-PHENYLALANYL-N-[(3S)-6-CARBAMIMIDAMIDO-1-CHLORO-2-OXO HEXAN-3-YL]-L-PROLINAMIDE (three-letter code: 0G7) (formula: C₂₁H₃₁ClN₆O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	Cl	N	O	0
			31	21	1	6	3	

- Molecule 5 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 6 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	G	1	Total	C	N	O	0	0
			14	8	1	5		

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	67.66 Å 67.66 Å 329.73 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.20 47.34 – 3.19	Depositor EDS
% Data completeness (in resolution range)	94.2 (50.00-3.20) 88.9 (47.34-3.19)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.49 (at 3.19 Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.220 , 0.263 0.231 , 0.277	Depositor DCC
R_{free} test set	611 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	56.8	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 37.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 12905 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4723	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0G7, MAN, NAG, TYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.42	0/290	0.69	0/384
2	B	0.44	0/2148	0.70	0/2903
3	G	0.37	0/2235	0.65	0/3050
All	All	0.41	0/4673	0.68	0/6337

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	287	0	278	32	0
2	B	2093	0	2063	110	0
3	G	2237	0	2225	179	0
4	B	31	0	28	20	0
5	B	61	0	52	1	0
6	G	14	0	13	3	0
All	All	4723	0	4659	303	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 303 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:126:ARG:CD	2:B:126:ARG:H	1.56	1.13
2:B:126:ARG:HD2	2:B:126:ARG:N	1.66	1.11
3:G:25:LEU:HD23	3:G:25:LEU:H	1.18	1.09
2:B:195:SER:HB3	4:B:301:OG7:O2	1.55	1.03
3:G:124:GLY:HA2	3:G:146:PRO:O	1.63	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	34/36 (94%)	28 (82%)	5 (15%)	1 (3%)	6	36
2	B	257/259 (99%)	240 (93%)	15 (6%)	2 (1%)	24	69
3	G	279/290 (96%)	209 (75%)	49 (18%)	21 (8%)	1	9
All	All	570/585 (97%)	477 (84%)	69 (12%)	24 (4%)	3	26

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	77(A)	ARG
3	G	2	PRO
3	G	9	VAL
3	G	189	LYS
3	G	234	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	31/31 (100%)	29 (94%)	2 (6%)	21	61
2	B	225/225 (100%)	216 (96%)	9 (4%)	38	77
3	G	251/256 (98%)	237 (94%)	14 (6%)	26	68
All	All	507/512 (99%)	482 (95%)	25 (5%)	31	72

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	1	HIS
3	G	8	LYS
3	G	242	ASN
3	G	7	SER
3	G	11	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	61	ASN
3	G	86	HIS
3	G	247	GLN
3	G	71	GLN
3	G	97	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TYS	G	276	3	15,16,17	1.79	2 (13%)	16,22,24	1.09	1 (6%)
3	TYS	G	278	3	15,16,17	1.47	1 (6%)	16,22,24	0.99	1 (6%)
3	TYS	G	279	3	15,16,17	1.59	2 (13%)	16,22,24	1.09	2 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TYS	G	276	3	-	0/9/11/13	0/1/1/1
3	TYS	G	278	3	-	0/9/11/13	0/1/1/1
3	TYS	G	279	3	-	0/9/11/13	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	276	TYS	OH-S	-5.93	1.52	1.63
3	G	279	TYS	OH-S	-5.21	1.54	1.63
3	G	278	TYS	OH-S	-4.94	1.54	1.63
3	G	276	TYS	OH-CZ	-2.80	1.38	1.42
3	G	279	TYS	OH-CZ	-2.50	1.38	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	279	TYS	OH-CZ-CE2	-2.01	114.77	118.74
3	G	278	TYS	OH-CZ-CE1	2.23	123.14	118.74
3	G	279	TYS	OH-CZ-CE1	2.65	123.98	118.74
3	G	276	TYS	OH-CZ-CE1	2.77	124.22	118.74

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	276	TYS	1	0
3	G	278	TYS	3	0
3	G	279	TYS	1	0

5.5 Carbohydrates [i](#)

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	NAG	B	302	2,5	14,14,15	1.80	2 (14%)	15,19,21	2.07	5 (33%)
5	NAG	B	303	5	14,14,15	0.65	0	15,19,21	1.12	2 (13%)
5	MAN	B	304	5	11,11,12	0.91	0	14,15,17	0.83	0
5	MAN	B	305	5	11,11,12	0.79	0	14,15,17	0.77	1 (7%)
5	MAN	B	306	5	11,11,12	0.65	0	14,15,17	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	302	2,5	-	0/6/23/26	0/1/1/1
5	NAG	B	303	5	-	0/6/23/26	0/1/1/1
5	MAN	B	304	5	-	0/2/19/22	0/1/1/1
5	MAN	B	305	5	-	0/2/19/22	0/1/1/1
5	MAN	B	306	5	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	302	NAG	C8-C7	2.67	1.55	1.50
5	B	302	NAG	C1-C2	5.38	1.59	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	302	NAG	C2-N2-C7	-3.15	118.99	123.04
5	B	302	NAG	O7-C7-C8	-2.43	117.60	122.06
5	B	302	NAG	O5-C5-C6	-2.35	102.25	107.35
5	B	303	NAG	C4-C3-C2	-2.14	107.90	111.23
5	B	303	NAG	C3-C4-C5	-2.00	106.70	110.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	303	NAG	1	0
5	B	304	MAN	1	0

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	0G7	B	301	-	27,32,32	0.81	2 (7%)	35,42,42	1.03	2 (5%)
6	NAG	G	601	3	14,14,15	1.34	3 (21%)	15,19,21	1.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	0G7	B	301	-	-	1/30/43/43	0/2/2/2
6	NAG	G	601	3	-	0/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	301	0G7	C3-CL1	2.03	1.83	1.76
6	G	601	NAG	C4-C3	2.05	1.57	1.52
6	G	601	NAG	C3-C2	2.15	1.57	1.52
4	B	301	0G7	O2-C2	2.80	1.26	1.21
6	G	601	NAG	C1-C2	3.29	1.57	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	B	301	0G7	C3-C2-CA2	2.54	120.77	116.24
4	B	301	0G7	CB-CA-C	3.04	115.61	108.80

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	301	0G7	CA2-C2-C3-CL1

There are no ring outliers.

2 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	301	0G7	20	0
6	G	601	NAG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	36/36 (100%)	0.15	4 (11%) 7 4	40, 63, 136, 153	0
2	B	259/259 (100%)	-0.42	0 100 100	26, 49, 76, 101	0
3	G	281/290 (96%)	0.16	12 (4%) 39 25	38, 81, 127, 139	0
All	All	576/585 (98%)	-0.10	16 (2%) 56 42	26, 62, 122, 153	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1(H)	THR	8.7
3	G	15	VAL	3.5
3	G	234	VAL	3.4
3	G	270	GLU	3.1
1	A	1(G)	PHE	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	TYS	G	278	16/17	0.87	0.24	-	84,94,100,101	4
3	TYS	G	279	16/17	0.94	0.21	-	67,75,87,89	0
3	TYS	G	276	16/17	0.84	0.33	-	97,103,120,122	4

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	NAG	B	302	14/15	0.92	0.19	-	93,101,107,118	0
5	MAN	B	306	11/12	0.55	0.37	-	128,134,137,139	0
5	NAG	B	303	14/15	0.85	0.20	-	99,104,108,112	0
5	MAN	B	304	11/12	0.81	0.20	-	104,112,116,118	0
5	MAN	B	305	11/12	0.72	0.29	-	78,90,98,99	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	0G7	B	301	31/31	0.86	0.25	0.31	20,36,44,56	0
6	NAG	G	601	14/15	0.82	0.17	-	86,102,107,112	0

6.5 Other polymers [i](#)

There are no such residues in this entry.