



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:44 AM GMT

PDB ID : 3PO5
Title : Structure of a mutant of the large fragment of DNA polymerase I from *Thermus*
Auquaticus in complex with an abasic site and ddATP
Authors : Marx, A.; Diederichs, K.; Obeid, S.
Deposited on : 2010-11-22
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	<div> <div>8%</div> <div>74%</div> <div>22%</div> <div>..</div> </div>
2	B	12	<div> <div>58%</div> <div>42%</div> </div>
3	C	13	<div> <div>46%</div> <div>46%</div> <div>8%</div> </div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4871 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	528	Total	C	N	O	S	0	0	0
			4197	2666	756	764	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	614	LYS	ILE	ENGINEERED MUTATION	UNP P19821
A	747	LYS	MET	ENGINEERED MUTATION	UNP P19821

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(2DA))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	P	0	0	0
			242	115	50	66	11			


- Molecule 3 is a DNA chain called DNA (5'-D(P*(3DR)P*TP*GP*CP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	P	0	0	0
			257	121	43	80	13			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is 2',3'-DIDEOXYADENOSINE TRIPHOSPHATE (three-letter code: DDS) (formula: C₁₀H₁₆N₅O₁₁P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			29	10	5	11	3		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	96	Total	O	0	0
			96	96		
8	B	12	Total	O	0	0
			12	12		
8	C	23	Total	O	0	0
			23	23		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase I



- Molecule 2: DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(2DA))-3')



- Molecule 3: DNA (5'-D(P*(3DR)P*TP*GP*CP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	109.30 Å 109.30 Å 90.32 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.33 – 2.39 47.33 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.33-2.39) 99.7 (47.33-2.39)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.39 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: DEV_572)	Depositor
R, R_{free}	0.210 , 0.266 0.209 , 0.269	Depositor DCC
R_{free} test set	1287 reflections (5.44%)	DCC
Wilson B-factor (Å ²)	36.8	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.3	EDS
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 24931 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4871	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, PGE, 3DR, DDS, 2DA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/4284	0.44	0/5799
2	B	0.59	0/249	1.23	1/382 (0.3%)
3	C	0.64	0/274	1.34	3/421 (0.7%)
All	All	0.34	0/4807	0.61	4/6602 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	214	DG	C1'-O4'-C4'	-6.65	103.45	110.10
3	C	214	DG	O4'-C1'-N9	6.54	112.58	108.00
3	C	206	DG	O4'-C4'-C3'	-5.48	102.31	104.50
2	B	108	DG	O4'-C1'-N9	5.16	111.61	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4197	0	4246	114	0
2	B	242	0	134	5	0
3	C	257	0	144	4	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	8	0	8	2	0
6	A	29	0	12	4	0
7	B	6	0	8	0	0
8	A	96	0	0	0	0
8	B	12	0	0	0	0
8	C	23	0	0	0	0
All	All	4871	0	4552	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (124) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:102:DA:H2''	2:B:103:DC:H5'	1.33	1.05
6:A:835:DDS:H8	6:A:835:DDS:H5'	1.43	0.96
6:A:835:DDS:C8	6:A:835:DDS:H5'	2.02	0.89
2:B:102:DA:C2'	2:B:103:DC:H5'	2.11	0.80
1:A:374:MET:CE	1:A:387:PRO:HA	2.18	0.74
1:A:567:THR:HG22	1:A:567:THR:O	1.87	0.72
1:A:626:GLU:O	1:A:630:ARG:HD3	1.88	0.71
1:A:588:THR:HG23	1:A:589:PRO:HD2	1.73	0.70
1:A:785:ASP:OD1	1:A:786:GLU:HG3	1.93	0.68
1:A:354:LYS:HE2	1:A:445:GLU:OE1	1.94	0.66
1:A:685:PRO:HB2	1:A:688:GLU:HG3	1.79	0.65
1:A:632:PHE:CE1	1:A:638:ILE:HG12	2.32	0.65
1:A:567:THR:CG2	1:A:567:THR:O	2.45	0.64
1:A:302:PRO:HB2	1:A:328:ARG:HD3	1.81	0.62
1:A:376:LEU:HD22	1:A:420:LEU:HD22	1.82	0.62
1:A:386:THR:HG23	1:A:387:PRO:HD2	1.83	0.60
1:A:638:ILE:N	1:A:638:ILE:HD12	2.17	0.60
2:B:101:DG:H2''	2:B:102:DA:O5'	2.03	0.59
1:A:503:ILE:CD1	1:A:522:LEU:HD11	2.32	0.58
1:A:374:MET:HE1	1:A:387:PRO:HA	1.85	0.58
1:A:335:ALA:HB1	1:A:341:ALA:HB2	1.86	0.58
1:A:519:LEU:CB	1:A:533:LEU:HD11	2.33	0.57
1:A:815:VAL:HG23	1:A:816:PRO:HD2	1.86	0.56
2:B:102:DA:H2''	2:B:103:DC:C5'	2.24	0.56
1:A:299:TRP:CZ2	1:A:341:ALA:HB1	2.39	0.56
1:A:609:LEU:HD11	1:A:789:LEU:HD22	1.88	0.56
1:A:636:ARG:NE	1:A:641:GLU:HG3	2.21	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:557:THR:OG1	1:A:561:HIS:HE1	1.90	0.55
1:A:661:ALA:O	1:A:665:ILE:HG21	2.07	0.55
1:A:447:THR:HG21	1:A:778:ARG:HD3	1.88	0.54
1:A:518:VAL:O	1:A:522:LEU:HD13	2.08	0.54
1:A:576:SER:O	3:C:208:DG:H4'	2.07	0.54
1:A:401:GLU:HG3	1:A:404:GLU:H	1.72	0.54
1:A:374:MET:CE	1:A:387:PRO:CA	2.84	0.53
1:A:627:ASN:O	1:A:631:VAL:HG23	2.09	0.53
1:A:466:GLU:HG2	1:A:538:LEU:HD21	1.90	0.53
1:A:519:LEU:HB2	1:A:533:LEU:HD11	1.89	0.53
1:A:657:LEU:C	1:A:659:ARG:H	2.12	0.53
1:A:632:PHE:CB	1:A:815:VAL:CG2	2.87	0.52
1:A:780:LEU:HD11	1:A:790:GLU:HB2	1.91	0.52
1:A:299:TRP:CE2	1:A:341:ALA:HB1	2.44	0.51
1:A:609:LEU:CD1	1:A:789:LEU:HD22	2.41	0.51
1:A:588:THR:HG23	1:A:589:PRO:CD	2.41	0.50
1:A:643:ALA:HB1	1:A:658:MET:O	2.11	0.50
1:A:657:LEU:C	1:A:659:ARG:N	2.64	0.50
1:A:673:MET:HG3	1:A:677:ARG:HB2	1.94	0.50
1:A:661:ALA:O	1:A:665:ILE:CG2	2.59	0.49
1:A:526:HIS:ND1	1:A:527:PRO:HD2	2.27	0.49
1:A:751:MET:HB3	1:A:752:PRO:HD3	1.93	0.49
1:A:299:TRP:CG	1:A:300:PRO:HA	2.46	0.49
1:A:678:LEU:O	1:A:682:LEU:HB2	2.13	0.49
1:A:549:LEU:HD22	1:A:560:LEU:HD21	1.93	0.49
1:A:632:PHE:CB	1:A:815:VAL:HG23	2.43	0.49
1:A:630:ARG:HD2	1:A:630:ARG:N	2.27	0.48
1:A:480:HIS:HE1	1:A:497:GLU:OE2	1.96	0.48
1:A:335:ALA:HB1	1:A:341:ALA:CB	2.42	0.48
1:A:632:PHE:HB3	1:A:815:VAL:CG2	2.44	0.48
1:A:295:GLU:HG2	1:A:332:VAL:HG12	1.95	0.48
1:A:386:THR:HG22	1:A:388:GLU:N	2.28	0.48
1:A:537:GLU:OE1	1:A:588:THR:HG21	2.14	0.48
1:A:503:ILE:CD1	1:A:522:LEU:CD1	2.92	0.47
1:A:671:TYR:HE2	6:A:835:DDS:H2	1.79	0.47
1:A:658:MET:SD	1:A:658:MET:O	2.72	0.47
1:A:351:LEU:HD11	1:A:412:LEU:HD12	1.95	0.47
1:A:380:LEU:HG	1:A:420:LEU:HD11	1.95	0.47
1:A:643:ALA:HB2	1:A:659:ARG:HA	1.96	0.47
1:A:302:PRO:HD3	1:A:333:HIS:NE2	2.29	0.47
1:A:643:ALA:CB	1:A:659:ARG:HA	2.45	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:808:GLU:HG2	1:A:819:VAL:HG23	1.96	0.47
1:A:679:SER:HB3	1:A:689:ALA:HB2	1.97	0.47
1:A:616:LEU:HD23	1:A:638:ILE:HG21	1.98	0.46
1:A:549:LEU:HB2	1:A:550:PRO:HD3	1.97	0.46
1:A:632:PHE:HB2	1:A:815:VAL:HG23	1.98	0.46
3:C:210:DC:H2'	3:C:211:DG:C8	2.51	0.46
1:A:374:MET:HE1	1:A:386:THR:O	2.16	0.46
1:A:297:ALA:HB1	1:A:298:PRO:HD2	1.98	0.46
1:A:396:GLY:HA3	1:A:408:LEU:CD1	2.46	0.45
1:A:380:LEU:HD12	1:A:416:LEU:HD12	1.98	0.45
1:A:630:ARG:CD	1:A:630:ARG:N	2.80	0.45
1:A:526:HIS:CG	1:A:527:PRO:HD2	2.52	0.45
1:A:699:SER:C	1:A:701:PRO:HD3	2.37	0.45
5:A:834:PGE:H32	3:C:209:DC:H4'	1.97	0.45
1:A:490:LEU:O	1:A:494:LEU:HD13	2.16	0.45
1:A:632:PHE:HE1	1:A:638:ILE:HG12	1.80	0.45
1:A:822:GLY:HA3	1:A:830:ALA:O	2.17	0.45
1:A:445:GLU:O	1:A:561:HIS:HD2	2.00	0.45
1:A:682:LEU:HD23	1:A:684:ILE:CD1	2.47	0.45
6:A:835:DDS:O4'	2:B:112:2DA:H2''	2.17	0.44
1:A:544:THR:O	5:A:834:PGE:H4	2.18	0.44
1:A:386:THR:CG2	1:A:388:GLU:OE1	2.65	0.44
1:A:374:MET:HE3	1:A:387:PRO:CA	2.47	0.44
1:A:434:GLU:HA	1:A:434:GLU:OE1	2.17	0.44
1:A:519:LEU:HB3	1:A:533:LEU:HD11	2.00	0.44
1:A:684:ILE:HB	1:A:685:PRO:HD2	2.00	0.43
1:A:379:LEU:CD1	1:A:430:TYR:HB2	2.48	0.43
1:A:815:VAL:CG2	1:A:816:PRO:HD2	2.46	0.43
1:A:435:ARG:N	1:A:436:PRO:HD2	2.33	0.43
1:A:769:PHE:N	1:A:770:PRO:HD2	2.33	0.43
1:A:637:ASP:O	1:A:641:GLU:CB	2.67	0.43
1:A:315:GLU:HA	1:A:316:PRO:HD2	1.94	0.43
1:A:519:LEU:HD13	1:A:533:LEU:CD1	2.50	0.42
1:A:665:ILE:HD11	1:A:696:TYR:CD1	2.54	0.42
1:A:665:ILE:HD11	1:A:696:TYR:CG	2.54	0.42
1:A:668:GLY:HA3	1:A:673:MET:HE2	2.01	0.42
1:A:684:ILE:HB	1:A:685:PRO:CD	2.50	0.42
1:A:374:MET:HE1	1:A:386:THR:C	2.41	0.41
1:A:676:HIS:O	1:A:677:ARG:C	2.58	0.41
1:A:354:LYS:NZ	1:A:354:LYS:HB3	2.34	0.41
1:A:621:HIS:CD2	1:A:814:ALA:H	2.38	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:685:PRO:CB	1:A:688:GLU:HG3	2.47	0.41
1:A:374:MET:HE1	1:A:387:PRO:CA	2.50	0.41
1:A:492:ARG:O	1:A:496:ASP:HB2	2.21	0.41
1:A:634:GLU:OE1	1:A:636:ARG:HD2	2.20	0.41
3:C:214:DG:C8	3:C:215:DT:H72	2.55	0.41
1:A:637:ASP:O	1:A:641:GLU:HB3	2.21	0.41
1:A:537:GLU:HG3	1:A:538:LEU:N	2.32	0.40
1:A:588:THR:HG22	1:A:590:LEU:H	1.86	0.40
1:A:588:THR:CG2	1:A:589:PRO:N	2.84	0.40
1:A:295:GLU:HG3	1:A:330:GLY:O	2.21	0.40
1:A:602:GLU:CD	1:A:602:GLU:H	2.25	0.40
1:A:638:ILE:CD1	1:A:638:ILE:N	2.83	0.40
1:A:778:ARG:HD2	1:A:790:GLU:OE1	2.22	0.40
1:A:490:LEU:HD11	1:A:494:LEU:HD11	2.03	0.40
1:A:554:HIS:HA	1:A:555:PRO:HD3	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	524/540 (97%)	494 (94%)	29 (6%)	1 (0%)	52	69

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	586	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	431/441 (98%)	415 (96%)	16 (4%)	41 62

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	354	LYS
1	A	416	LEU
1	A	426	LEU
1	A	427	LEU
1	A	487	ARG
1	A	537	GLU
1	A	583	ASN
1	A	609	LEU
1	A	657	LEU
1	A	658	MET
1	A	660	ARG
1	A	665	ILE
1	A	681	GLU
1	A	727	ARG
1	A	787	LEU
1	A	789	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	480	HIS
1	A	561	HIS
1	A	583	ASN
1	A	621	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	2DA	B	112	3,2	14,22,23	1.10	1 (7%)	13,31,34	2.48	4 (30%)
3	3DR	C	204	3	7,11,12	0.42	0	8,14,17	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2DA	B	112	3,2	-	0/3/18/19	0/3/3/3
3	3DR	C	204	3	-	0/3/15/16	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	112	2DA	C5-C4	3.11	1.47	1.40

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	112	2DA	N3-C2-N1	-7.05	123.50	128.89
2	B	112	2DA	C4-C5-N7	-2.43	107.24	109.48
2	B	112	2DA	C2'-C1'-N9	-2.30	107.83	112.49
2	B	112	2DA	C3'-C2'-C1'	2.24	105.21	102.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	112	2DA	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	PGE	A	833	-	3,3,9	0.45	0	2,2,8	0.43	0
5	PGE	A	834	-	3,3,9	0.53	0	2,2,8	0.33	0
6	DDS	A	835	4	23,31,31	1.20	3 (13%)	28,48,48	2.38	5 (17%)
7	GOL	B	1	-	5,5,5	0.22	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PGE	A	833	-	-	0/1/1/7	0/0/0/0
5	PGE	A	834	-	-	0/1/1/7	0/0/0/0
6	DDS	A	835	4	-	0/18/31/31	0/3/3/3
7	GOL	B	1	-	-	0/4/4/4	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	835	DDS	C5'-C4'	-2.78	1.42	1.50
6	A	835	DDS	C3'-C2'	-2.24	1.47	1.54
6	A	835	DDS	C6-N6	2.72	1.43	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	835	DDS	N3-C2-N1	-9.50	121.62	128.89
6	A	835	DDS	PB-O3B-PG	-3.39	121.30	132.67
6	A	835	DDS	PB-O3A-PA	-3.07	124.11	132.73
6	A	835	DDS	C4-C5-N7	-2.55	107.13	109.48
6	A	835	DDS	C3'-C2'-C1'	3.72	106.87	102.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	834	PGE	2	0
6	A	835	DDS	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	528/540 (97%)	0.62	45 (8%) 13 13	26, 49, 105, 132	0
2	B	11/12 (91%)	-0.16	0 100 100	28, 32, 62, 71	0
3	C	12/13 (92%)	-0.39	0 100 100	24, 32, 51, 54	0
All	All	551/565 (97%)	0.58	45 (8%) 14 14	24, 48, 105, 132	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	683	ALA	8.3
1	A	656	PRO	6.5
1	A	661	ALA	5.9
1	A	645	TRP	5.4
1	A	689	ALA	5.3
1	A	636	ARG	4.8
1	A	644	SER	4.7
1	A	688	GLU	4.7
1	A	692	PHE	4.4
1	A	643	ALA	4.1
1	A	499	GLY	4.0
1	A	686	TYR	4.0
1	A	345	LEU	3.9
1	A	657	LEU	3.8
1	A	680	GLN	3.7
1	A	695	ARG	3.6
1	A	642	THR	3.6
1	A	658	MET	3.4
1	A	682	LEU	3.3
1	A	662	ALA	3.3
1	A	677	ARG	3.3
1	A	696	TYR	3.2
1	A	673	MET	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	667	PHE	3.2
1	A	691	ALA	3.0
1	A	631	VAL	2.9
1	A	693	ILE	2.9
1	A	676	HIS	2.8
1	A	697	PHE	2.8
1	A	421	GLU	2.8
1	A	340	LYS	2.7
1	A	503	ILE	2.7
1	A	684	ILE	2.7
1	A	299	TRP	2.6
1	A	704	ARG	2.6
1	A	335	ALA	2.5
1	A	698	GLN	2.3
1	A	630	ARG	2.2
1	A	670	LEU	2.1
1	A	627	ASN	2.1
1	A	341	ALA	2.1
1	A	330	GLY	2.1
1	A	685	PRO	2.1
1	A	333	HIS	2.0
1	A	365	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	3DR	C	204	11/12	0.90	0.17	-	66,67,85,86	0
2	2DA	B	112	20/21	0.98	0.11	-	27,30,40,42	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	DDS	A	835	29/29	0.88	0.14	-0.76	50,72,94,271	0
5	PGE	A	834	4/10	0.96	0.10	-1.95	29,29,30,30	0
5	PGE	A	833	4/10	0.97	0.07	-4.60	46,47,47,47	0
4	MG	A	1	1/1	0.85	0.11	-	64,64,64,64	0
7	GOL	B	1	6/6	0.85	0.19	-	43,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.