



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:35 PM GMT

PDB ID : 1PQC  
Title : HUMAN LXR BETA HORMONE RECEPTOR COMPLEXED WITH  
T0901317  
Authors : Farnegardh, M.; Bonn, T.; Sun, S.; Ljunggren, J.; Ahola, H.; Wilhelmsson, A.;  
Gustafsson, J.-A.; Carlquist, M.  
Deposited on : 2003-06-18  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	<div> <div>5%</div> <div> <div></div> <div>77%</div> <div>13%</div> <div>• 8%</div> </div> </div>
1	B	253	<div> <div>2%</div> <div> <div></div> <div>78%</div> <div>15%</div> <div>• 6%</div> </div> </div>
1	C	253	<div> <div>3%</div> <div> <div></div> <div>69%</div> <div>21%</div> <div>• 9%</div> </div> </div>
1	D	253	<div> <div>10%</div> <div> <div></div> <div>68%</div> <div>18%</div> <div>• 13%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7782 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Oxysterols receptor LXR-beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	232	Total	C	N	O	S	0	0	0
			1891	1210	331	343	7			
1	B	238	Total	C	N	O	S	0	0	0
			1925	1230	337	351	7			
1	C	231	Total	C	N	O	S	0	0	0
			1887	1207	333	340	7			
1	D	219	Total	C	N	O	S	0	0	0
			1774	1133	313	321	7			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	209	GLY	-	INSERTION	UNP P55055
A	210	SER	-	INSERTION	UNP P55055
A	211	HIS	-	INSERTION	UNP P55055
A	212	MET	-	INSERTION	UNP P55055
B	209	GLY	-	INSERTION	UNP P55055
B	210	SER	-	INSERTION	UNP P55055
B	211	HIS	-	INSERTION	UNP P55055
B	212	MET	-	INSERTION	UNP P55055
C	209	GLY	-	INSERTION	UNP P55055
C	210	SER	-	INSERTION	UNP P55055
C	211	HIS	-	INSERTION	UNP P55055
C	212	MET	-	INSERTION	UNP P55055
D	209	GLY	-	INSERTION	UNP P55055
D	210	SER	-	INSERTION	UNP P55055
D	211	HIS	-	INSERTION	UNP P55055
D	212	MET	-	INSERTION	UNP P55055

- Molecule 2 is N-(2,2,2-TRIFLUOROETHYL)-N-{4-[2,2,2-TRIFLUORO-1-HYDROXY-1-(TRIFLUOROMETHYL)ETHYL]PHENYL}BENZENESULFONAMIDE (three-letter code: 444) (formula: C<sub>17</sub>H<sub>12</sub>F<sub>9</sub>NO<sub>3</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 31	C 17	F 9	N 1	O 3	S 1	0	0
2	B	1	Total 31	C 17	F 9	N 1	O 3	S 1	0	0
2	C	1	Total 31	C 17	F 9	N 1	O 3	S 1	0	0
2	D	1	Total 31	C 17	F 9	N 1	O 3	S 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	36	Total	O	0	0
			36	36		
3	B	72	Total	O	0	0
			72	72		
3	C	52	Total	O	0	0
			52	52		
3	D	21	Total	O	0	0
			21	21		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Oxysterols receptor LXR-beta



#### • Molecule 1: Oxysterols receptor LXR-beta




#### • Molecule 1: Oxysterols receptor LXR-beta



#### • Molecule 1: Oxysterols receptor LXR-beta





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.72Å 103.26Å 176.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.80 40.48 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.00-2.80) 99.9 (40.48-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.74 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.1.19	Depositor
R, $R_{free}$	0.195 , 0.262 0.244 , 0.244	Depositor DCC
$R_{free}$ test set	1353 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	52.1	Xtriage
Anisotropy	0.243	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 27.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 27100 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	7782	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 444

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	0/1928	0.84	5/2607 (0.2%)
1	B	0.71	0/1963	0.87	4/2656 (0.2%)
1	C	0.71	0/1922	0.86	4/2595 (0.2%)
1	D	0.68	1/1804 (0.1%)	0.82	6/2436 (0.2%)
All	All	0.69	1/7617 (0.0%)	0.85	19/10294 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	287	LYS	CD-CE	5.13	1.64	1.51

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	284	ASP	CB-CG-OD2	7.14	124.73	118.30
1	C	299	ASP	CB-CG-OD2	6.63	124.27	118.30
1	B	332	ASP	CB-CG-OD2	6.44	124.10	118.30
1	C	262	ASP	CB-CG-OD2	6.24	123.92	118.30
1	B	338	ASP	CB-CG-OD2	6.18	123.86	118.30
1	D	458	ASP	CB-CG-OD2	5.98	123.68	118.30
1	A	262	ASP	CB-CG-OD2	5.96	123.66	118.30
1	D	414	ASP	CB-CG-OD2	5.80	123.53	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	262	ASP	CB-CG-OD2	5.79	123.51	118.30
1	A	245	ASP	CB-CG-OD2	5.71	123.43	118.30
1	D	366	ASP	CB-CG-OD2	5.62	123.36	118.30
1	C	366	ASP	CB-CG-OD2	5.48	123.23	118.30
1	A	332	ASP	CB-CG-OD2	5.48	123.23	118.30
1	B	299	ASP	CB-CG-OD2	5.37	123.14	118.30
1	D	299	ASP	CB-CG-OD2	5.36	123.12	118.30
1	A	446	ASP	CB-CG-OD2	5.30	123.07	118.30
1	C	446	ASP	CB-CG-OD2	5.22	123.00	118.30
1	D	284	ASP	CB-CG-OD2	5.12	122.90	118.30
1	A	299	ASP	CB-CG-OD2	5.06	122.86	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	248	LYS	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1891	0	1913	11	0
1	B	1925	0	1936	18	0
1	C	1887	0	1914	24	0
1	D	1774	0	1788	15	0
2	A	31	0	12	0	0
2	B	31	0	12	0	0
2	C	31	0	12	0	0
2	D	31	0	12	0	0
3	A	36	0	0	0	0
3	B	72	0	0	1	0
3	C	52	0	0	0	0
3	D	21	0	0	0	0
All	All	7782	0	7599	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:257:ASP:HB3	1:B:258:PRO:HD2	1.67	0.77
1:B:257:ASP:CB	1:B:258:PRO:HD2	2.18	0.73
1:C:351:ASN:O	1:C:355:GLU:HG3	1.98	0.64
1:C:409:ILE:O	1:C:410:LYS:C	2.35	0.63
1:B:257:ASP:HB3	1:B:258:PRO:CD	2.30	0.61
1:B:257:ASP:CB	1:B:258:PRO:CD	2.78	0.60
1:D:233:ALA:O	1:D:237:GLN:HG3	2.00	0.60
1:B:437:GLU:OE2	3:B:50:HOH:O	2.16	0.59
1:A:349:PHE:CE1	1:A:353:ILE:HD11	2.38	0.58
1:D:252:TRP:CZ2	1:D:267:ARG:HD3	2.41	0.55
1:B:418:PHE:HB3	1:B:419:PRO:HD3	1.89	0.55
1:B:403:LEU:HD21	1:B:421:MET:HE2	1.90	0.53
1:C:373:LEU:HD22	1:C:425:LEU:HD21	1.90	0.53
1:B:241:ARG:NH1	1:B:245:ASP:OD2	2.41	0.53
1:C:321:ASN:HD22	1:C:324:THR:H	1.56	0.53
1:D:242:SER:OG	1:D:281:GLU:OE2	2.19	0.53
1:C:418:PHE:CZ	1:C:422:LEU:HD11	2.44	0.52
1:C:315:GLU:OE1	1:C:319:ARG:NH1	2.42	0.52
1:B:246:GLN:HE22	1:B:332:ASP:HB2	1.75	0.52
1:A:253:PRO:HG2	1:A:263:ALA:HB1	1.91	0.52
1:C:349:PHE:CZ	1:C:353:ILE:HD11	2.46	0.51
1:D:291:GLY:O	1:D:294:GLN:HG2	2.11	0.51
1:C:311:ILE:HG12	1:C:374:ILE:HG23	1.92	0.51
1:D:252:TRP:CE2	1:D:267:ARG:HD3	2.46	0.50
1:A:416:LEU:CD1	1:B:397:GLN:HE21	2.25	0.50
1:C:401:GLU:HG2	1:D:416:LEU:HD21	1.93	0.49
1:C:397:GLN:HB3	1:C:398:PRO:HD3	1.93	0.49
1:D:356:PHE:CE1	1:D:360:MET:HE1	2.48	0.49
1:B:397:GLN:HB3	1:B:398:PRO:HD3	1.94	0.49
1:A:263:ALA:HA	1:A:266:GLN:HB2	1.95	0.48
1:D:311:ILE:HG12	1:D:374:ILE:HG23	1.96	0.48
1:A:416:LEU:HD13	1:B:397:GLN:HE21	1.79	0.48
1:B:410:LYS:HG2	1:B:411:ARG:HG2	1.96	0.48
1:C:345:LEU:N	1:C:345:LEU:HD23	2.29	0.48
1:C:409:ILE:HG22	1:C:410:LYS:N	2.30	0.47
1:C:220:LEU:HD13	1:C:225:GLU:CG	2.43	0.47
1:C:328:THR:HG23	1:C:334:THR:HG22	1.96	0.47
1:C:330:LEU:O	1:C:331:LYS:C	2.50	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:403:LEU:HD11	1:C:421:MET:HE1	1.97	0.46
1:B:246:GLN:NE2	1:B:330:LEU:O	2.49	0.45
1:C:220:LEU:HD13	1:C:225:GLU:HG3	1.98	0.45
1:C:337:LYS:HG2	1:C:354:PHE:CD1	2.52	0.45
1:A:252:TRP:HE3	1:A:270:HIS:CD2	2.35	0.45
1:C:321:ASN:HD21	1:C:323:GLU:HB3	1.83	0.44
1:C:401:GLU:CG	1:D:416:LEU:HD21	2.48	0.44
1:D:417:ARG:O	1:D:421:MET:HG3	2.18	0.44
1:A:337:LYS:HG3	1:A:354:PHE:CD1	2.53	0.43
1:D:349:PHE:CE1	1:D:353:ILE:HD11	2.53	0.43
1:C:423:MET:HE1	1:D:422:LEU:HD22	2.01	0.43
1:C:380:SER:O	1:C:383:ARG:HG2	2.19	0.43
1:D:314:LEU:HD13	1:D:428:LEU:HD11	2.01	0.42
1:C:340:PHE:HB3	1:C:350:ILE:HG12	2.01	0.42
1:D:351:ASN:HB2	1:D:352:PRO:HD3	2.01	0.42
1:D:282:ILE:HG12	1:D:311:ILE:HG21	2.01	0.41
1:C:363:LEU:HD23	1:C:363:LEU:HA	1.93	0.41
1:A:252:TRP:CE3	1:A:270:HIS:CD2	3.09	0.41
1:A:275:ALA:O	1:A:279:VAL:HG23	2.20	0.41
1:B:275:ALA:O	1:B:279:VAL:HG23	2.20	0.41
1:A:408:ARG:O	1:A:412:PRO:HA	2.21	0.41
1:B:349:PHE:CZ	1:B:353:ILE:HD11	2.55	0.41
1:B:436:SER:HA	1:B:439:VAL:HG22	2.02	0.40
1:A:363:LEU:HA	1:A:363:LEU:HD23	1.96	0.40
1:B:416:LEU:C	1:B:419:PRO:HD2	2.41	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	228/253 (90%)	221 (97%)	6 (3%)	1 (0%)	39 74

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	234/253 (92%)	227 (97%)	5 (2%)	2 (1%)	21	55
1	C	225/253 (89%)	216 (96%)	7 (3%)	2 (1%)	21	55
1	D	209/253 (83%)	202 (97%)	7 (3%)	0	100	100
All	All	896/1012 (88%)	866 (97%)	25 (3%)	5 (1%)	30	65

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	446	ASP
1	C	410	LYS
1	B	331	LYS
1	B	257	ASP
1	C	253	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/222 (93%)	185 (90%)	21 (10%)	9	26
1	B	208/222 (94%)	191 (92%)	17 (8%)	14	38
1	C	205/222 (92%)	188 (92%)	17 (8%)	14	38
1	D	191/222 (86%)	170 (89%)	21 (11%)	8	23
All	All	810/888 (91%)	734 (91%)	76 (9%)	11	31

All (76) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	220	LEU
1	A	241	ARG
1	A	249	VAL
1	A	265	GLN
1	A	266	GLN
1	A	272	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	298	GLU
1	A	314	LEU
1	A	318	ARG
1	A	325	GLU
1	A	330	LEU
1	A	331	LYS
1	A	334	THR
1	A	337	LYS
1	A	353	ILE
1	A	355	GLU
1	A	417	ARG
1	A	431	LEU
1	A	443	ARG
1	A	447	LYS
1	A	454	SER
1	B	227	MET
1	B	235	GLN
1	B	236	LEU
1	B	242	SER
1	B	244	SER
1	B	257	ASP
1	B	264	ARG
1	B	265	GLN
1	B	278	SER
1	B	314	LEU
1	B	319	ARG
1	B	327	ILE
1	B	331	LYS
1	B	332	ASP
1	B	417	ARG
1	B	431	LEU
1	B	445	GLN
1	C	220	LEU
1	C	227	MET
1	C	235	GLN
1	C	242	SER
1	C	243	PHE
1	C	259	GLN
1	C	261	ARG
1	C	264	ARG
1	C	272	THR
1	C	287	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	314	LEU
1	C	330	LEU
1	C	347	VAL
1	C	373	LEU
1	C	393	GLU
1	C	408	ARG
1	C	447	LYS
1	D	239	ASN
1	D	242	SER
1	D	264	ARG
1	D	280	GLN
1	D	287	LYS
1	D	294	GLN
1	D	298	GLU
1	D	301	ILE
1	D	307	SER
1	D	321	ASN
1	D	342	ARG
1	D	348	GLU
1	D	408	ARG
1	D	411	ARG
1	D	413	GLN
1	D	423	MET
1	D	439	VAL
1	D	442	LEU
1	D	443	ARG
1	D	448	LYS
1	D	455	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	237	GLN
1	A	266	GLN
1	A	396	GLN
1	B	239	ASN
1	B	246	GLN
1	B	397	GLN
1	C	259	GLN
1	C	321	ASN
1	C	346	GLN
1	D	235	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	266	GLN
1	D	294	GLN
1	D	321	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	444	A	1500	-	32,32,32	2.66	8 (25%)	51,52,52	2.41	12 (23%)
2	444	B	2500	-	32,32,32	2.56	8 (25%)	51,52,52	2.30	10 (19%)
2	444	C	3500	-	32,32,32	2.44	8 (25%)	51,52,52	1.96	11 (21%)
2	444	D	4500	-	32,32,32	2.06	7 (21%)	51,52,52	1.75	7 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	444	A	1500	-	-	0/44/45/45	0/2/2/2
2	444	B	2500	-	-	0/44/45/45	0/2/2/2
2	444	C	3500	-	-	0/44/45/45	0/2/2/2
2	444	D	4500	-	-	0/44/45/45	0/2/2/2

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1500	444	C05-C04	2.14	1.43	1.38
2	C	3500	444	C05-C04	2.22	1.43	1.38
2	B	2500	444	C33-C26	2.31	1.57	1.53
2	B	2500	444	C28-C23	2.33	1.43	1.39
2	B	2500	444	C25-C26	2.34	1.42	1.39
2	D	4500	444	C28-C23	2.36	1.43	1.39
2	A	1500	444	C28-C23	2.42	1.44	1.39
2	D	4500	444	O42-C33	2.43	1.47	1.42
2	A	1500	444	C27-C26	2.51	1.43	1.39
2	A	1500	444	C25-C26	2.60	1.43	1.39
2	C	3500	444	C28-C23	2.68	1.44	1.39
2	B	2500	444	C02-C01	2.74	1.43	1.38
2	C	3500	444	C27-C26	2.77	1.43	1.39
2	A	1500	444	C16-N15	2.80	1.53	1.48
2	D	4500	444	C16-N15	2.81	1.53	1.48
2	D	4500	444	C27-C26	2.82	1.43	1.39
2	B	2500	444	C27-C26	3.19	1.44	1.39
2	D	4500	444	C02-C01	3.35	1.44	1.38
2	A	1500	444	C02-C01	3.37	1.44	1.38
2	B	2500	444	C16-N15	3.42	1.54	1.48
2	C	3500	444	C02-C01	3.48	1.44	1.38
2	D	4500	444	C25-C26	3.50	1.44	1.39
2	C	3500	444	C23-N15	3.57	1.49	1.44
2	C	3500	444	C25-C26	3.58	1.44	1.39
2	C	3500	444	C16-N15	4.05	1.55	1.48
2	B	2500	444	C23-N15	4.40	1.50	1.44
2	A	1500	444	C23-N15	4.82	1.50	1.44
2	D	4500	444	S12-N15	7.35	1.75	1.64
2	C	3500	444	S12-N15	9.43	1.77	1.64
2	B	2500	444	S12-N15	10.92	1.80	1.64
2	A	1500	444	S12-N15	11.45	1.80	1.64

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2500	444	C16-N15-C23	-6.92	109.96	117.34
2	C	3500	444	C16-N15-C23	-6.18	110.76	117.34
2	B	2500	444	C38-C33-C34	-6.13	105.20	110.55
2	A	1500	444	O14-S12-O13	-6.10	108.94	119.47
2	A	1500	444	C16-N15-C23	-5.95	110.99	117.34
2	D	4500	444	C38-C33-C34	-5.92	105.38	110.55
2	D	4500	444	O13-S12-C01	-4.81	101.76	108.00
2	C	3500	444	C38-C33-C34	-4.64	106.49	110.55
2	A	1500	444	C38-C33-C34	-4.62	106.51	110.55
2	A	1500	444	O13-S12-C01	-4.52	102.14	108.00
2	B	2500	444	O13-S12-C01	-4.28	102.44	108.00
2	C	3500	444	O14-S12-O13	-3.87	112.79	119.47
2	B	2500	444	O14-S12-O13	-3.77	112.96	119.47
2	D	4500	444	C16-N15-C23	-3.38	113.74	117.34
2	C	3500	444	C01-S12-N15	-3.06	102.04	106.81
2	A	1500	444	C38-C33-C26	-2.94	106.25	110.88
2	D	4500	444	C23-N15-S12	-2.93	112.03	118.09
2	A	1500	444	F22-C19-C16	-2.91	106.97	112.08
2	B	2500	444	C01-S12-N15	-2.50	102.91	106.81
2	C	3500	444	C38-C33-C26	-2.39	107.10	110.88
2	D	4500	444	C24-C23-N15	-2.39	116.72	120.34
2	C	3500	444	C24-C23-N15	-2.33	116.81	120.34
2	D	4500	444	F22-C19-C16	-2.31	108.02	112.08
2	B	2500	444	C24-C23-N15	-2.29	116.87	120.34
2	A	1500	444	C16-N15-S12	-2.02	113.48	117.50
2	C	3500	444	C27-C26-C33	2.51	126.25	120.52
2	B	2500	444	C27-C26-C33	2.52	126.27	120.52
2	C	3500	444	F41-C38-C33	2.61	117.34	111.92
2	A	1500	444	C27-C26-C33	2.63	126.53	120.52
2	C	3500	444	C28-C23-N15	2.81	124.61	120.34
2	A	1500	444	F41-C38-C33	2.84	117.82	111.92
2	A	1500	444	C28-C23-N15	2.91	124.76	120.34
2	B	2500	444	C28-C23-N15	3.32	125.38	120.34
2	C	3500	444	O13-S12-N15	4.23	112.24	106.78
2	D	4500	444	O13-S12-N15	4.69	112.84	106.78
2	B	2500	444	O13-S12-N15	5.07	113.32	106.78
2	A	1500	444	O14-S12-N15	5.44	113.80	106.78
2	C	3500	444	O14-S12-N15	5.84	114.31	106.78
2	B	2500	444	O14-S12-N15	7.70	116.72	106.78
2	A	1500	444	O13-S12-N15	8.29	117.47	106.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/253 (91%)	0.19	12 (5%) 31 20	13, 25, 36, 42	0
1	B	238/253 (94%)	0.06	5 (2%) 67 56	12, 23, 44, 57	0
1	C	231/253 (91%)	0.10	7 (3%) 54 41	13, 24, 39, 62	0
1	D	219/253 (86%)	0.41	25 (11%) 7 3	16, 23, 34, 48	0
All	All	920/1012 (90%)	0.19	49 (5%) 30 20	12, 23, 39, 62	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	241	ARG	5.5
1	A	322	HIS	4.1
1	D	262	ASP	3.9
1	D	344	GLY	3.8
1	D	263	ALA	3.8
1	A	444	LEU	3.7
1	D	252	TRP	3.7
1	D	451	PRO	3.6
1	D	265	GLN	3.6
1	D	442	LEU	3.4
1	A	264	ARG	3.4
1	A	446	ASP	3.3
1	A	222	ALA	3.2
1	D	240	LYS	3.2
1	C	331	LYS	3.2
1	C	413	GLN	3.1
1	D	266	GLN	3.1
1	B	446	ASP	3.1
1	A	324	THR	3.1
1	D	345	LEU	3.0
1	C	408	ARG	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	413	GLN	3.0
1	D	452	LEU	3.0
1	D	249	VAL	2.9
1	D	440	PHE	2.8
1	D	242	SER	2.7
1	D	271	PHE	2.7
1	B	247	PRO	2.7
1	C	332	ASP	2.6
1	D	264	ARG	2.5
1	D	260	SER	2.4
1	C	220	LEU	2.4
1	A	447	LYS	2.4
1	B	243	PHE	2.4
1	C	259	GLN	2.3
1	D	343	ALA	2.3
1	B	330	LEU	2.3
1	D	341	HIS	2.2
1	D	268	PHE	2.2
1	D	441	ALA	2.2
1	B	333	PHE	2.2
1	C	329	PHE	2.2
1	A	440	PHE	2.2
1	D	415	GLN	2.1
1	A	341	HIS	2.1
1	A	413	GLN	2.1
1	A	262	ASP	2.1
1	D	412	PRO	2.0
1	A	326	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	444	D	4500	31/31	0.96	0.27	0.07	15,27,47,47	0
2	444	C	3500	31/31	0.97	0.20	0.03	19,27,51,51	0
2	444	B	2500	31/31	0.95	0.19	-0.61	18,27,51,52	0
2	444	A	1500	31/31	0.95	0.17	-0.97	14,22,36,38	0

## 6.5 Other polymers

There are no such residues in this entry.