



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:58 PM GMT

PDB ID : 6PTD  
Title : PHOSPHATIDYLINOSITOL-SPECIFIC PHOSPHOLIPASE C MUTANT  
H32L  
Authors : Heinz, D.W.  
Deposited on : 1997-07-18  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2506 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHATIDYLINOSITOL-SPECIFIC PHOSPHOLIPASE C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	0	0	0
			2420	1548	398	467	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	LEU	HIS	ENGINEERED	UNP P14262


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	86	Total	O	0	0
			86	86		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PHOSPHATIDYLINOSITOL-SPECIFIC PHOSPHOLIPASE C



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.20 Å 46.00 Å 161.20 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	18.90 – 2.60 18.72 – 2.60	Depositor EDS
% Data completeness (in resolution range)	83.5 (18.90-2.60) 83.6 (18.72-2.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.46 (at 2.59 Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.182 , 0.294 0.184 , 0.284	Depositor DCC
$R_{free}$ test set	434 reflections (4.77%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtriage
Anisotropy	0.596	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 78.0	EDS
Estimated twinning fraction	0.029 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 9104 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2506	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/2483	0.63	1/3369 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	LEU	CA-CB-CG	5.45	127.83	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2420	0	2344	87	0
2	A	86	0	0	5	0
All	All	2506	0	2344	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:VAL:HG22	1:A:194:VAL:HG22	1.60	0.84

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:GLU:HA	1:A:199:LYS:HD3	1.61	0.81
1:A:183:THR:HG22	1:A:197:GLN:HG2	1.63	0.80
1:A:6:GLU:HB2	1:A:16:PRO:HG3	1.65	0.78
1:A:69:ARG:HB3	1:A:117:GLU:HB2	1.67	0.75
1:A:71:ARG:HG3	1:A:118:TYR:HB3	1.66	0.73
1:A:28:ILE:HG21	1:A:64:ARG:HB2	1.71	0.72
1:A:116:LYS:HE3	1:A:164:TYR:HB2	1.74	0.69
1:A:188:VAL:HG22	1:A:194:VAL:CG2	2.22	0.69
1:A:221:ASN:HD21	1:A:224:ASP:HB2	1.59	0.67
1:A:160:LEU:CD1	1:A:162:LYS:HG2	2.25	0.66
1:A:18:PRO:HB2	1:A:21:ILE:HD12	1.78	0.65
1:A:194:VAL:HG13	1:A:228:LEU:HB3	1.81	0.62
1:A:24:ALA:HB2	1:A:228:LEU:HD22	1.82	0.61
1:A:252:ILE:O	1:A:256:ILE:HG12	2.00	0.61
1:A:185:THR:HA	1:A:194:VAL:O	2.01	0.61
1:A:204:TYR:O	1:A:208:VAL:HG23	2.01	0.60
1:A:54:ASP:O	1:A:58:GLN:HG3	2.02	0.60
1:A:182:GLU:HB2	2:A:362:HOH:O	2.04	0.58
1:A:238:GLY:O	1:A:243:ASN:HB3	2.05	0.57
1:A:129:SER:O	1:A:133:GLU:HG3	2.05	0.56
1:A:28:ILE:CG2	1:A:64:ARG:HB2	2.36	0.55
1:A:269:GLY:HA2	2:A:368:HOH:O	2.06	0.55
1:A:186:THR:HG22	1:A:187:THR:H	1.72	0.55
1:A:228:LEU:HD11	1:A:270:TRP:CZ3	2.42	0.53
1:A:215:MET:O	1:A:219:MET:HG3	2.07	0.53
1:A:260:ILE:O	1:A:265:PRO:HD3	2.08	0.53
1:A:36:THR:HB	1:A:49:MET:HA	1.90	0.52
1:A:41:ASN:O	1:A:45:GLN:HG3	2.10	0.52
1:A:73:THR:HA	1:A:121:MET:SD	2.50	0.52
1:A:188:VAL:CG2	1:A:194:VAL:HG22	2.36	0.51
1:A:242:TRP:CE3	1:A:242:TRP:HA	2.45	0.51
1:A:207:LYS:HG2	1:A:235:LEU:HG	1.93	0.51
1:A:160:LEU:HD11	1:A:162:LYS:HG2	1.92	0.50
1:A:257:ALA:CB	1:A:282:PRO:HB2	2.42	0.50
1:A:5:ASN:H	1:A:5:ASN:HD22	1.60	0.50
1:A:242:TRP:HE3	1:A:242:TRP:HA	1.75	0.50
1:A:184:PHE:CZ	1:A:196:VAL:HB	2.47	0.50
1:A:39:LEU:HD12	1:A:49:MET:H	1.78	0.49
1:A:253:ASN:HB2	1:A:254:PRO:HD3	1.92	0.49
1:A:116:LYS:HB3	2:A:358:HOH:O	2.12	0.49
1:A:12:LYS:HB3	1:A:12:LYS:NZ	2.28	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:LEU:HD11	1:A:270:TRP:CH2	2.48	0.48
1:A:39:LEU:HD22	1:A:44:LYS:HD3	1.95	0.48
1:A:221:ASN:ND2	1:A:224:ASP:HB2	2.25	0.48
1:A:257:ALA:HB3	1:A:282:PRO:HB2	1.95	0.48
1:A:188:VAL:HB	1:A:189:ASN:H	1.37	0.48
1:A:176:PHE:CD2	1:A:188:VAL:HG13	2.48	0.48
1:A:174:ASN:HD22	1:A:174:ASN:N	2.11	0.48
1:A:267:ARG:HD2	2:A:368:HOH:O	2.13	0.47
1:A:6:GLU:HB2	1:A:16:PRO:CG	2.42	0.47
1:A:132:PHE:CE2	1:A:137:PHE:HB2	2.50	0.47
1:A:160:LEU:HD13	1:A:162:LYS:HG2	1.96	0.46
1:A:21:ILE:HA	1:A:22:PRO:HD3	1.78	0.46
1:A:5:ASN:HD22	1:A:5:ASN:N	2.13	0.46
1:A:183:THR:HG23	1:A:199:LYS:HE2	1.97	0.46
1:A:22:PRO:HD2	1:A:225:LEU:HD22	1.98	0.46
1:A:133:GLU:HA	1:A:137:PHE:HB3	1.96	0.46
1:A:25:ARG:HB2	1:A:225:LEU:HD23	1.98	0.46
1:A:181:ASN:O	1:A:182:GLU:HB3	2.16	0.45
1:A:69:ARG:O	1:A:80:LEU:HA	2.17	0.45
1:A:24:ALA:CB	1:A:228:LEU:HD22	2.46	0.45
1:A:46:VAL:HG23	1:A:47:TRP:CD1	2.52	0.45
1:A:41:ASN:ND2	1:A:44:LYS:HD2	2.32	0.45
1:A:12:LYS:HA	1:A:108:GLU:CD	2.37	0.44
1:A:5:ASN:ND2	1:A:6:GLU:HG2	2.33	0.44
1:A:176:PHE:HD2	1:A:188:VAL:HG13	1.82	0.44
1:A:15:GLN:NE2	1:A:107:SER:HB3	2.33	0.44
1:A:236:SER:O	1:A:237:SER:C	2.56	0.44
1:A:139:ASP:OD1	1:A:141:ILE:HG13	2.18	0.43
1:A:83:GLY:H	1:A:117:GLU:CG	2.31	0.43
1:A:240:THR:HG22	1:A:241:ALA:N	2.33	0.43
1:A:224:ASP:O	1:A:267:ARG:NH2	2.52	0.43
1:A:251:TYR:O	1:A:255:GLU:HB2	2.19	0.43
1:A:70:GLY:HA2	1:A:79:VAL:O	2.20	0.42
1:A:95:ILE:HG22	1:A:99:LYS:HE3	2.00	0.42
1:A:214:THR:HB	1:A:231:ASN:HD21	1.83	0.42
1:A:28:ILE:HG22	1:A:29:PRO:O	2.19	0.42
1:A:281:SER:CB	1:A:282:PRO:HD3	2.50	0.42
1:A:286:GLN:HG3	2:A:378:HOH:O	2.19	0.42
1:A:260:ILE:HG23	1:A:268:VAL:HG21	2.02	0.41
1:A:95:ILE:O	1:A:99:LYS:HG3	2.21	0.41
1:A:10:TRP:O	1:A:13:TRP:HD1	2.04	0.41

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:SER:HB3	1:A:248:TYR:HE2	1.85	0.40
1:A:174:ASN:N	1:A:174:ASN:ND2	2.70	0.40
1:A:272:ILE:N	1:A:272:ILE:HD12	2.37	0.40
1:A:18:PRO:HB2	1:A:21:ILE:CD1	2.50	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/298 (99%)	253 (86%)	34 (12%)	7 (2%)	7	13

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	177	TYR
1	A	188	VAL
1	A	237	SER
1	A	238	GLY
1	A	49	MET
1	A	189	ASN
1	A	275	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	268/270 (99%)	239 (89%)	29 (11%)	8 15

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	6	GLU
1	A	12	LYS
1	A	19	ASP
1	A	33	ASP
1	A	52	GLU
1	A	69	ARG
1	A	72	LEU
1	A	81	HIS
1	A	91	LEU
1	A	100	GLN
1	A	114	LEU
1	A	116	LYS
1	A	119	GLU
1	A	129	SER
1	A	141	ILE
1	A	148	ASN
1	A	153	ASP
1	A	187	THR
1	A	188	VAL
1	A	194	VAL
1	A	202	VAL
1	A	218	THR
1	A	228	LEU
1	A	242	TRP
1	A	243	ASN
1	A	281	SER
1	A	284	LEU
1	A	296	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	15	GLN
1	A	41	ASN
1	A	100	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	148	ASN
1	A	174	ASN
1	A	191	ASN
1	A	197	GLN
1	A	221	ASN
1	A	227	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	296/298 (99%)	-0.62	6 (2%) 68 63	3, 22, 65, 104	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	GLY	3.4
1	A	191	ASN	3.0
1	A	1	ALA	2.8
1	A	242	TRP	2.6
1	A	240	THR	2.2
1	A	241	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.