



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:45 PM GMT

PDB ID : 4PU3
Title : Shewanella oneidensis MR-1 Toxin Antitoxin System HipA, HipB and its operator DNA complex (space group P212121)
Authors : Wen, Y.; Behiels, E.; Felix, J.; Elegheert, J.; Vergauwen, B.; Devreese, B.; Savvides, S.
Deposited on : 2014-03-12
Resolution : 3.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1476 (3.50-3.30)
Clashscore	102246	1611 (3.50-3.30)
Ramachandran outliers	100387	1571 (3.50-3.30)
Sidechain outliers	100360	1571 (3.50-3.30)
RSRZ outliers	91569	1485 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	454	<div> <div>9%</div> <div>87%</div> <div>7%</div> <div>6%</div> </div>
1	B	454	<div> <div>7%</div> <div>90%</div> <div>•</div> <div>6%</div> </div>
2	C	118	<div> <div>4%</div> <div>54%</div> <div>• •</div> <div>40%</div> </div>
2	D	118	<div> <div>3%</div> <div>58%</div> <div>•</div> <div>40%</div> </div>
3	Q	26	<div> <div>4%</div> <div>85%</div> <div>8%</div> <div>• •</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	P	26	 A horizontal bar chart showing the quality of chain 4. The bar is divided into three segments: a green segment representing 85%, a yellow segment representing 8%, and a grey segment representing 8%. The percentages are labeled below the corresponding segments.

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 17163 atoms, of which 8358 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toxin-antitoxin system toxin HipA family.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	426	Total	C	H	N	O	P	S	0	0	0
			6706	2147	3342	572	626	1	18			
1	B	426	Total	C	H	N	O	P	S	0	0	0
			6663	2143	3308	573	621	1	17			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP Q8EIX3
A	-19	GLY	-	EXPRESSION TAG	UNP Q8EIX3
A	-18	SER	-	EXPRESSION TAG	UNP Q8EIX3
A	-17	SER	-	EXPRESSION TAG	UNP Q8EIX3
A	-16	HIS	-	EXPRESSION TAG	UNP Q8EIX3
A	-15	HIS	-	EXPRESSION TAG	UNP Q8EIX3
A	-14	HIS	-	EXPRESSION TAG	UNP Q8EIX3
A	-13	HIS	-	EXPRESSION TAG	UNP Q8EIX3
A	-12	HIS	-	EXPRESSION TAG	UNP Q8EIX3
A	-11	HIS	-	EXPRESSION TAG	UNP Q8EIX3
A	-10	SER	-	EXPRESSION TAG	UNP Q8EIX3
A	-9	SER	-	EXPRESSION TAG	UNP Q8EIX3
A	-8	GLY	-	EXPRESSION TAG	UNP Q8EIX3
A	-7	LEU	-	EXPRESSION TAG	UNP Q8EIX3
A	-6	VAL	-	EXPRESSION TAG	UNP Q8EIX3
A	-5	PRO	-	EXPRESSION TAG	UNP Q8EIX3
A	-4	ARG	-	EXPRESSION TAG	UNP Q8EIX3
A	-3	GLY	-	EXPRESSION TAG	UNP Q8EIX3
A	-2	SER	-	EXPRESSION TAG	UNP Q8EIX3
A	-1	HIS	-	EXPRESSION TAG	UNP Q8EIX3
A	0	MET	-	EXPRESSION TAG	UNP Q8EIX3
B	-20	MET	-	EXPRESSION TAG	UNP Q8EIX3
B	-19	GLY	-	EXPRESSION TAG	UNP Q8EIX3
B	-18	SER	-	EXPRESSION TAG	UNP Q8EIX3
B	-17	SER	-	EXPRESSION TAG	UNP Q8EIX3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	EXPRESSION TAG	UNP Q8EIX3
B	-15	HIS	-	EXPRESSION TAG	UNP Q8EIX3
B	-14	HIS	-	EXPRESSION TAG	UNP Q8EIX3
B	-13	HIS	-	EXPRESSION TAG	UNP Q8EIX3
B	-12	HIS	-	EXPRESSION TAG	UNP Q8EIX3
B	-11	HIS	-	EXPRESSION TAG	UNP Q8EIX3
B	-10	SER	-	EXPRESSION TAG	UNP Q8EIX3
B	-9	SER	-	EXPRESSION TAG	UNP Q8EIX3
B	-8	GLY	-	EXPRESSION TAG	UNP Q8EIX3
B	-7	LEU	-	EXPRESSION TAG	UNP Q8EIX3
B	-6	VAL	-	EXPRESSION TAG	UNP Q8EIX3
B	-5	PRO	-	EXPRESSION TAG	UNP Q8EIX3
B	-4	ARG	-	EXPRESSION TAG	UNP Q8EIX3
B	-3	GLY	-	EXPRESSION TAG	UNP Q8EIX3
B	-2	SER	-	EXPRESSION TAG	UNP Q8EIX3
B	-1	HIS	-	EXPRESSION TAG	UNP Q8EIX3
B	0	MET	-	EXPRESSION TAG	UNP Q8EIX3

- Molecule 2 is a protein called Toxin-antitoxin system antidote transcriptional repressor Xre family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	D	71	Total	C	H	N	O	S	0	0	0
			1118	347	577	91	100	3			
2	C	71	Total	C	H	N	O	S	0	0	0
			1118	347	577	91	100	3			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-20	MET	-	EXPRESSION TAG	UNP Q8EIX4
D	-19	GLY	-	EXPRESSION TAG	UNP Q8EIX4
D	-18	SER	-	EXPRESSION TAG	UNP Q8EIX4
D	-17	SER	-	EXPRESSION TAG	UNP Q8EIX4
D	-16	HIS	-	EXPRESSION TAG	UNP Q8EIX4
D	-15	HIS	-	EXPRESSION TAG	UNP Q8EIX4
D	-14	HIS	-	EXPRESSION TAG	UNP Q8EIX4
D	-13	HIS	-	EXPRESSION TAG	UNP Q8EIX4
D	-12	HIS	-	EXPRESSION TAG	UNP Q8EIX4
D	-11	HIS	-	EXPRESSION TAG	UNP Q8EIX4
D	-10	SER	-	EXPRESSION TAG	UNP Q8EIX4
D	-9	SER	-	EXPRESSION TAG	UNP Q8EIX4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-8	GLY	-	EXPRESSION TAG	UNP Q8EIX4
D	-7	LEU	-	EXPRESSION TAG	UNP Q8EIX4
D	-6	VAL	-	EXPRESSION TAG	UNP Q8EIX4
D	-5	PRO	-	EXPRESSION TAG	UNP Q8EIX4
D	-4	ARG	-	EXPRESSION TAG	UNP Q8EIX4
D	-3	GLY	-	EXPRESSION TAG	UNP Q8EIX4
D	-2	SER	-	EXPRESSION TAG	UNP Q8EIX4
D	-1	HIS	-	EXPRESSION TAG	UNP Q8EIX4
D	0	MET	-	EXPRESSION TAG	UNP Q8EIX4
D	1	MET	-	EXPRESSION TAG	UNP Q8EIX4
D	2	ASN	-	EXPRESSION TAG	UNP Q8EIX4
D	3	GLY	-	EXPRESSION TAG	UNP Q8EIX4
D	4	THR	-	EXPRESSION TAG	UNP Q8EIX4
D	5	ASP	-	EXPRESSION TAG	UNP Q8EIX4
D	6	ILE	-	EXPRESSION TAG	UNP Q8EIX4
D	7	LYS	-	EXPRESSION TAG	UNP Q8EIX4
D	8	ALA	-	EXPRESSION TAG	UNP Q8EIX4
D	9	LYS	-	EXPRESSION TAG	UNP Q8EIX4
D	10	VAL	-	EXPRESSION TAG	UNP Q8EIX4
D	11	TYR	-	EXPRESSION TAG	UNP Q8EIX4
D	12	GLU	-	EXPRESSION TAG	UNP Q8EIX4
D	13	ASP	-	EXPRESSION TAG	UNP Q8EIX4
D	14	THR	-	EXPRESSION TAG	UNP Q8EIX4
D	15	LEU	-	EXPRESSION TAG	UNP Q8EIX4
D	16	LEU	-	EXPRESSION TAG	UNP Q8EIX4
D	17	GLU	-	EXPRESSION TAG	UNP Q8EIX4
D	18	THR	-	EXPRESSION TAG	UNP Q8EIX4
D	19	ILE	-	EXPRESSION TAG	UNP Q8EIX4
C	-20	MET	-	EXPRESSION TAG	UNP Q8EIX4
C	-19	GLY	-	EXPRESSION TAG	UNP Q8EIX4
C	-18	SER	-	EXPRESSION TAG	UNP Q8EIX4
C	-17	SER	-	EXPRESSION TAG	UNP Q8EIX4
C	-16	HIS	-	EXPRESSION TAG	UNP Q8EIX4
C	-15	HIS	-	EXPRESSION TAG	UNP Q8EIX4
C	-14	HIS	-	EXPRESSION TAG	UNP Q8EIX4
C	-13	HIS	-	EXPRESSION TAG	UNP Q8EIX4
C	-12	HIS	-	EXPRESSION TAG	UNP Q8EIX4
C	-11	HIS	-	EXPRESSION TAG	UNP Q8EIX4
C	-10	SER	-	EXPRESSION TAG	UNP Q8EIX4
C	-9	SER	-	EXPRESSION TAG	UNP Q8EIX4
C	-8	GLY	-	EXPRESSION TAG	UNP Q8EIX4
C	-7	LEU	-	EXPRESSION TAG	UNP Q8EIX4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-6	VAL	-	EXPRESSION TAG	UNP Q8EIX4
C	-5	PRO	-	EXPRESSION TAG	UNP Q8EIX4
C	-4	ARG	-	EXPRESSION TAG	UNP Q8EIX4
C	-3	GLY	-	EXPRESSION TAG	UNP Q8EIX4
C	-2	SER	-	EXPRESSION TAG	UNP Q8EIX4
C	-1	HIS	-	EXPRESSION TAG	UNP Q8EIX4
C	0	MET	-	EXPRESSION TAG	UNP Q8EIX4
C	1	MET	-	EXPRESSION TAG	UNP Q8EIX4
C	2	ASN	-	EXPRESSION TAG	UNP Q8EIX4
C	3	GLY	-	EXPRESSION TAG	UNP Q8EIX4
C	4	THR	-	EXPRESSION TAG	UNP Q8EIX4
C	5	ASP	-	EXPRESSION TAG	UNP Q8EIX4
C	6	ILE	-	EXPRESSION TAG	UNP Q8EIX4
C	7	LYS	-	EXPRESSION TAG	UNP Q8EIX4
C	8	ALA	-	EXPRESSION TAG	UNP Q8EIX4
C	9	LYS	-	EXPRESSION TAG	UNP Q8EIX4
C	10	VAL	-	EXPRESSION TAG	UNP Q8EIX4
C	11	TYR	-	EXPRESSION TAG	UNP Q8EIX4
C	12	GLU	-	EXPRESSION TAG	UNP Q8EIX4
C	13	ASP	-	EXPRESSION TAG	UNP Q8EIX4
C	14	THR	-	EXPRESSION TAG	UNP Q8EIX4
C	15	LEU	-	EXPRESSION TAG	UNP Q8EIX4
C	16	LEU	-	EXPRESSION TAG	UNP Q8EIX4
C	17	GLU	-	EXPRESSION TAG	UNP Q8EIX4
C	18	THR	-	EXPRESSION TAG	UNP Q8EIX4
C	19	ILE	-	EXPRESSION TAG	UNP Q8EIX4


- Molecule 3 is a DNA chain called Operator DNA.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	Q	25	Total	C	H	N	O	P	0	0	0
			799	247	282	101	144	25			

- Molecule 4 is a DNA chain called Operator DNA.



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	P	24	Total	C	H	N	O	P	0	0	0
			759	235	272	79	149	24			

- Molecule 1: Toxin-antitoxin system toxin HipA family






● Molecule 3: Operator DNA



● Molecule 4: Operator DNA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.82Å 122.45Å 189.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.38 – 3.39 49.38 – 3.39	Depositor EDS
% Data completeness (in resolution range)	98.8 (49.38-3.39) 98.9 (49.38-3.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 3.40Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1287)	Depositor
R, R_{free}	0.256 , 0.315 0.255 , 0.321	Depositor DCC
R_{free} test set	959 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	75.0	Xtriage
Anisotropy	0.496	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 28.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	1 of 19211 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	17163	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.72 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.2648e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/3426	0.42	0/4638
1	B	0.23	0/3420	0.41	0/4635
2	C	0.22	0/545	0.39	0/732
2	D	0.22	0/545	0.40	0/732
3	Q	0.48	0/582	1.14	2/896 (0.2%)
4	P	0.47	0/543	1.27	2/835 (0.2%)
All	All	0.27	0/9061	0.59	4/12468 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	P	6	DG	O4'-C1'-N9	6.22	112.35	108.00
4	P	6	DG	C1'-O4'-C4'	-5.89	104.21	110.10
3	Q	14	DA	O4'-C1'-N9	5.53	111.87	108.00
3	Q	6	DT	O4'-C4'-C3'	-5.25	102.40	104.50

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	112	LEU	Peptide
1	A	131	ASP	Peptide
1	B	137	MET	Peptide
1	B	143	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3364	3342	3332	16	0
1	B	3355	3308	3298	10	0
2	C	541	577	577	4	0
2	D	541	577	577	2	0
3	Q	517	282	282	2	0
4	P	487	272	273	1	0
All	All	8805	8358	8339	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:GLN:OE1	1:A:380:ARG:NH2	2.23	0.72
1:B:146:LEU:CB	1:B:147:SEP:HA	2.24	0.67
2:D:55:LYS:HE2	3:Q:6:DT:H72	1.78	0.65
1:A:335:TYR:O	1:A:340:GLN:NE2	2.31	0.64
2:C:36:ARG:NH1	2:C:79:LEU:O	2.31	0.63
2:C:20:MET:N	2:C:20:MET:SD	2.70	0.63
1:B:335:TYR:O	1:B:340:GLN:NE2	2.34	0.60
1:A:99:THR:O	1:A:224:ARG:NH2	2.35	0.59
1:B:137:MET:HB3	1:B:138:GLU:HB2	1.84	0.58
1:A:132:PRO:HA	1:A:133:THR:C	2.23	0.58
1:B:146:LEU:CB	1:B:147:SEP:CA	2.84	0.55
1:A:131:ASP:HB3	1:A:132:PRO:CD	2.37	0.55
1:B:290:GLN:OE1	1:B:377:ASN:ND2	2.39	0.54
1:B:269:GLY:O	1:B:274:ARG:NH1	2.42	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:ASP:HB2	1:A:142:GLY:HA3	1.92	0.51
1:A:145:ARG:NH1	1:A:339:GLU:OE1	2.44	0.50
3:Q:15:DG:N2	4:P:12:DT:O2	2.45	0.49
1:A:132:PRO:HB3	1:A:134:MET:HA	1.93	0.49
1:A:138:GLU:OE2	1:A:138:GLU:N	2.43	0.49
1:A:162:LYS:NZ	1:A:163:GLU:OE2	2.47	0.48
1:A:12:HIS:CE1	1:A:106:VAL:HG11	2.48	0.48
2:C:37:ARG:NE	2:C:62:GLU:OE2	2.47	0.47
1:B:137:MET:HB3	1:B:138:GLU:CB	2.45	0.47
1:A:269:GLY:O	1:A:274:ARG:NH1	2.48	0.46
1:A:221:ARG:NH2	1:A:223:ASP:OD2	2.46	0.45
2:D:37:ARG:NE	2:D:62:GLU:OE2	2.50	0.45
1:B:359:GLN:OE1	1:B:359:GLN:N	2.47	0.44
1:B:221:ARG:NH2	1:B:223:ASP:OD2	2.50	0.44
2:C:22:SER:HB2	2:C:23:PRO:HD2	1.99	0.44
1:A:146:LEU:O	1:A:148:VAL:N	2.50	0.43
1:B:138:GLU:OE1	1:B:140:TRP:NE1	2.52	0.43
1:A:385:ILE:HB	1:A:386:PRO:HD3	2.03	0.40
1:A:38:TRP:NE1	1:A:43:PHE:O	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/454 (93%)	413 (98%)	10 (2%)	0	100	100
1	B	423/454 (93%)	413 (98%)	9 (2%)	1 (0%)	52	87
2	C	67/118 (57%)	67 (100%)	0	0	100	100
2	D	67/118 (57%)	67 (100%)	0	0	100	100
All	All	980/1144 (86%)	960 (98%)	19 (2%)	1 (0%)	56	89

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	148	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/386 (93%)	355 (99%)	4 (1%)	80	92
1	B	354/386 (92%)	352 (99%)	2 (1%)	90	96
2	C	59/101 (58%)	57 (97%)	2 (3%)	44	79
2	D	59/101 (58%)	59 (100%)	0	100	100
All	All	831/974 (85%)	823 (99%)	8 (1%)	82	93

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	131	ASP
1	A	145	ARG
1	A	151	VAL
1	A	290	GLN
1	B	131	ASP
1	B	290	GLN
2	C	20	MET
2	C	36	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	SEP	A	147	1	8,9,10	1.53	1 (12%)	8,12,14	1.41	2 (25%)
1	SEP	B	147	1	8,9,10	1.55	1 (12%)	8,12,14	1.46	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	147	1	-	0/6/8/10	0/0/0/0
1	SEP	B	147	1	-	0/6/8/10	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	147	SEP	P-O1P	3.17	1.61	1.51
1	B	147	SEP	P-O1P	3.21	1.61	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	147	SEP	O-C-CA	-2.15	119.90	125.49
1	A	147	SEP	OG-CB-CA	2.77	110.64	108.27
1	B	147	SEP	OG-CB-CA	3.18	110.99	108.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	147	SEP	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	425/454 (93%)	0.56	39 (9%)	11 11	9, 48, 115, 135	0
1	B	425/454 (93%)	0.59	34 (8%)	15 14	6, 34, 110, 150	0
2	C	71/118 (60%)	0.38	5 (7%)	19 18	6, 19, 47, 55	0
2	D	71/118 (60%)	0.35	3 (4%)	40 35	9, 21, 50, 70	0
3	Q	25/26 (96%)	0.35	1 (4%)	42 37	22, 56, 97, 123	0
4	P	24/26 (92%)	0.25	0	100 100	21, 56, 82, 95	0
All	All	1041/1196 (87%)	0.53	82 (7%)	15 14	6, 38, 111, 150	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	170	THR	8.1
1	B	137	MET	7.0
1	B	169	GLY	6.1
1	B	231	GLN	6.1
1	A	108	LYS	5.6
1	B	139	ILE	5.3
1	A	113	PRO	4.8
1	A	83	VAL	4.7
1	B	72	LYS	4.6
1	B	85	LYS	4.4
1	B	135	TRP	4.4
1	A	114	GLU	4.3
1	A	112	LEU	4.3
1	B	82	GLY	4.2
1	B	230	GLU	4.2
1	A	134	MET	4.2
1	A	26	THR	4.1
1	B	113	PRO	4.0
1	B	73	GLY	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	212	GLY	4.0
1	A	164	PHE	3.8
2	C	85	SER	3.7
1	A	215	LYS	3.6
1	A	77	LEU	3.6
1	B	83	VAL	3.5
1	A	82	GLY	3.5
1	A	432	GLU	3.3
1	B	116	GLN	3.3
2	D	95	GLY	3.3
2	D	84	VAL	3.3
1	B	229	GLY	3.1
1	A	85	LYS	3.0
1	A	209	VAL	2.9
1	A	72	LYS	2.9
1	B	213	ARG	2.9
1	B	84	SER	2.9
1	A	109	GLY	2.8
1	B	433	VAL	2.8
1	B	86	GLY	2.8
2	C	84	VAL	2.8
1	B	111	LEU	2.8
2	C	83	ILE	2.8
1	A	207	ASP	2.7
1	A	211	PHE	2.7
1	B	214	TYR	2.7
1	A	151	VAL	2.7
1	A	132	PRO	2.7
1	A	103	ILE	2.6
1	A	84	SER	2.6
1	A	120	ILE	2.6
1	A	163	GLU	2.6
1	B	211	PHE	2.5
1	A	162	LYS	2.5
1	A	183	HIS	2.5
1	A	170	THR	2.5
1	A	216	ALA	2.4
1	A	229	GLY	2.4
1	A	73	GLY	2.3
1	B	89	PHE	2.3
1	A	81	LEU	2.3
1	A	179	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	141	ASP	2.3
1	B	151	VAL	2.2
1	A	184	HIS	2.2
1	B	215	LYS	2.2
1	B	227	ILE	2.2
1	A	181	LYS	2.2
1	A	433	VAL	2.2
2	D	85	SER	2.2
1	B	171	LEU	2.2
3	Q	25	DT	2.1
1	A	137	MET	2.1
1	B	114	GLU	2.1
1	A	75	ASP	2.1
1	A	71	ASN	2.1
2	C	73	PHE	2.1
1	B	115	THR	2.1
1	B	205	ASN	2.1
2	C	97	TYR	2.1
1	A	214	TYR	2.1
1	B	210	HIS	2.0
1	B	133	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	SEP	A	147	10/11	0.78	0.50	-	49,64,77,84	0
1	SEP	B	147	10/11	0.73	0.38	-	41,49,64,80	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.