



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:40 PM GMT

PDB ID : 1PVO
Title : X-ray crystal structure of Rho transcription termination factor in complex with ssRNA substrate and ANPPNP
Authors : Skordalakes, E.; Berger, J.M.
Deposited on : 2003-06-27
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)
RNA backbone	2183	1036 (3.40-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	2	<div> <div>50%</div> <div>50%</div> </div>
1	H	2	<div> <div>50%</div> <div>100%</div> <div>50%</div> </div>
1	J	2	<div> <div>100%</div> <div>100%</div> </div>
1	K	2	<div> <div>100%</div> <div>100%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	L	2	 50% 50%
2	A	419	 8% 55% 34% 9% •
2	B	419	 8% 43% 32% 11% 15%
2	C	419	 2% 53% 35% 9% •
2	D	419	 4% 47% 39% 11% •
2	E	419	 3% 55% 35% 7% •
2	F	419	 6% 51% 34% 11% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ANP	A	601	X	-	-	-
3	ANP	B	602	X	-	-	-
3	ANP	C	603	X	-	-	-
3	ANP	D	604	X	-	-	-
3	ANP	E	605	X	-	-	-
3	ANP	F	606	X	-	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 19246 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(P*UP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	2	Total	C	N	O	P	0	0	0
			40	18	5	15	2			
1	H	2	Total	C	N	O	P	0	0	0
			40	18	5	15	2			
1	J	2	Total	C	N	O	P	0	0	0
			40	18	5	15	2			
1	K	2	Total	C	N	O	P	0	0	0
			40	18	5	15	2			
1	L	2	Total	C	N	O	P	0	0	0
			40	18	5	15	2			

- Molecule 2 is a protein called Transcription termination factor rho.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	408	Total	C	N	O	S	0	0	0
			3208	2025	562	604	17			
2	B	358	Total	C	N	O	S	0	0	0
			2813	1776	494	529	14			
2	C	408	Total	C	N	O	S	0	0	0
			3208	2025	562	604	17			
2	D	408	Total	C	N	O	S	0	0	0
			3208	2025	562	604	17			
2	E	407	Total	C	N	O	S	0	0	0
			3201	2020	561	603	17			
2	F	408	Total	C	N	O	S	0	0	0
			3208	2025	562	604	17			

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
3	B	1	Total 31	C 10	N 6	O 12	P 3	0	0
3	C	1	Total 31	C 10	N 6	O 12	P 3	0	0
3	D	1	Total 31	C 10	N 6	O 12	P 3	0	0
3	E	1	Total 31	C 10	N 6	O 12	P 3	0	0
3	F	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O 1 1	0	0
4	C	5	Total O 5 5	0	0
4	D	4	Total O 4 4	0	0
4	E	2	Total O 2 2	0	0
4	F	2	Total O 2 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-R(P*UP*C)-3'




- Molecule 1: 5'-R(P*UP*C)-3'



- Molecule 1: 5'-R(P*UP*C)-3'



- Molecule 1: 5'-R(P*UP*C)-3'




- Molecule 1: 5'-R(P*UP*C)-3'




- Molecule 2: Transcription termination factor rho







• Molecule 2: Transcription termination factor rho




• Molecule 2: Transcription termination factor rho







• Molecule 2: Transcription termination factor rho




• Molecule 2: Transcription termination factor rho





• Molecule 2: Transcription termination factor rho



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	119.25Å 204.65Å 147.79Å 90.00° 96.54° 90.00°	Depositor
Resolution (Å)	20.00 – 3.00 29.88 – 3.00	Depositor EDS
% Data completeness (in resolution range)	95.7 (20.00-3.00) 96.3 (29.88-3.00)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 3.00Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.271 , 0.304 0.238 , 0.242	Depositor DCC
R_{free} test set	3396 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	89.5	Xtriage
Anisotropy	0.178	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 114.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 67632 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	19246	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	G	0.69	0/43	1.40	1/64 (1.6%)
1	H	0.80	0/43	1.34	1/64 (1.6%)
1	J	0.82	0/43	1.05	0/64
1	K	0.98	0/43	1.54	1/64 (1.6%)
1	L	0.69	0/43	1.39	1/64 (1.6%)
2	A	0.42	0/3253	0.71	11/4379 (0.3%)
2	B	0.43	0/2854	0.71	9/3844 (0.2%)
2	C	0.56	0/3253	0.76	7/4379 (0.2%)
2	D	0.51	0/3253	0.75	10/4379 (0.2%)
2	E	0.49	0/3246	0.74	11/4369 (0.3%)
2	F	0.42	0/3253	0.70	14/4379 (0.3%)
All	All	0.48	0/19327	0.74	66/26049 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	E	0	3
2	F	1	1
All	All	1	5

There are no bond length outliers.

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	77	ASP	CB-CG-OD2	7.04	124.64	118.30
1	G	1	U	P-O3'-C3'	6.65	127.68	119.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	95	ASP	CB-CG-OD2	6.63	124.26	118.30
2	A	265	ASP	CB-CG-OD2	6.28	123.95	118.30
2	F	233	ASP	CB-CG-OD2	6.25	123.92	118.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	F	355	PHE	CA

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	355	PHE	Peptide
2	E	266	SER	Peptide
2	E	355	PHE	Peptide
2	E	371	LEU	Peptide
2	F	24	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	40	0	22	0	0
1	H	40	0	22	2	0
1	J	40	0	22	3	0
1	K	40	0	22	4	0
1	L	40	0	22	3	0
2	A	3208	0	3284	116	0
2	B	2813	0	2870	126	0
2	C	3208	0	3284	152	0
2	D	3208	0	3284	149	0
2	E	3201	0	3275	142	0
2	F	3208	0	3284	124	0
3	A	31	0	7	3	0
3	B	31	0	7	0	0
3	C	31	0	7	0	0
3	D	31	0	7	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	31	0	7	0	0
3	F	31	0	7	0	0
4	A	1	0	0	0	0
4	C	5	0	0	0	0
4	D	4	0	0	0	0
4	E	2	0	0	1	0
4	F	2	0	0	0	0
All	All	19246	0	19433	778	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 778 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:272:ARG:HH11	2:B:272:ARG:HG3	1.01	1.10
2:C:355:PHE:HB2	2:C:356:PRO:HD3	1.26	1.09
2:F:353:ARG:HH11	2:F:353:ARG:HG3	1.06	1.09
2:C:30:ARG:HG2	2:C:30:ARG:HH11	1.14	1.09
2:C:272:ARG:HH11	2:C:272:ARG:HG3	0.92	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	400/419 (96%)	367 (92%)	30 (8%)	3 (1%)	24 66
2	B	350/419 (84%)	313 (89%)	25 (7%)	12 (3%)	5 25
2	C	400/419 (96%)	361 (90%)	32 (8%)	7 (2%)	11 45
2	D	400/419 (96%)	363 (91%)	27 (7%)	10 (2%)	7 34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	399/419 (95%)	370 (93%)	25 (6%)	4 (1%)	19	61
2	F	400/419 (96%)	351 (88%)	35 (9%)	14 (4%)	4	24
All	All	2349/2514 (93%)	2125 (90%)	174 (7%)	50 (2%)	9	40

5 of 50 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	356	PRO
2	C	50	PHE
2	C	355	PHE
2	D	47	GLU
2	D	181	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	350/359 (98%)	271 (77%)	79 (23%)	1	5
2	B	306/359 (85%)	242 (79%)	64 (21%)	1	7
2	C	350/359 (98%)	284 (81%)	66 (19%)	2	10
2	D	350/359 (98%)	273 (78%)	77 (22%)	1	6
2	E	349/359 (97%)	289 (83%)	60 (17%)	2	12
2	F	350/359 (98%)	280 (80%)	70 (20%)	1	8
All	All	2055/2154 (95%)	1639 (80%)	416 (20%)	1	8

5 of 416 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	286	THR
2	D	115	LYS
2	F	266	SER
2	C	326	LYS
2	D	5	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	275	ASN
2	C	409	ASN
2	E	306	ASN
2	C	344	HIS
2	D	32	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	G	2/2 (100%)	1 (50%)	1 (50%)
1	H	2/2 (100%)	1 (50%)	1 (50%)
1	J	1/2 (50%)	0	0
1	K	2/2 (100%)	1 (50%)	1 (50%)
1	L	2/2 (100%)	1 (50%)	1 (50%)
All	All	9/10 (90%)	4 (44%)	4 (44%)

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	G	2	C
1	H	2	C
1	K	2	C
1	L	2	C

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	G	1	U
1	H	1	U
1	K	1	U
1	L	1	U

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ANP	A	601	-	27,33,33	3.52	11 (40%)	30,52,52	3.53	17 (56%)
3	ANP	B	602	-	27,33,33	3.59	10 (37%)	30,52,52	3.79	16 (53%)
3	ANP	C	603	-	27,33,33	3.59	9 (33%)	30,52,52	3.60	15 (50%)
3	ANP	D	604	-	27,33,33	3.59	11 (40%)	30,52,52	3.68	16 (53%)
3	ANP	E	605	-	27,33,33	3.62	11 (40%)	30,52,52	3.64	16 (53%)
3	ANP	F	606	-	27,33,33	3.58	11 (40%)	30,52,52	3.66	15 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	601	-	3/3/7/8	1/12/38/38	0/3/3/3
3	ANP	B	602	-	3/3/7/8	1/12/38/38	0/3/3/3
3	ANP	C	603	-	3/3/7/8	0/12/38/38	0/3/3/3
3	ANP	D	604	-	2/2/7/8	0/12/38/38	0/3/3/3
3	ANP	E	605	-	2/2/7/8	0/12/38/38	0/3/3/3
3	ANP	F	606	-	2/2/7/8	0/12/38/38	0/3/3/3

The worst 5 of 63 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	604	ANP	O2'-C2'	-8.75	1.22	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	ANP	O3'-C3'	-8.73	1.22	1.43
3	E	605	ANP	O2'-C2'	-8.73	1.22	1.43
3	B	602	ANP	O2'-C2'	-8.72	1.22	1.43
3	C	603	ANP	O3'-C3'	-8.67	1.22	1.43

The worst 5 of 95 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	605	ANP	C4'-O4'-C1'	-10.72	97.93	109.72
3	D	604	ANP	C4'-O4'-C1'	-10.59	98.09	109.72
3	F	606	ANP	C4'-O4'-C1'	-9.95	98.78	109.72
3	C	603	ANP	C4'-O4'-C1'	-9.93	98.81	109.72
3	B	602	ANP	C4'-O4'-C1'	-9.47	99.32	109.72

5 of 15 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	B	602	ANP	C4'
3	B	602	ANP	C1'
3	B	602	ANP	C3'
3	F	606	ANP	C4'
3	F	606	ANP	C3'

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	602	ANP	O1B-PB-N3B-PG
3	A	601	ANP	O1B-PB-N3B-PG

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	ANP	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	2/2 (100%)	1.06	0 100 100	150, 150, 150, 150	0
1	H	2/2 (100%)	2.65	2 (100%) 0 0	150, 150, 150, 150	0
1	J	2/2 (100%)	4.42	2 (100%) 0 0	150, 150, 150, 150	0
1	K	2/2 (100%)	4.12	2 (100%) 0 0	150, 150, 150, 150	0
1	L	2/2 (100%)	1.43	0 100 100	150, 150, 150, 150	0
2	A	408/419 (97%)	0.41	33 (8%) 15 5	42, 66, 114, 127	0
2	B	358/419 (85%)	0.53	33 (9%) 11 4	35, 65, 112, 127	0
2	C	408/419 (97%)	0.14	10 (2%) 61 30	20, 43, 96, 103	0
2	D	408/419 (97%)	0.15	16 (3%) 43 18	18, 46, 72, 124	0
2	E	407/419 (97%)	0.10	14 (3%) 49 21	21, 46, 85, 123	0
2	F	408/419 (97%)	0.33	25 (6%) 25 9	39, 65, 105, 119	0
All	All	2407/2524 (95%)	0.28	137 (5%) 27 10	18, 56, 104, 150	0

The worst 5 of 137 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	51	GLY	8.4
2	A	286	THR	6.7
2	A	412	PHE	6.5
2	B	88	ARG	6.4
1	K	2	C	6.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ANP	F	606	31/31	0.80	0.25	0.32	150,150,150,150	0
3	ANP	A	601	31/31	0.72	0.31	-0.20	150,150,150,150	0
3	ANP	E	605	31/31	0.82	0.21	-0.29	150,150,150,150	0
3	ANP	D	604	31/31	0.85	0.20	-0.37	150,150,150,150	0
3	ANP	B	602	31/31	0.84	0.18	-0.92	150,150,150,150	0
3	ANP	C	603	31/31	0.84	0.18	-0.95	150,150,150,150	0

6.5 Other polymers [i](#)

There are no such residues in this entry.