



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:13 AM GMT

PDB ID : 2PV7  
Title : Crystal structure of chorismate mutase / prephenate dehydrogenase (tyrA) (1574749) from Haemophilus influenzae RD at 2.00 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-05-09  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	<div> <div>6%</div> <div> <div></div> <div>84%</div> <div>8%</div> <div>7%</div> </div> </div>
1	B	298	<div> <div>5%</div> <div> <div></div> <div>85%</div> <div>9%</div> <div>6%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called T-protein [Includes: Chorismate mutase (EC 5.4.99.5) (CM) and Prephenate dehydrogenase (EC 1.3.1.12) (PDH)].

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	Se	0	1	0
			2211	1424	373	406	1	7			
1	B	280	Total	C	N	O	S	Se	0	7	1
			2265	1460	382	414	1	8			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	80	GLY	-	LEADER SEQUENCE	UNP P43902
A	81	MSE	MET	MODIFIED RESIDUE	UNP P43902
A	172	MSE	MET	MODIFIED RESIDUE	UNP P43902
A	188	MSE	MET	MODIFIED RESIDUE	UNP P43902
A	202	MSE	MET	MODIFIED RESIDUE	UNP P43902
A	210	MSE	MET	MODIFIED RESIDUE	UNP P43902
A	252	MSE	MET	MODIFIED RESIDUE	UNP P43902
A	294	MSE	MET	MODIFIED RESIDUE	UNP P43902
A	311	MSE	MET	MODIFIED RESIDUE	UNP P43902
B	80	GLY	-	LEADER SEQUENCE	UNP P43902
B	81	MSE	MET	MODIFIED RESIDUE	UNP P43902
B	172	MSE	MET	MODIFIED RESIDUE	UNP P43902
B	188	MSE	MET	MODIFIED RESIDUE	UNP P43902
B	202	MSE	MET	MODIFIED RESIDUE	UNP P43902
B	210	MSE	MET	MODIFIED RESIDUE	UNP P43902
B	252	MSE	MET	MODIFIED RESIDUE	UNP P43902
B	294	MSE	MET	MODIFIED RESIDUE	UNP P43902
B	311	MSE	MET	MODIFIED RESIDUE	UNP P43902

- Molecule 2 is TYROSINE (three-letter code: TYR) (formula: C<sub>9</sub>H<sub>11</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 13	C 9	N 1	O 3	0	0
2	B	1	Total 13	C 9	N 1	O 3	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 44	C 21	N 7	O 14	P 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	156	Total	O	0	0
			156	156		
4	B	237	Total	O	0	0
			237	237		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T-protein [Includes: Chorismate mutase (EC 5.4.99.5) (CM) and Prephenate dehydrogenase (EC 1.3.1.12) (PDH)]



- Molecule 1: T-protein [Includes: Chorismate mutase (EC 5.4.99.5) (CM) and Prephenate dehydrogenase (EC 1.3.1.12) (PDH)]



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.79 Å   127.79 Å   100.62 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	29.70 – 2.00 29.70 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.70-2.00) 99.7 (29.70-2.00)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.161   ,   0.194 0.175   ,   0.202	Depositor DCC
$R_{free}$ test set	2870 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.0	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36   ,   41.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 56542 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4983	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	0/2253	0.76	3/3043 (0.1%)
1	B	0.80	1/2326 (0.0%)	0.76	1/3140 (0.0%)
All	All	0.81	1/4579 (0.0%)	0.76	4/6183 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	358	GLU	CG-CD	5.20	1.59	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	259	ARG	NE-CZ-NH1	6.81	123.70	120.30
1	B	259	ARG	NE-CZ-NH1	5.55	123.07	120.30
1	A	331	LEU	CA-CB-CG	5.47	127.89	115.30
1	A	365	ARG	NE-CZ-NH2	-5.13	117.73	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2211	0	2202	21	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2265	0	2271	20	0
2	A	13	0	8	1	0
2	B	13	0	8	1	0
3	A	44	0	26	1	0
3	B	44	0	26	0	0
4	A	156	0	0	0	0
4	B	237	0	0	2	0
All	All	4983	0	4541	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:ILE:HD12	1:A:280:LEU:HD11	1.60	0.84
1:A:261:PHE:HE2	1:B:327:TYR:CZ	2.14	0.65
1:B:275:ILE:HD12	1:B:280:LEU:HD11	1.79	0.65
2:A:401:TYR:N	1:B:288:TYR:HH	1.99	0.61
1:A:337:ASN:HD21	1:B:311[A]:MSE:HE3	1.72	0.54
1:A:206:ASP:O	1:B:365[A]:ARG:HD3	2.08	0.54
1:A:371:ALA:HA	1:B:370:GLN:OE1	2.07	0.53
1:A:288:TYR:HH	2:B:401:TYR:N	2.06	0.53
1:A:169:THR:O	1:A:172:MSE:HG3	2.11	0.50
1:A:261:PHE:CE2	1:B:327:TYR:CZ	2.98	0.49
1:B:103:ILE:HB	1:B:127:ILE:HD13	1.95	0.49
1:A:261:PHE:HE2	1:B:327:TYR:CE1	2.31	0.48
1:A:350:ARG:C	1:A:350:ARG:HD3	2.34	0.48
1:A:197:LEU:HD13	1:A:226:TYR:CE1	2.51	0.46
1:A:103:ILE:HB	1:A:127:ILE:HD13	1.99	0.45
1:A:303:ALA:HB2	1:B:346:PHE:CD1	2.53	0.44
1:B:223:PRO:HA	1:B:226:TYR:CZ	2.53	0.44
1:A:130:LEU:C	1:A:130:LEU:HD23	2.39	0.43
1:A:223:PRO:HA	1:A:226:TYR:CZ	2.54	0.43
1:A:201:PRO:O	3:A:400:NAD:H5N	2.19	0.43
1:A:323:LEU:HA	1:B:269:HIS:CD2	2.54	0.42
1:B:348:LYS:HE2	4:B:457:HOH:O	2.18	0.42
1:B:182[B]:ARG:HH22	1:B:313:LYS:HD3	1.85	0.42
1:B:203:PHE:CD2	1:B:213:GLN:HB3	2.55	0.42
1:B:182[B]:ARG:NH2	4:B:579:HOH:O	2.53	0.41
1:A:261:PHE:CE2	1:B:327:TYR:CE1	3.09	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:334:PHE:CE1	1:B:311[B]:MSE:HE3	2.55	0.40
1:A:323:LEU:HD23	1:B:327:TYR:OH	2.22	0.40
1:A:311:MSE:HG2	1:B:334:PHE:HD1	1.85	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/298 (92%)	265 (97%)	8 (3%)	0	100	100
1	B	285/298 (96%)	281 (99%)	4 (1%)	0	100	100
All	All	558/596 (94%)	546 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/245 (95%)	229 (98%)	4 (2%)	68	71
1	B	241/245 (98%)	236 (98%)	5 (2%)	61	63
All	All	474/490 (97%)	465 (98%)	9 (2%)	65	67

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	181	LYS
1	A	209	SER
1	A	331	LEU
1	A	350	ARG
1	B	130	LEU
1	B	181	LYS
1	B	209	SER
1	B	270	LEU
1	B	350	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	337	ASN
1	A	366	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAD	A	400	-	38,48,48	1.59	3 (7%)	47,73,73	2.34	9 (19%)
2	TYR	A	401	-	10,13,13	0.96	1 (10%)	11,17,17	0.46	0
3	NAD	B	400	-	38,48,48	1.56	4 (10%)	47,73,73	2.95	8 (17%)
2	TYR	B	401	-	10,13,13	0.81	0	11,17,17	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	400	-	-	0/22/62/62	0/5/5/5
2	TYR	A	401	-	-	0/4/8/8	0/1/1/1
3	NAD	B	400	-	-	0/22/62/62	0/5/5/5
2	TYR	B	401	-	-	0/4/8/8	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	400	NAD	PA-O2A	-2.30	1.45	1.54
2	A	401	TYR	CE1-CD1	2.13	1.42	1.38
3	A	400	NAD	C2A-N1A	2.58	1.38	1.33
3	B	400	NAD	C2A-N1A	2.76	1.39	1.33
3	A	400	NAD	C2A-N3A	2.98	1.37	1.32
3	B	400	NAD	C2A-N3A	3.80	1.38	1.32
3	B	400	NAD	O7N-C7N	6.60	1.38	1.24
3	A	400	NAD	O7N-C7N	7.25	1.39	1.24

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	400	NAD	N3A-C2A-N1A	-14.51	117.78	128.89
3	A	400	NAD	N3A-C2A-N1A	-12.26	119.50	128.89
3	B	400	NAD	O7N-C7N-C3N	-6.76	112.21	119.59
3	B	400	NAD	C1B-N9A-C4A	-5.15	119.17	126.94
3	B	400	NAD	C4B-O4B-C1B	-4.16	105.15	109.72
3	A	400	NAD	O7N-C7N-C3N	-3.30	115.99	119.59
3	A	400	NAD	C1B-N9A-C4A	-2.61	123.01	126.94
3	A	400	NAD	O3-PN-O5D	-2.26	96.93	102.94
3	B	400	NAD	PN-O3-PA	-2.12	126.78	132.73
3	A	400	NAD	PN-O3-PA	-2.02	127.06	132.73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	A	400	NAD	O4D-C1D-N1N	2.24	110.59	108.13
3	A	400	NAD	O2D-C2D-C3D	2.34	119.45	111.83
3	B	400	NAD	O4D-C1D-N1N	2.60	110.98	108.13
3	A	400	NAD	O5D-PN-O1N	2.71	120.13	109.62
3	B	400	NAD	C2A-N1A-C6A	3.15	124.39	118.77
3	A	400	NAD	C3N-C7N-N7N	4.72	122.98	117.82
3	B	400	NAD	C3N-C7N-N7N	7.73	126.28	117.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	400	NAD	1	0
2	A	401	TYR	1	0
2	B	401	TYR	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	269/298 (90%)	0.41	17 (6%) 23 24	26, 34, 44, 77	0
1	B	273/298 (91%)	0.09	16 (5%) 26 27	27, 33, 43, 80	0
All	All	542/596 (90%)	0.25	33 (6%) 25 26	26, 33, 44, 80	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	370	GLN	6.0
1	B	149	ILE	4.5
1	A	149	ILE	4.2
1	B	199	LEU	3.7
1	A	177	LEU	3.7
1	B	177	LEU	3.5
1	A	137	VAL	3.3
1	A	371	ALA	3.2
1	A	138	ALA	3.2
1	A	133	GLU	3.2
1	A	370	GLN	3.2
1	A	199	LEU	3.1
1	A	150	VAL	2.9
1	A	215	VAL	2.9
1	A	175	ALA	2.8
1	A	132	ARG	2.8
1	B	150	VAL	2.6
1	A	136	ALA	2.6
1	B	371	ALA	2.5
1	A	208	ALA	2.5
1	B	105[A]	ILE	2.4
1	B	148	VAL	2.4
1	A	229	LEU	2.4
1	B	112	LEU	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	229	LEU	2.3
1	B	369	GLN	2.2
1	B	258	LEU	2.2
1	B	197	LEU	2.1
1	A	92	GLY	2.1
1	B	175	ALA	2.1
1	A	106	VAL	2.1
1	B	103	ILE	2.0
1	B	215	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	TYR	A	401	13/13	0.92	0.17	1.82	36,40,42,44	0
2	TYR	B	401	13/13	0.95	0.17	0.88	31,35,38,38	0
3	NAD	B	400	44/44	0.94	0.11	-0.26	32,36,41,43	0
3	NAD	A	400	44/44	0.90	0.14	-0.31	36,45,50,59	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.