



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:14 AM GMT

PDB ID : 2PZM
Title : Crystal structure of the Bordetella bronchiseptica enzyme WbmG in complex with NAD and UDP
Authors : Harmer, N.J.; King, J.D.; Palmer, C.M.; Maskell, D.; Blundell, T.L.
Deposited on : 2007-05-18
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	
1	B	330	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	703	-	-	-	X
4	UDP	A	602	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	UDP	B	602	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5357 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative nucleotide sugar epimerase/ dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	6	0
			2384	1528	392	456	8			
1	B	313	Total	C	N	O	S	0	6	0
			2364	1516	391	448	9			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP O87988
A	-18	GLY	-	EXPRESSION TAG	UNP O87988
A	-17	SER	-	EXPRESSION TAG	UNP O87988
A	-16	SER	-	EXPRESSION TAG	UNP O87988
A	-15	HIS	-	EXPRESSION TAG	UNP O87988
A	-14	HIS	-	EXPRESSION TAG	UNP O87988
A	-13	HIS	-	EXPRESSION TAG	UNP O87988
A	-12	HIS	-	EXPRESSION TAG	UNP O87988
A	-11	HIS	-	EXPRESSION TAG	UNP O87988
A	-10	HIS	-	EXPRESSION TAG	UNP O87988
A	-9	SER	-	EXPRESSION TAG	UNP O87988
A	-8	SER	-	EXPRESSION TAG	UNP O87988
A	-7	GLY	-	EXPRESSION TAG	UNP O87988
A	-6	LEU	-	EXPRESSION TAG	UNP O87988
A	-5	VAL	-	EXPRESSION TAG	UNP O87988
A	-4	PRO	-	EXPRESSION TAG	UNP O87988
A	-3	ARG	-	EXPRESSION TAG	UNP O87988
A	-2	GLY	-	EXPRESSION TAG	UNP O87988
A	-1	SER	-	EXPRESSION TAG	UNP O87988
A	0	HIS	-	EXPRESSION TAG	UNP O87988
B	-19	MET	-	EXPRESSION TAG	UNP O87988
B	-18	GLY	-	EXPRESSION TAG	UNP O87988
B	-17	SER	-	EXPRESSION TAG	UNP O87988
B	-16	SER	-	EXPRESSION TAG	UNP O87988
B	-15	HIS	-	EXPRESSION TAG	UNP O87988

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	EXPRESSION TAG	UNP O87988
B	-13	HIS	-	EXPRESSION TAG	UNP O87988
B	-12	HIS	-	EXPRESSION TAG	UNP O87988
B	-11	HIS	-	EXPRESSION TAG	UNP O87988
B	-10	HIS	-	EXPRESSION TAG	UNP O87988
B	-9	SER	-	EXPRESSION TAG	UNP O87988
B	-8	SER	-	EXPRESSION TAG	UNP O87988
B	-7	GLY	-	EXPRESSION TAG	UNP O87988
B	-6	LEU	-	EXPRESSION TAG	UNP O87988
B	-5	VAL	-	EXPRESSION TAG	UNP O87988
B	-4	PRO	-	EXPRESSION TAG	UNP O87988
B	-3	ARG	-	EXPRESSION TAG	UNP O87988
B	-2	GLY	-	EXPRESSION TAG	UNP O87988
B	-1	SER	-	EXPRESSION TAG	UNP O87988
B	0	HIS	-	EXPRESSION TAG	UNP O87988


- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD)

(formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C₉H₁₄N₂O₁₂P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			8	4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	254	Total	O	0	0
			254	254		
5	B	219	Total	O	0	0
			219	219		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative nucleotide sugar epimerase/ dehydratase



- Molecule 1: Putative nucleotide sugar epimerase/ dehydratase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	58.21Å 140.48Å 184.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 28.75 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-2.00) 99.7 (28.75-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.171 , 0.221 0.171 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 51115 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5357	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, SO4, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/2460	0.70	2/3361 (0.1%)
1	B	0.70	0/2439	0.73	2/3332 (0.1%)
All	All	0.71	0/4899	0.72	4/6693 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	59	LEU	CA-CB-CG	7.17	131.78	115.30
1	B	56	ASP	CB-CG-OD1	5.28	123.05	118.30
1	B	21	LEU	CA-CB-CG	5.15	127.14	115.30
1	A	21	LEU	CA-CB-CG	5.06	126.94	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	-2	GLY	Peptide
1	B	-3	ARG	Peptide
1	B	-4	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2384	0	2347	20	0
1	B	2364	0	2335	17	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
3	A	44	0	26	0	0
3	B	44	0	26	0	0
4	A	25	0	11	0	0
4	B	8	0	3	0	0
5	A	254	0	0	0	1
5	B	219	0	0	3	0
All	All	5357	0	4748	37	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:GLU:HG3	1:A:301:ILE:HB	1.64	0.79
1:B:1:MET:H	1:B:25:HIS:HD2	1.29	0.78
1:A:308:PRO:O	1:A:309:LYS:HB2	1.94	0.66
1:B:55:THR:HG23	5:B:765:HOH:O	1.96	0.64
1:A:183:LEU:HD12	1:A:238:VAL:HG23	1.79	0.64
1:A:100:ALA:HB2	1:A:151:LEU:HD23	1.79	0.63
1:A:229:HIS:HD2	1:A:233:GLU:OE1	1.85	0.60
1:A:178:THR:O	1:A:182:ARG:HG2	2.04	0.58
1:B:94[A]:GLN:OE1	5:B:884:HOH:O	2.17	0.58
1:B:129:PRO:HD2	1:B:132:SER:OG	2.04	0.56
1:A:130:ILE:HG21	1:A:269:LYS:HD2	1.88	0.55
1:A:182:ARG:HB2	1:A:189:CYS:SG	2.47	0.55
1:A:271[B]:GLU:HG3	1:A:276:TRP:O	2.07	0.55
1:A:179:PHE:HB3	1:A:238:VAL:HG21	1.90	0.53
1:A:9:ALA:HB2	1:A:41:LEU:HD11	1.93	0.51
1:B:1:MET:H	1:B:25:HIS:CD2	2.18	0.49
1:B:-4:PRO:O	1:B:0:HIS:N	2.44	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:-4:PRO:O	1:B:-1:SER:N	2.45	0.49
1:A:39:GLU:CG	1:A:301:ILE:HB	2.41	0.48
1:B:130:ILE:HD11	1:B:220:VAL:HG23	1.96	0.48
1:A:179:PHE:CB	1:A:238:VAL:HG21	2.44	0.47
1:B:181:LYS:HG3	1:B:298:VAL:HG21	1.97	0.46
1:A:269:LYS:O	1:A:273:GLU:HG3	2.17	0.45
1:B:135:ALA:N	1:B:136:PRO:CD	2.80	0.45
1:B:213:GLN:HB2	1:B:216:ARG:HG3	1.99	0.44
1:A:60:LEU:HD12	1:A:98:ASN:HB3	2.00	0.44
1:B:2:ARG:HB2	1:B:69:PRO:HA	1.99	0.44
1:B:60:LEU:HD12	1:B:98:ASN:HB3	1.99	0.44
1:A:-4:PRO:HB2	1:A:-1[A]:SER:OG	2.18	0.43
1:A:213:GLN:HB2	1:A:216:ARG:HG3	2.00	0.43
1:B:126:VAL:HA	1:B:127:PRO:C	2.40	0.42
1:A:226:GLY:HA2	1:A:265:LEU:O	2.20	0.41
1:B:55:THR:CG2	5:B:765:HOH:O	2.64	0.41
1:B:53:SER:OG	1:B:55:THR:HB	2.21	0.41
1:A:267:PRO:O	1:A:271[B]:GLU:OE1	2.38	0.41
1:B:156:VAL:O	1:B:158:VAL:HG23	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:925:HOH:O	5:A:925:HOH:O 3_554	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	320/330 (97%)	310 (97%)	8 (2%)	2 (1%)	30 22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	317/330 (96%)	311 (98%)	4 (1%)	2 (1%)	30	22
All	All	637/660 (96%)	621 (98%)	12 (2%)	4 (1%)	30	22

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	-2	GLY
1	B	79	TYR
1	A	79	TYR
1	A	308	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/269 (94%)	245 (96%)	9 (4%)	43	40
1	B	252/269 (94%)	243 (96%)	9 (4%)	42	39
All	All	506/538 (94%)	488 (96%)	18 (4%)	46	39

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	LEU
1	A	44	VAL
1	A	59	LEU
1	A	116	THR
1	A	187	GLN
1	A	214[A]	GLU
1	A	214[B]	GLU
1	A	248	GLU
1	A	273	GLU
1	B	21	LEU
1	B	37	LYS
1	B	44	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	55	THR
1	B	245	THR
1	B	246	LEU
1	B	252	VAL
1	B	271[A]	GLU
1	B	271[B]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	187	GLN
1	A	229	HIS
1	B	25	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAD	A	601	-	38,48,48	1.58	4 (10%)	47,73,73	2.66	8 (17%)
4	UDP	A	602	-	18,26,26	1.16	2 (11%)	26,40,40	1.63	4 (15%)
2	SO4	A	701	-	4,4,4	0.32	0	6,6,6	0.42	0
2	SO4	A	702	-	4,4,4	0.56	0	6,6,6	0.50	0
3	NAD	B	601	-	38,48,48	1.76	6 (15%)	47,73,73	2.39	10 (21%)
4	UDP	B	602	-	4,8,26	1.92	1 (25%)	6,10,40	6.47	4 (66%)
2	SO4	B	703	-	4,4,4	0.23	0	6,6,6	1.00	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	601	-	-	0/22/62/62	0/5/5/5
4	UDP	A	602	-	-	0/12/32/32	0/2/2/2
2	SO4	A	701	-	-	0/0/0/0	0/0/0/0
2	SO4	A	702	-	-	0/0/0/0	0/0/0/0
3	NAD	B	601	-	-	0/22/62/62	0/5/5/5
4	UDP	B	602	-	-	0/0/0/32	0/1/1/2
2	SO4	B	703	-	-	0/0/0/0	0/0/0/0

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	NAD	O4D-C4D	-2.40	1.39	1.45
3	B	601	NAD	PN-O2N	-2.03	1.46	1.54
3	B	601	NAD	C4A-N3A	-2.01	1.32	1.35
4	A	602	UDP	C6-N1	2.23	1.38	1.35
3	B	601	NAD	O4B-C1B	2.32	1.44	1.41
4	A	602	UDP	C4-N3	2.77	1.38	1.33
3	B	601	NAD	C2A-N1A	3.13	1.39	1.33
3	A	601	NAD	C2A-N1A	3.13	1.39	1.33
4	B	602	UDP	C4-N3	3.18	1.39	1.33
3	A	601	NAD	C2A-N3A	3.21	1.37	1.32
3	B	601	NAD	C2A-N3A	4.15	1.39	1.32
3	A	601	NAD	O7N-C7N	6.69	1.38	1.24
3	B	601	NAD	O7N-C7N	7.33	1.39	1.24

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	NAD	N3A-C2A-N1A	-14.80	117.56	128.89
3	B	601	NAD	N3A-C2A-N1A	-13.08	118.88	128.89
4	B	602	UDP	N1-C2-N3	-11.27	121.14	128.33
3	A	601	NAD	O7N-C7N-C3N	-4.66	114.50	119.59
3	A	601	NAD	C4A-C5A-N7A	-4.01	105.79	109.48
3	B	601	NAD	O7N-C7N-C3N	-3.65	115.60	119.59
3	B	601	NAD	PN-O3-PA	-3.18	123.80	132.73
4	B	602	UDP	C5-C6-N1	-3.14	120.31	123.90
3	A	601	NAD	C1B-N9A-C4A	-2.43	123.28	126.94
3	B	601	NAD	C4A-C5A-N7A	-2.38	107.29	109.48
3	B	601	NAD	C1B-N9A-C4A	-2.29	123.48	126.94
2	B	703	SO4	O2-S-O1	-2.11	102.82	109.50
3	B	601	NAD	C5N-C4N-C3N	-2.04	117.76	120.33
4	A	602	UDP	O2B-PB-O3A	2.02	114.27	105.09
3	B	601	NAD	O4D-C1D-N1N	2.24	110.60	108.13
4	A	602	UDP	O3A-PA-O5'	2.25	108.92	102.94
3	A	601	NAD	C2A-N1A-C6A	2.30	122.88	118.77
3	B	601	NAD	O2N-PN-O1N	2.55	126.35	112.53
4	A	602	UDP	O4'-C1'-N1	2.72	113.83	108.08
3	A	601	NAD	O4D-C1D-N1N	2.99	111.42	108.13
3	A	601	NAD	C4D-O4D-C1D	3.06	113.08	109.72
3	B	601	NAD	C3N-C7N-N7N	3.16	121.27	117.82
3	B	601	NAD	C4D-O4D-C1D	3.19	113.22	109.72
3	A	601	NAD	C3N-C7N-N7N	5.25	123.56	117.82
4	B	602	UDP	C4-N3-C2	5.63	119.71	114.14
4	A	602	UDP	C4-N3-C2	6.34	120.42	114.14
4	B	602	UDP	C6-N1-C2	8.93	118.79	114.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/330 (95%)	0.73	37 (11%) 6 7	16, 25, 40, 55	0
1	B	313/330 (94%)	0.61	28 (8%) 12 13	15, 27, 43, 47	0
All	All	629/660 (95%)	0.67	65 (10%) 9 9	15, 26, 42, 55	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	308	PRO	6.0
1	B	114	PHE	4.9
1	A	163	LEU	4.6
1	B	-6	LEU	4.6
1	B	163	LEU	4.4
1	B	73	VAL	4.3
1	A	12	LEU	4.1
1	A	166	VAL	4.0
1	B	112	LEU	3.9
1	A	114	PHE	3.9
1	A	112	LEU	3.7
1	A	307	ALA	3.6
1	A	73	VAL	3.5
1	B	16	LEU	3.5
1	A	117	ALA	3.2
1	A	161	LEU	3.2
1	A	264	VAL	3.1
1	B	166	VAL	3.1
1	B	12	LEU	3.1
1	A	-5	VAL	3.1
1	A	309	LYS	3.0
1	A	5	ILE	2.9
1	B	11	CYS	2.9
1	B	161	LEU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	113	ASN	2.9
1	A	-6	LEU	2.9
1	A	245	THR	2.8
1	A	-3	ARG	2.8
1	A	146	ALA	2.8
1	A	275	GLY	2.8
1	A	76	ALA	2.7
1	A	243	GLY	2.7
1	A	263	VAL	2.6
1	A	13	GLY	2.6
1	B	5	ILE	2.6
1	A	119	CYS	2.5
1	A	75	SER	2.5
1	B	75	SER	2.5
1	A	16	LEU	2.5
1	B	145	THR	2.5
1	A	17	ILE	2.4
1	B	223	VAL	2.4
1	A	160	SER	2.4
1	B	74	HIS	2.4
1	A	11	CYS	2.4
1	B	72	VAL	2.3
1	A	272	THR	2.3
1	A	195	VAL	2.3
1	B	-5	VAL	2.3
1	B	6	THR	2.3
1	A	164	ALA	2.3
1	B	43	PRO	2.2
1	B	208	ALA	2.2
1	A	265	LEU	2.2
1	B	204	PHE	2.2
1	B	249	PRO	2.2
1	A	246	LEU	2.1
1	A	238	VAL	2.1
1	A	115	GLN	2.1
1	B	66	SER	2.1
1	B	4	LEU	2.0
1	B	151	LEU	2.0
1	B	275	GLY	2.0
1	B	116	THR	2.0
1	A	7	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	UDP	B	602	8/25	0.89	0.47	14.13	58,59,60,61	0
2	SO4	B	703	5/5	0.97	0.24	6.95	42,43,45,48	0
4	UDP	A	602	25/25	0.79	0.30	3.45	75,77,92,92	0
2	SO4	A	702	5/5	0.98	0.19	0.93	42,42,47,47	0
3	NAD	A	601	44/44	0.96	0.10	-1.28	19,22,27,28	0
3	NAD	B	601	44/44	0.97	0.09	-1.40	17,22,26,28	0
2	SO4	A	701	5/5	0.78	0.34	-	81,81,82,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.