



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:46 AM GMT

PDB ID : 3PZP
Title : Human DNA polymerase kappa extending opposite a cis-syn thymine dimer
Authors : Vasquez-Del Carpio, R.; Silverstein, T.D.; Lone, S.; Johnson, R.E.; Prakash, S.; Prakash, L.; Aggarwal, A.K.
Deposited on : 2010-12-14
Resolution : 3.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1004 (3.40-3.28)
Clashscore	102246	1072 (3.40-3.28)
Ramachandran outliers	100387	1055 (3.40-3.28)
Sidechain outliers	100360	1054 (3.40-3.28)
RSRZ outliers	91569	1009 (3.40-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	517	 8% 51% 30% 15%
1	B	517	 8% 54% 27% 14%
2	P	13	 8% 31% 62% 8%
2	Q	13	 38% 38% 15% 8%
3	T	17	 12% 47% 29% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	U	17	 29% 35% 18% 18%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase kappa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	440	Total	C	N	O	S	0	0	0
			3352	2105	577	648	22			
1	B	443	Total	C	N	O	S	0	0	0
			3192	1994	546	631	21			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	GLY	-	EXPRESSION TAG	UNP Q9UBT6
A	13	PRO	-	EXPRESSION TAG	UNP Q9UBT6
A	14	GLY	-	EXPRESSION TAG	UNP Q9UBT6
A	15	GLY	-	EXPRESSION TAG	UNP Q9UBT6
A	16	ASP	-	EXPRESSION TAG	UNP Q9UBT6
A	17	PRO	-	EXPRESSION TAG	UNP Q9UBT6
A	18	HIS	-	EXPRESSION TAG	UNP Q9UBT6
B	12	GLY	-	EXPRESSION TAG	UNP Q9UBT6
B	13	PRO	-	EXPRESSION TAG	UNP Q9UBT6
B	14	GLY	-	EXPRESSION TAG	UNP Q9UBT6
B	15	GLY	-	EXPRESSION TAG	UNP Q9UBT6
B	16	ASP	-	EXPRESSION TAG	UNP Q9UBT6
B	17	PRO	-	EXPRESSION TAG	UNP Q9UBT6
B	18	HIS	-	EXPRESSION TAG	UNP Q9UBT6


- Molecule 2 is a DNA chain called 5'-D(*GP*GP*GP*GP*GP*AP*AP*GP*GP*AP*CP*C P*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	P	13	Total	C	N	O	P	0	0	0
			272	128	61	71	12			
2	Q	12	Total	C	N	O	P	0	0	0
			250	118	56	65	11			

- Molecule 3 is a DNA chain called 5'-D(*TP*TP*CP*CP*(TTD)P*GP*GP*TP*CP*CP*T P*TP*CP*CP*CP*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	T	15	Total 312	C 151	N 47	O 99	P 15	0	0	0
3	U	14	Total 293	C 142	N 44	O 93	P 14	0	0	0

- Molecule 4 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: C₁₀H₁₆N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	Total 30	C 10	N 5	O 12	P 3	0	0
4	B	1	Total 30	C 10	N 5	O 12	P 3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total 2	Mg 2	0	0
5	A	2	Total 2	Mg 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	5	Total 5	O 5	0	0
6	B	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase kappa



- Molecule 1: DNA polymerase kappa





● Molecule 2: 5'-D(*GP*GP*GP*GP*GP*AP*AP*GP*GP*AP*CP*CP*A)-3'



● Molecule 2: 5'-D(*GP*GP*GP*GP*GP*AP*AP*GP*GP*AP*CP*CP*A)-3'



● Molecule 3: 5'-D(*TP*TP*CP*CP*(TTD)P*GP*GP*TP*CP*CP*TP*TP*CP*CP*CP*CP*C)-3'



● Molecule 3: 5'-D(*TP*TP*CP*CP*(TTD)P*GP*GP*TP*CP*CP*TP*TP*CP*CP*CP*CP*C)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	117.14Å 154.48Å 217.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.67 – 3.34 45.63 – 3.34	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.67-3.34) 99.2 (45.63-3.34)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 3.32Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.242 , 0.288 0.244 , 0.288	Depositor DCC
R_{free} test set	1452 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	98.9	Xtrriage
Anisotropy	0.051	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 95.2	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Outliers	0 of 28882 reflections	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7743	wwPDB-VP
Average B, all atoms (Å ²)	118.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, DTP, TTD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.56	0/3403	0.71	1/4607 (0.0%)
1	B	0.47	0/3240	0.63	0/4406
2	P	1.19	0/308	1.63	11/475 (2.3%)
2	Q	0.99	1/283 (0.4%)	1.55	5/436 (1.1%)
3	T	1.17	1/300 (0.3%)	2.13	17/456 (3.7%)
3	U	1.12	1/279 (0.4%)	1.86	10/424 (2.4%)
All	All	0.65	3/7813 (0.0%)	0.96	44/10804 (0.4%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	4	DC	C3'-O3'	5.37	1.50	1.44
3	U	12	DT	C1'-N1	5.05	1.55	1.49
2	Q	11	DC	C3'-O3'	-5.03	1.37	1.44

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	U	5	DC	O4'-C4'-C3'	-10.84	99.50	106.00
3	T	15	DC	O4'-C4'-C3'	-10.57	99.66	106.00
3	T	16	DC	O4'-C1'-N1	9.58	114.70	108.00
2	P	4	DG	O4'-C1'-N9	9.08	114.36	108.00
3	T	11	DC	O4'-C1'-N1	9.07	114.35	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3352	0	3229	157	0
1	B	3192	0	2896	141	0
2	P	272	0	143	6	0
2	Q	250	0	132	6	0
3	T	312	0	183	6	0
3	U	293	0	172	5	0
4	A	30	0	12	1	0
4	B	30	0	12	1	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	5	0	0	0	0
6	B	3	0	0	1	0
All	All	7743	0	6779	307	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 307 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:457:THR:HG23	1:B:471:ALA:HB2	1.34	1.05
1:B:455:THR:HG22	1:B:473:THR:HB	1.47	0.96
1:A:455:THR:HG23	1:A:515:SER:OG	1.64	0.95
1:B:193:MET:HE2	1:B:408:THR:HG23	1.54	0.90
1:A:83:GLN:HE22	1:A:380:ARG:HH21	1.23	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	436/517 (84%)	367 (84%)	61 (14%)	8 (2%)	11	46
1	B	439/517 (85%)	367 (84%)	61 (14%)	11 (2%)	7	39
All	All	875/1034 (85%)	734 (84%)	122 (14%)	19 (2%)	8	42

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	477	VAL
1	B	351	ILE
1	B	465	PHE
1	A	410	ASP
1	B	306	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/460 (76%)	314 (90%)	35 (10%)	9	35
1	B	306/460 (66%)	281 (92%)	25 (8%)	14	48
All	All	655/920 (71%)	595 (91%)	60 (9%)	11	40

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	477	VAL
1	A	514	SER
1	B	462	ASN
1	A	484	ILE
1	B	23	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	518	ASN
1	B	24	ASN
1	B	332	GLN
1	A	334	GLN
1	B	291	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	TTD	T	6	3	40,45,46	1.45	8 (20%)	58,74,77	2.00	10 (17%)
3	TTD	U	6	3	40,45,46	1.42	6 (15%)	58,74,77	2.32	9 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TTD	T	6	3	-	0/22/109/110	0/3/6/6
3	TTD	U	6	3	-	0/22/109/110	0/3/6/6

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	U	6	TTD	C5-C6	-3.79	1.50	1.55
3	T	6	TTD	C2T-N3T	-3.49	1.31	1.38
3	U	6	TTD	C2T-N3T	-3.30	1.32	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	6	TTD	C4-N3	-3.19	1.32	1.37
3	T	6	TTD	C5-C6	-3.07	1.51	1.55

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	U	6	TTD	O4T-C4T-C5T	-7.75	116.60	122.92
3	U	6	TTD	C2R-C1R-N1T	-7.17	105.82	115.64
3	T	6	TTD	C2'-C1'-N1	-5.92	107.53	115.64
3	T	6	TTD	O4T-C4T-C5T	-5.69	118.28	122.92
3	T	6	TTD	C2R-C1R-N1T	-5.49	108.12	115.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	T	6	TTD	2	0
3	U	6	TTD	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DTP	A	1	5	24,32,32	0.98	1 (4%)	32,50,50	1.66	4 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DTP	B	2	5	24,32,32	0.94	1 (4%)	32,50,50	1.87	5 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTP	A	1	5	-	0/18/34/34	0/3/3/3
4	DTP	B	2	5	-	0/18/34/34	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1	DTP	C5-C4	2.39	1.45	1.40
4	B	2	DTP	C5-C4	3.15	1.47	1.40

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2	DTP	N3-C2-N1	-7.94	122.81	128.89
4	A	1	DTP	N3-C2-N1	-6.31	124.06	128.89
4	A	1	DTP	C2'-C1'-N9	-3.41	105.86	114.16
4	A	1	DTP	PB-O3B-PG	-3.08	122.35	132.67
4	A	1	DTP	C4-C5-N7	-2.69	107.01	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1	DTP	1	0
4	B	2	DTP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	440/517 (85%)	-0.06	7 (1%) 74 74	51, 107, 160, 223	0
1	B	443/517 (85%)	-0.17	5 (1%) 82 83	81, 129, 180, 234	0
2	P	13/13 (100%)	0.00	1 (7%) 16 16	77, 111, 135, 136	0
2	Q	12/13 (92%)	0.03	0 100 100	101, 121, 146, 153	0
3	T	14/17 (82%)	-0.04	0 100 100	86, 111, 133, 135	0
3	U	13/17 (76%)	-0.01	0 100 100	104, 122, 147, 147	0
All	All	935/1094 (85%)	-0.11	13 (1%) 78 79	51, 120, 173, 234	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	410	ASP	3.2
1	B	130	VAL	2.7
1	B	409	ARG	2.7
1	A	220	ARG	2.5
1	A	476	SER	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	TTD	U	6	40/41	0.93	0.23	-	98,107,111,112	0
3	TTD	T	6	40/41	0.96	0.22	-	79,85,89,91	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	MG	A	3	1/1	0.94	0.30	1.45	76,76,76,76	0
5	MG	B	4	1/1	0.68	0.29	1.00	81,81,81,81	0
4	DTP	A	1	30/30	0.98	0.21	0.44	72,75,77,77	0
4	DTP	B	2	30/30	0.94	0.18	-0.49	112,116,121,121	0
5	MG	A	2	1/1	0.99	0.17	-0.92	70,70,70,70	0
5	MG	B	1	1/1	0.92	0.13	-	86,86,86,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.