



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:10 PM GMT

PDB ID : 1Q45  
Title : 12-0xo-phytodienoate reductase isoform 3  
Authors : Phillips Jr., G.N.; Johnson, K.A.; Bingman, C.A.; Smith, D.W.; Center for Eukaryotic Structural Genomics (CESG)  
Deposited on : 2003-08-01  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	391	<div> <div>4%</div> <div> <div></div> <div>73%</div> <div>19%</div> <div>• 7%</div> </div> </div>
1	B	391	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>19%</div> <div>• 6%</div> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 12-oxophytodienoate-10,11-reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	365	Total	C	N	O	S	0	1	0
			2804	1775	496	521	12			
1	B	367	Total	C	N	O	S	0	0	0
			2830	1795	495	528	12			

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	203	Total 203	O 203	0	0
3	B	207	Total 207	O 207	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: 12-oxophytodienoate-10,11-reductase



#### • Molecule 1: 12-oxophytodienoate-10,11-reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.11Å 85.07Å 121.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.92 – 2.00 24.90 – 2.00	Depositor EDS
% Data completeness (in resolution range)	91.3 (24.92-2.00) 91.3 (24.90-2.00)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	0.13	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.15 (at 1.99Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.186 , 0.236 0.196 , 0.238	Depositor DCC
$R_{free}$ test set	2572 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.2	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 37.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 50628 reflections (0.004%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6106	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.91	1/2873 (0.0%)	0.92	7/3892 (0.2%)
1	B	0.91	2/2897 (0.1%)	0.92	14/3926 (0.4%)
All	All	0.91	3/5770 (0.1%)	0.92	21/7818 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	311	MET	CG-SD	6.56	1.98	1.81
1	B	354	SER	CB-OG	-5.22	1.35	1.42
1	B	226	VAL	CB-CG1	-5.19	1.42	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	351	ASP	CB-CG-OD2	9.84	127.16	118.30
1	B	203	ASP	CB-CG-OD2	9.51	126.86	118.30
1	A	381	ASP	CB-CG-OD2	8.62	126.06	118.30
1	A	204	ARG	NE-CZ-NH2	-7.20	116.70	120.30
1	B	359	ASP	CB-CG-OD2	6.65	124.28	118.30
1	B	170	LEU	CA-CB-CG	6.63	130.55	115.30
1	A	206	ASP	CB-CG-OD2	6.56	124.20	118.30
1	A	247	ASP	CB-CG-OD2	6.30	123.97	118.30
1	A	252	ASP	CB-CG-OD2	6.23	123.90	118.30
1	A	203	ASP	CB-CG-OD2	6.15	123.84	118.30
1	B	144	ASP	CB-CG-OD2	6.05	123.74	118.30
1	B	381	ASP	CB-CG-OD2	5.80	123.52	118.30
1	B	247	ASP	CB-CG-OD1	-5.70	113.17	118.30
1	B	351	ASP	CB-CG-OD2	5.64	123.38	118.30
1	B	22	ASP	CB-CG-OD2	5.62	123.36	118.30
1	B	247	ASP	CB-CG-OD2	5.54	123.29	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	344	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	B	252	ASP	CB-CG-OD2	5.43	123.19	118.30
1	B	181	ASP	CB-CG-OD2	5.14	122.93	118.30
1	B	167	ASP	CB-CG-OD2	5.08	122.87	118.30
1	B	344	ARG	NE-CZ-NH2	-5.01	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2804	0	2758	51	0
1	B	2830	0	2768	52	0
2	A	31	0	19	1	0
2	B	31	0	19	3	0
3	A	203	0	0	1	0
3	B	207	0	0	4	0
All	All	6106	0	5564	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:344:ARG:HD3	2:B:9401:FMN:O1P	1.82	0.80
1:A:50:TYR:O	1:A:54:THR:HG22	1.87	0.74
1:A:156:LEU:O	1:A:204:ARG:NH2	2.13	0.73
1:A:97:HIS:HE1	1:A:181:ASP:OD2	1.71	0.73
1:B:97:HIS:HE1	1:B:181:ASP:OD2	1.74	0.70
1:B:135:GLU:HB3	1:B:149:LYS:HD3	1.75	0.69
1:B:188:ALA:HB1	1:B:189:HIS:CD2	2.31	0.66
1:A:80:ILE:HD12	1:A:89:TRP:CD1	2.31	0.66
1:A:324:ASN:C	1:A:324:ASN:HD22	1.99	0.65

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:39:ASN:H	1:B:76:HIS:HD2	1.42	0.65
1:B:271:ASN:ND2	3:B:574:HOH:O	2.29	0.65
1:B:50:TYR:O	1:B:54:THR:HG22	1.98	0.64
1:A:324:ASN:ND2	1:A:327:LEU:H	1.94	0.64
1:B:112:ARG:H	1:B:195:GLN:HE22	1.47	0.62
1:B:183:ILE:HD11	1:B:226:VAL:HG11	1.83	0.60
1:A:324:ASN:HD21	1:A:327:LEU:H	1.48	0.60
1:B:48:GLU:O	1:B:52:GLN:HG3	2.03	0.59
1:A:183:ILE:HD11	1:A:226:VAL:HG11	1.84	0.59
1:A:52:GLN:HE21	1:A:53:ARG:HH22	1.50	0.59
1:B:54:THR:HG21	1:B:96:VAL:HA	1.84	0.58
1:A:97:HIS:CE1	1:A:181:ASP:OD2	2.56	0.56
1:B:40:GLY:HA2	1:B:77:VAL:O	2.06	0.56
1:A:54:THR:HG21	1:A:96:VAL:HA	1.89	0.55
1:A:268:GLN:HE21	1:A:275:LEU:H	1.54	0.55
1:A:108:TRP:CD1	1:A:186:HIS:CD2	2.95	0.54
1:A:188:ALA:HB1	1:A:189:HIS:CD2	2.43	0.54
1:B:268:GLN:HE21	1:B:275:LEU:H	1.55	0.53
1:A:345:LEU:O	1:A:349:ASN:HB2	2.09	0.53
1:B:140:VAL:HB	1:B:150:TYR:CE1	2.43	0.53
1:B:112:ARG:HB3	1:B:127:SER:HB2	1.90	0.52
1:B:214:ASN:ND2	1:B:217:ARG:HH11	2.07	0.51
1:B:137:ARG:HB2	1:B:138:TRP:CD1	2.46	0.51
1:B:252:ASP:OD1	1:B:255:SER:OG	2.29	0.51
1:A:358:ILE:HD11	1:A:385:LEU:CD1	2.41	0.51
1:A:232:ALA:O	1:A:275:LEU:HA	2.12	0.50
1:A:112:ARG:H	1:A:195:GLN:HE22	1.59	0.50
1:A:183:ILE:C	1:A:183:ILE:HD12	2.32	0.50
1:A:270:VAL:O	1:A:270:VAL:HG12	2.12	0.50
1:B:268:GLN:NE2	1:B:275:LEU:H	2.10	0.50
1:A:198:LYS:HB3	1:A:249:THR:O	2.12	0.49
1:A:238:ARG:HE	1:A:279:HIS:CD2	2.30	0.49
1:A:140:VAL:HB	1:A:150:TYR:CE1	2.47	0.49
1:B:54:THR:HG22	1:B:60:LEU:HD11	1.95	0.49
1:B:76:HIS:HE1	3:B:663:HOH:O	1.96	0.49
1:A:279:HIS:CE1	1:A:318:MET:SD	3.06	0.49
1:B:198:LYS:HG3	1:B:248:ALA:HB1	1.94	0.49
1:A:97:HIS:HD2	3:A:591:HOH:O	1.95	0.49
1:A:40:GLY:HA2	1:A:77:VAL:O	2.13	0.49
1:B:21:PHE:CE2	1:B:181:ASP:HB3	2.48	0.48
1:A:204:ARG:HB2	1:A:209:GLY:HA3	1.94	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:39:ASN:H	1:B:76:HIS:CD2	2.28	0.48
1:A:324:ASN:C	1:A:324:ASN:ND2	2.66	0.48
1:A:187:GLY:HA2	1:A:193:ILE:HB	1.95	0.48
1:A:135:GLU:O	1:A:139:ARG:NH1	2.46	0.47
1:A:270:VAL:CG1	1:A:270:VAL:O	2.61	0.47
1:B:66:MET:SD	1:B:72:ALA:HB2	2.55	0.47
1:B:97:HIS:HD2	3:B:902:HOH:O	1.99	0.46
1:A:254:LEU:HA	1:A:305:LEU:HD13	1.97	0.46
1:A:377:VAL:HA	1:A:381:ASP:OD1	2.16	0.46
1:B:189:HIS:HE1	2:B:9401:FMN:H1'1	1.82	0.45
1:A:135:GLU:HB2	1:A:149:LYS:HE2	1.97	0.45
1:B:69:PRO:HB2	1:B:120:PRO:HG3	1.98	0.45
1:B:93:VAL:HG12	1:B:97:HIS:CD2	2.52	0.45
1:A:112:ARG:HB3	1:A:127:SER:HB2	1.97	0.45
1:B:322:GLY:HA2	2:B:9401:FMN:O3P	2.17	0.45
1:B:16:TYR:HB2	1:B:26:ARG:NE	2.32	0.45
1:B:285:TYR:HE2	1:B:299:ASP:HB3	1.82	0.45
1:B:201:ILE:HD11	1:B:248:ALA:HA	1.98	0.45
1:A:193:ILE:HG21	1:A:219:LEU:HD13	1.98	0.45
1:B:177:ARG:HD3	3:B:935:HOH:O	2.16	0.45
1:A:219:LEU:O	1:A:223:VAL:HG23	2.16	0.44
1:B:32:MET:O	1:B:34:ARG:HG2	2.18	0.44
1:B:318:MET:HA	1:B:339:LEU:O	2.17	0.44
1:B:137:ARG:HB2	1:B:138:TRP:NE1	2.33	0.44
1:B:97:HIS:CE1	1:B:181:ASP:OD2	2.64	0.43
1:A:268:GLN:NE2	1:A:275:LEU:H	2.15	0.43
1:B:345:LEU:O	1:B:349:ASN:HB2	2.18	0.43
1:B:252:ASP:CG	1:B:255:SER:OG	2.57	0.43
1:A:303:ALA:O	1:A:307:LYS:HG3	2.18	0.43
1:B:187:GLY:HA2	1:B:193:ILE:HB	2.01	0.43
1:B:80:ILE:HD13	1:B:89:TRP:CD1	2.54	0.43
1:B:33:THR:O	1:B:34:ARG:HD2	2.19	0.43
1:A:18:MET:HE1	1:A:21:PHE:CD1	2.54	0.42
1:B:189:HIS:N	1:B:189:HIS:CD2	2.86	0.42
1:B:50:TYR:O	1:B:54:THR:CG2	2.66	0.42
1:A:324:ASN:H	1:A:324:ASN:ND2	2.16	0.42
1:A:254:LEU:HA	1:A:305:LEU:CD1	2.49	0.42
1:B:126:ILE:HD12	1:B:156:LEU:HD21	2.01	0.42
1:A:371:TYR:CZ	2:A:7401:FMN:HM72	2.55	0.42
1:A:268:GLN:NE2	1:A:313:TYR:OH	2.47	0.42
1:B:60:LEU:O	1:B:103:ILE:HA	2.20	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:285:TYR:CE2	1:B:299:ASP:HB3	2.55	0.41
1:A:18:MET:CE	1:A:21:PHE:HD1	2.32	0.41
1:A:358:ILE:HD11	1:A:385:LEU:HD13	2.02	0.41
1:B:161:ILE:N	1:B:162:PRO:HD2	2.34	0.41
1:B:345:LEU:HD23	1:B:345:LEU:HA	1.95	0.41
1:B:238:ARG:HA	1:B:279:HIS:O	2.21	0.41
1:A:60:LEU:O	1:A:103:ILE:HA	2.20	0.41
1:A:348:ALA:HB2	1:A:365:TYR:HA	2.04	0.40
1:A:324:ASN:HD21	1:A:327:LEU:HG	1.86	0.40
1:A:261:VAL:HG11	1:A:312:ALA:CB	2.52	0.40
1:A:137:ARG:CZ	1:A:137:ARG:HB2	2.51	0.40
1:B:64:GLY:HA3	1:B:108:TRP:CE3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	362/391 (93%)	351 (97%)	11 (3%)	0	100	100
1	B	363/391 (93%)	350 (96%)	13 (4%)	0	100	100
All	All	725/782 (93%)	701 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/317 (94%)	282 (95%)	15 (5%)	29	23
1	B	298/317 (94%)	283 (95%)	15 (5%)	30	24
All	All	595/634 (94%)	565 (95%)	30 (5%)	29	24

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	LYS
1	A	20	ARG
1	A	54	THR
1	A	80	ILE
1	A	137	ARG
1	A	186	HIS
1	A	212	ILE
1	A	241	PRO
1	A	274	LYS
1	A	284	ARG
1	A	310	ARG
1	A	324	ASN
1	A	346	PHE
1	A	354	SER
1	A	367	ARG
1	B	48	GLU
1	B	54	THR
1	B	135	GLU
1	B	137	ARG
1	B	170	LEU
1	B	212	ILE
1	B	249	THR
1	B	255	SER
1	B	304	LYS
1	B	307	LYS
1	B	310	ARG
1	B	344	ARG
1	B	346	PHE
1	B	354	SER
1	B	358	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	52	GLN
1	A	97	HIS
1	A	195	GLN
1	A	221	GLN
1	A	268	GLN
1	A	314	ASN
1	A	324	ASN
1	A	334	GLN
1	B	76	HIS
1	B	97	HIS
1	B	174	ASN
1	B	195	GLN
1	B	214	ASN
1	B	268	GLN
1	B	314	ASN
1	B	373	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FMN	A	7401	-	32,33,33	1.20	4 (12%)	34,50,50	2.20	9 (26%)
2	FMN	B	9401	-	32,33,33	1.31	4 (12%)	34,50,50	2.43	8 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	7401	-	-	0/18/18/18	0/3/3/3
2	FMN	B	9401	-	-	0/18/18/18	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	9401	FMN	C10-N10	-3.79	1.34	1.39
2	A	7401	FMN	C6-C5A	-2.91	1.37	1.41
2	A	7401	FMN	C4A-C10	-2.14	1.36	1.40
2	B	9401	FMN	C6-C5A	-2.13	1.38	1.41
2	B	9401	FMN	O5'-C5'	-2.11	1.36	1.44
2	A	7401	FMN	P-O1P	-2.06	1.44	1.50
2	A	7401	FMN	C4-N3	2.76	1.38	1.33
2	B	9401	FMN	C4A-N5	2.99	1.37	1.33

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	7401	FMN	N3-C2-N1	-5.44	118.53	127.69
2	B	9401	FMN	N3-C2-N1	-4.35	120.36	127.69
2	B	9401	FMN	C4A-C4-N3	-4.04	118.24	123.52
2	A	7401	FMN	C4A-C4-N3	-3.68	118.72	123.52
2	A	7401	FMN	O3P-P-O5'	-2.94	98.15	106.72
2	B	9401	FMN	C6-C5A-N5	-2.22	116.14	118.92
2	A	7401	FMN	O2P-P-O5'	2.10	112.85	106.72
2	B	9401	FMN	O2'-C2'-C1'	2.42	115.91	109.93
2	A	7401	FMN	C6-C5A-C9A	2.45	121.81	119.11
2	A	7401	FMN	O5'-P-O1P	2.62	113.67	107.08
2	A	7401	FMN	C4A-N5-C5A	2.90	120.14	116.72
2	A	7401	FMN	C1'-N10-C9A	3.02	122.33	118.83
2	B	9401	FMN	C5A-C9A-N10	3.41	120.13	117.58
2	B	9401	FMN	C6-C5A-C9A	4.11	123.64	119.11
2	B	9401	FMN	C1'-N10-C9A	4.56	124.12	118.83

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	7401	FMN	C4-N3-C2	7.75	121.63	115.16
2	B	9401	FMN	C4-N3-C2	8.87	122.56	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	7401	FMN	1	0
2	B	9401	FMN	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	365/391 (93%)	0.13	16 (4%) 38 39	20, 30, 48, 63	0
1	B	367/391 (93%)	-0.07	11 (2%) 54 55	19, 28, 41, 61	0
All	All	732/782 (93%)	0.03	27 (3%) 45 47	19, 29, 46, 63	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	270	VAL	8.5
1	A	272	GLY	6.5
1	A	271	ASN	5.2
1	B	270	VAL	4.5
1	A	269	GLY	4.2
1	B	286	HIS	4.0
1	B	269	GLY	3.1
1	A	284	ARG	3.1
1	B	287	ALA	3.0
1	A	273	SER	3.0
1	A	243	ILE	2.8
1	A	303	ALA	2.8
1	A	8	SER	2.7
1	A	137	ARG	2.7
1	A	244	ASP	2.6
1	A	20	ARG	2.5
1	A	300	GLU	2.5
1	B	300	GLU	2.5
1	A	9	ASN	2.5
1	B	9	ASN	2.3
1	B	137	ARG	2.3
1	B	271	ASN	2.2
1	A	192	LEU	2.2
1	A	301	GLU	2.2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	333	GLN	2.2
1	B	285	TYR	2.1
1	B	272	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FMN	B	9401	31/31	0.97	0.09	-1.17	15,23,28,30	0
2	FMN	A	7401	31/31	0.97	0.10	-1.24	16,22,26,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.