



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 04:22 PM BST

PDB ID : 1Q69  
Title : Solution structure of T-cell surface glycoprotein CD8 alpha chain and Proto-oncogene tyrosine-protein kinase LCK fragments  
Authors : Kim, P.W.; Sun, Z.Y.; Blacklow, S.C.; Wagner, G.; Eck, M.J.  
Deposited on : 2003-08-12

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 77%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	19	
2	B	29	

## 2 Ensemble composition and analysis

This entry contains 10 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *noe limit 0.5 angstrom, angle violation 5 degrees*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:192-A:200, B:19-B:29 (20)	0.21	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	5, 7, 10
2	1, 3, 9
3	2, 6, 8
Single-model clusters	4

### 3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 766 atoms, of which 378 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called T-cell surface glycoprotein CD8 alpha chain.

Mol	Chain	Residues	Atoms						Trace
1	A	19	Total	C	H	N	O	S	0
			329	92	173	38	24	2	

- Molecule 2 is a protein called Proto-oncogene tyrosine-protein kinase LCK.

Mol	Chain	Residues	Atoms						Trace
2	B	29	Total	C	H	N	O	S	0
			436	143	205	37	49	2	

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	14	LEU	MET	ENGINEERED	UNP P06239

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
3	A	1	Total	Zn
			1	1

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain



- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain




- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2.2 Score per residue for model 2

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain




- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2.3 Score per residue for model 3

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain




- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2.4 Score per residue for model 4

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain




- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2.5 Score per residue for model 5

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain




- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2.6 Score per residue for model 6

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain




- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2.7 Score per residue for model 7

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain




- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2.8 Score per residue for model 8

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain




- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2.9 Score per residue for model 9

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain




- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK




### 4.2.10 Score per residue for model 10 (medoid)

- Molecule 1: T-cell surface glycoprotein CD8 alpha chain



- Molecule 2: Proto-oncogene tyrosine-protein kinase LCK



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry, simulated annealing, molecular dynamics, torsion angle dynamics*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *The submitted conformer models are the 10 structures with the lowest energy..*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	.
X-PLOR	structure solution	.
X-PLOR	refinement	.

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5945
Number of chemical shift lists	2
Total number of shifts	457
Number of shifts mapped to atoms	457
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	77%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality ⓘ

### 6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	71	79	79	4±2
2	B	87	80	79	8±2
All	All	1590	1590	1580	87

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:28:VAL:HG12	2:B:29:PRO:HD2	0.96	1.37	5	10
2:B:28:VAL:HG12	2:B:29:PRO:CD	0.77	2.10	4	10
1:A:194:CYS:N	2:B:25:TYR:HB3	0.58	2.13	6	6
1:A:198:ARG:O	1:A:198:ARG:CG	0.58	2.52	2	1
2:B:27:ILE:HG22	2:B:27:ILE:O	0.55	2.01	3	4
2:B:27:ILE:O	2:B:27:ILE:HG22	0.53	2.02	10	6
2:B:28:VAL:CG1	2:B:29:PRO:CD	0.51	2.87	7	10
1:A:193:VAL:C	2:B:25:TYR:HB3	0.50	2.27	4	6
1:A:194:CYS:N	2:B:25:TYR:CB	0.50	2.75	7	7
1:A:194:CYS:SG	2:B:27:ILE:HD11	0.48	2.48	4	5
2:B:25:TYR:N	2:B:25:TYR:CD1	0.46	2.83	3	4

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:193:VAL:C	2:B:25:TYR:CB	0.45	2.85	3	1
1:A:197:PRO:O	1:A:198:ARG:CB	0.43	2.66	2	1
1:A:194:CYS:SG	2:B:27:ILE:CG1	0.43	3.07	5	3
1:A:193:VAL:O	2:B:25:TYR:CD2	0.41	2.73	2	3
2:B:19:VAL:HG13	2:B:26:PRO:HA	0.41	1.92	4	1
1:A:194:CYS:HB2	2:B:27:ILE:HG13	0.41	1.92	1	2
1:A:194:CYS:N	2:B:25:TYR:HB2	0.41	2.31	1	1
2:B:23:CYS:O	2:B:24:HIS:CB	0.41	2.68	7	1
1:A:194:CYS:CB	2:B:27:ILE:HG13	0.41	2.46	4	2
1:A:194:CYS:CB	2:B:27:ILE:CG1	0.41	2.99	1	1
2:B:20:CYS:HB2	2:B:27:ILE:HG13	0.40	1.93	1	1
1:A:198:ARG:O	1:A:198:ARG:HG2	0.40	2.16	2	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	9/19 (47%)	4±1 (44±7%)	4±1 (44±9%)	1±0 (11±5%)	1	9
2	B	11/29 (38%)	8±1 (69±12%)	3±1 (25±12%)	1±0 (6±4%)	3	20
All	All	200/480 (42%)	116 (58%)	67 (34%)	17 (8%)	2	13

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	198	ARG	7
2	B	20	CYS	7
1	A	192	ARG	3

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	9/18 (50%)	8±1 (84±7%)	1±1 (16±7%)	7	45
2	B	11/27 (41%)	9±1 (77±7%)	3±1 (23±7%)	4	30
All	All	200/450 (44%)	161 (80%)	39 (20%)	5	37

All 8 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	28	VAL	10
2	B	24	HIS	8
1	A	198	ARG	7
1	A	192	ARG	4
1	A	195	LYS	3
2	B	22	ASN	3
2	B	21	GLU	3
2	B	25	TYR	1

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

## 6.7 Other polymers ⓘ

There are no such molecules in this entry.

## 6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 77% for the well-defined parts and 73% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 5945

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	190
Number of shifts mapped to atoms	190
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	19	$0.23 \pm 0.31$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	18	$0.30 \pm 0.52$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	16	$-3.22 \pm 0.40$	Should be applied

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 36%, i.e. 94 atoms were assigned a chemical shift out of a possible 258. 2 out of 4 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	32/92 (35%)	16/36 (44%)	9/40 (22%)	7/16 (44%)
Sidechain	62/150 (41%)	38/91 (42%)	24/51 (47%)	0/8 (0%)

*Continued on next page...*

Continued from previous page...

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	0/16 (0%)	0/8 (0%)	0/6 (0%)	0/2 (0%)
Overall	94/258 (36%)	54/135 (40%)	33/97 (34%)	7/26 (27%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 32%, i.e. 195 atoms were assigned a chemical shift out of a possible 615. 3 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	70/230 (30%)	35/91 (38%)	19/96 (20%)	16/43 (37%)
Sidechain	125/349 (36%)	78/208 (38%)	47/119 (39%)	0/22 (0%)
Aromatic	0/36 (0%)	0/18 (0%)	0/13 (0%)	0/5 (0%)
Overall	195/615 (32%)	113/317 (36%)	66/228 (29%)	16/70 (23%)


#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



## 7.2 Chemical shift list 2

File name: BMRB entry 5945

Chemical shift list name: *assigned\_chem\_shift\_list\_2*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	267
Number of shifts mapped to atoms	267
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

### 7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	28	$-0.23 \pm 0.15$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	26	$0.81 \pm 0.49$	None needed (imprecise)
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	24	$-0.74 \pm 0.38$	None needed (imprecise)

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 40%, i.e. 104 atoms were assigned a chemical shift out of a possible 258. 1 out of 4 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	40/92 (43%)	20/36 (56%)	11/40 (28%)	9/16 (56%)
Sidechain	59/150 (39%)	37/91 (41%)	22/51 (43%)	0/8 (0%)
Aromatic	5/16 (31%)	5/8 (62%)	0/6 (0%)	0/2 (0%)
Overall	104/258 (40%)	62/135 (46%)	33/97 (34%)	9/26 (35%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 42%, i.e. 257 atoms were assigned a chemical shift out of a possible 615. 3 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	104/230 (45%)	52/91 (57%)	28/96 (29%)	24/43 (56%)

*Continued on next page...*

*Continued from previous page...*

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Sidechain	141/349 (40%)	88/208 (42%)	53/119 (45%)	0/22 (0%)
Aromatic	12/36 (33%)	11/18 (61%)	0/13 (0%)	1/5 (20%)
Overall	257/615 (42%)	151/317 (48%)	81/228 (36%)	25/70 (36%)

#### 7.2.4 Statistically unusual chemical shifts ⓘ

There are no statistically unusual chemical shifts.

#### 7.2.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:

