



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2016 – 09:12 PM EDT

PDB ID : 4Q9K
Title : P-glycoprotein cocrystallised with QZ-Leu
Authors : McGrath, A.P.; Szewczyk, P.; Chang, G.
Deposited on : 2014-05-01
Resolution : 3.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027457
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1317 (4.10-3.50)
Clashscore	102246	1458 (4.10-3.50)
Ramachandran outliers	100387	1397 (4.10-3.50)
Sidechain outliers	100360	1392 (4.10-3.50)
RSRZ outliers	91569	1325 (4.10-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1284	<div> <div>5%</div> <div>87%</div> <div>6%</div> <div>8%</div> </div>
2	B	6	<div> <div>17%</div> <div>100%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9246 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multidrug resistance protein 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1187	9207	5919	1562	1688	38	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	83	GLN	ASN	CONFLICT	UNP P21447
A	87	GLN	ASN	CONFLICT	UNP P21447
A	90	GLN	ASN	CONFLICT	UNP P21447
A	1277	LEU	-	EXPRESSION TAG	UNP P21447
A	1278	GLU	-	EXPRESSION TAG	UNP P21447
A	1279	HIS	-	EXPRESSION TAG	UNP P21447
A	1280	HIS	-	EXPRESSION TAG	UNP P21447
A	1281	HIS	-	EXPRESSION TAG	UNP P21447
A	1282	HIS	-	EXPRESSION TAG	UNP P21447
A	1283	HIS	-	EXPRESSION TAG	UNP P21447
A	1284	HIS	-	EXPRESSION TAG	UNP P21447

- Molecule 2 is a protein called (30F)L(30F)L(30F)L Peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
2	B	6	39	27	6	3	3	0	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.24Å 138.60Å 195.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.55 – 3.80 49.55 – 3.80	Depositor EDS
% Data completeness (in resolution range)	91.6 (49.55-3.80) 91.2 (49.55-3.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 3.77Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.292 , 0.321 0.299 , 0.332	Depositor DCC
R_{free} test set	1161 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	168.4	Xtriage
Anisotropy	0.321	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.23 , 94.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	9246	wwPDB-VP
Average B, all atoms (Å ²)	184.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 30F

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/9376	0.42	0/12671
2	B	0.64	0/18	0.82	0/21
All	All	0.26	0/9394	0.42	0/12692

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9207	0	9398	37	0
2	B	39	0	33	0	0
All	All	9246	0	9431	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:ASP:OD2	1:A:901:ARG:NH2	2.28	0.66
1:A:266:GLN:NE2	1:A:793:LEU:O	2.32	0.62
1:A:1050:GLN:NE2	1:A:1244:ASN:O	2.33	0.61
1:A:1207:GLU:OE1	1:A:1229:ARG:NH2	2.36	0.59
1:A:604:GLU:OE1	1:A:617:ILE:N	2.38	0.57
1:A:922:ILE:HB	1:A:923:PRO:HD3	1.88	0.56
1:A:1173:SER:OG	1:A:1176:GLN:OE1	2.24	0.55
1:A:734:VAL:O	1:A:738:GLY:N	2.40	0.55
1:A:387:ASN:O	1:A:450:ASP:N	2.40	0.54
1:A:1249:GLU:OE1	1:A:1262:ILE:N	2.41	0.53
1:A:1114:GLN:O	1:A:1178:GLN:NE2	2.43	0.52
1:A:477:ALA:O	1:A:478:THR:HG23	2.10	0.51
1:A:1204:THR:OG1	1:A:1205:GLU:N	2.45	0.50
1:A:93:GLU:N	1:A:93:GLU:OE1	2.44	0.50
1:A:801:ASP:OD2	1:A:1083:TYR:OH	2.28	0.49
1:A:694:TRP:O	1:A:697:LEU:N	2.46	0.49
1:A:1036:VAL:O	1:A:1051:GLY:N	2.46	0.49
1:A:938:PHE:O	1:A:941:THR:OG1	2.29	0.48
1:A:906:LEU:O	1:A:907:THR:OG1	2.30	0.48
1:A:729:SER:O	1:A:733:GLY:N	2.48	0.45
1:A:1165:VAL:HG12	1:A:1166:GLY:N	2.32	0.45
1:A:219:PRO:O	1:A:222:GLY:N	2.49	0.45
1:A:1122:SER:OG	1:A:1161:TYR:O	2.34	0.45
1:A:745:ARG:O	1:A:749:ASN:ND2	2.50	0.45
1:A:963:GLN:HG2	1:A:964:LEU:H	1.83	0.44
1:A:495:GLU:N	1:A:495:GLU:OE1	2.51	0.44
1:A:952:CYS:HB3	1:A:974:PHE:CZ	2.53	0.44
1:A:231:ILE:O	1:A:234:SER:OG	2.36	0.43
1:A:762:SER:HA	1:A:765:THR:HG22	2.00	0.43
1:A:544:ASN:ND2	1:A:544:ASN:O	2.53	0.42
1:A:704:TRP:N	1:A:705:PRO:HD2	2.34	0.42
1:A:721:GLN:HB3	1:A:722:PRO:HD3	2.01	0.41
1:A:481:ALA:O	1:A:485:ARG:N	2.54	0.41
1:A:694:TRP:CD1	1:A:694:TRP:C	2.94	0.41
1:A:397:TYR:O	1:A:399:SER:N	2.53	0.41
1:A:893:ALA:HB2	1:A:916:TYR:HE1	1.85	0.40
1:A:372:ASP:O	1:A:373:SER:CB	2.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1183/1284 (92%)	1127 (95%)	56 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	978/1065 (92%)	974 (100%)	4 (0%)	93	97
2	B	3/3 (100%)	3 (100%)	0	100	100
All	All	981/1068 (92%)	977 (100%)	4 (0%)	93	97

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	310	PHE
1	A	700	ASN
1	A	979	PHE
1	A	1223	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1250	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	30F	B	1	2	3,5,6	3.17	2 (66%)	1,5,7	0.56	0
2	30F	B	3	2	3,5,6	3.04	2 (66%)	1,5,7	0.97	0
2	30F	B	5	2	3,5,6	3.31	2 (66%)	1,5,7	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	30F	B	1	2	-	0/0/4/6	0/0/0/0
2	30F	B	3	2	-	0/0/4/6	0/0/0/0
2	30F	B	5	2	-	0/0/4/6	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	5	30F	C10-C08	-4.88	1.37	1.45
2	B	1	30F	C10-C08	-4.72	1.37	1.45
2	B	3	30F	C10-C08	-4.60	1.37	1.45
2	B	5	30F	C08-N04	-3.00	1.27	1.35
2	B	1	30F	C08-N04	-2.78	1.28	1.35
2	B	3	30F	C08-N04	-2.57	1.28	1.35

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1187/1284 (92%)	-0.04	59 (4%) 32 21	95, 172, 293, 445	0
2	B	3/6 (50%)	1.84	1 (33%) 0 1	193, 193, 193, 193	0
All	All	1190/1290 (92%)	-0.03	60 (5%) 32 21	95, 172, 293, 445	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	704	TRP	14.7
1	A	705	PRO	7.4
1	A	700	ASN	5.7
1	A	85	SER	5.7
1	A	1278	GLU	5.3
1	A	424	ASN	5.0
1	A	522	GLU	4.8
1	A	1275	ARG	4.7
1	A	226	GLY	4.6
1	A	90	GLN	4.5
1	A	88	SER	4.5
1	A	224	SER	4.4
1	A	708	VAL	4.4
1	A	86	LYS	4.4
1	A	231	ILE	4.3
1	A	706	TYR	4.2
1	A	1276	SER	4.1
1	A	699	LEU	4.0
1	A	701	SER	4.0
1	A	702	THR	3.9
1	A	524	GLY	3.9
1	A	87	GLN	3.9
1	A	230	LYS	3.5
2	B	4	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	703	GLU	3.3
1	A	95	ASP	3.3
1	A	41	TYR	3.2
1	A	707	PHE	3.1
1	A	234	SER	3.1
1	A	227	ILE	3.1
1	A	222	GLY	3.0
1	A	1059	GLY	2.9
1	A	1219	GLU	2.9
1	A	525	ALA	2.8
1	A	223	LEU	2.7
1	A	220	VAL	2.7
1	A	42	ALA	2.6
1	A	235	PHE	2.6
1	A	883	LYS	2.6
1	A	698	LYS	2.6
1	A	75	THR	2.6
1	A	225	ALA	2.5
1	A	84	VAL	2.5
1	A	523	ARG	2.4
1	A	526	GLN	2.4
1	A	389	GLU	2.4
1	A	91	MET	2.4
1	A	221	LEU	2.4
1	A	959	LEU	2.3
1	A	709	VAL	2.3
1	A	229	ALA	2.2
1	A	228	TRP	2.1
1	A	519	LEU	2.1
1	A	695	ARG	2.1
1	A	71	PHE	2.0
1	A	89	THR	2.0
1	A	236	THR	2.0
1	A	237	ASP	2.0
1	A	1042	THR	2.0
1	A	233	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	30F	B	1	6/7	0.91	0.61	-	193,193,193,193	0
2	30F	B	3	6/7	0.86	0.71	-	193,193,193,193	0
2	30F	B	5	6/7	0.78	0.64	-	193,193,193,193	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.