



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:44 PM GMT

PDB ID : 1QFY  
Title : PEA FNR Y308S MUTANT IN COMPLEX WITH NADP+  
Authors : Deng, Z.; Aliverti, A.; Zanetti, G.; Arakaki, A.K.; Ottado, J.; Orellano, E.G.;  
Calcaterra, N.B.; Ceccarelli, E.A.; Carrillo, N.; Karplus, P.A.  
Deposited on : 1999-04-18  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	<div> <div>5%</div> <div>80%</div> <div>18%</div> <div>.</div> </div>
1	B	308	<div> <div>5%</div> <div>80%</div> <div>15%</div> <div>• •</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5708 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (FERREDOXIN: NADP+ REDUCTASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	308	Total	C	N	O	S	0	0	0
			2439	1560	403	460	16			
1	B	295	Total	C	N	O	S	0	0	0
			2341	1498	385	442	16			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	308	SER	TYR	MUTATION	UNP P10933
B	808	SER	TYR	MUTATION	UNP P10933

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
4	B	1	Total 48	C 21	N 7	O 17	P 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	388	Total 388	O 388	0	0
5	B	333	Total 333	O 333	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (FERREDOXIN: NADP+ REDUCTASE)



- Molecule 1: PROTEIN (FERREDOXIN: NADP+ REDUCTASE)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.28 Å 110.65 Å 81.04 Å 90.00° 94.38° 90.00°	Depositor
Resolution (Å)	8.00 – 1.80 22.07 – 1.80	Depositor EDS
% Data completeness (in resolution range)	80.5 (8.00-1.80) 80.8 (22.07-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.60 (at 1.80 Å)	Xtriage
Refinement program	X-PLOR 3.8	Depositor
R, $R_{free}$	0.196 , 0.257 0.196 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	21.3	Xtriage
Anisotropy	0.819	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 118.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 65922 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5708	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, SO4, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.50	0/2492	0.69	0/3356
1	B	0.46	0/2392	0.70	1/3219 (0.0%)
All	All	0.48	0/4884	0.70	1/6575 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	538	LEU	CA-CB-CG	7.36	132.23	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2439	0	2446	48	0
1	B	2341	0	2338	45	0
2	A	5	0	0	0	0
3	A	53	0	31	2	0
3	B	53	0	31	0	0
4	A	48	0	25	1	0
4	B	48	0	25	6	0
5	A	388	0	0	6	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	333	0	0	4	0
All	All	5708	0	4896	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:541:LYS:HD2	1:B:550:GLU:HG2	1.54	0.89
1:A:183:HIS:H	1:A:187:GLN:NE2	1.78	0.81
1:A:21:ILE:HD13	1:A:147:LYS:HG2	1.65	0.77
1:B:675:LEU:HD23	1:B:678:MET:HE2	1.66	0.76
4:B:810:NAP:H3D	5:B:1723:HOH:O	1.86	0.74
1:A:208:GLU:HG2	1:A:212:LYS:HE3	1.72	0.72
1:B:742:GLN:HE22	4:B:810:NAP:C2A	2.04	0.70
1:B:768:LEU:HD23	4:B:810:NAP:H4D	1.74	0.69
1:B:675:LEU:HD23	1:B:678:MET:CE	2.23	0.68
1:A:21:ILE:CD1	1:A:147:LYS:HG2	2.23	0.67
1:A:175:LEU:HD23	1:A:178:MET:CE	2.25	0.67
1:B:683:HIS:H	1:B:687:GLN:NE2	1.93	0.65
1:A:183:HIS:H	1:A:187:GLN:HE21	1.42	0.63
1:B:795:ARG:NH1	1:B:799:LYS:HZ2	1.96	0.63
1:A:295:ARG:HE	1:A:299:LYS:HE3	1.63	0.62
1:A:268:LEU:HD23	4:A:310:NAP:H4D	1.82	0.60
1:A:40:THR:HG23	5:B:1302:HOH:O	2.01	0.60
1:A:175:LEU:HD23	1:A:178:MET:HE2	1.84	0.60
1:A:70:ILE:HG23	1:A:91:ILE:HD11	1.84	0.59
1:B:740:TYR:HB3	1:B:742:GLN:HE21	1.68	0.59
1:A:216:LYS:NZ	1:A:216:LYS:HB3	2.18	0.58
1:B:741:ILE:HG22	4:B:810:NAP:H2A	1.86	0.57
1:A:27:LYS:HB3	1:A:29:LYS:NZ	2.20	0.57
1:A:16:LYS:O	1:A:153:LYS:HE3	2.05	0.57
1:B:527:LYS:HB3	1:B:529:LYS:HZ3	1.70	0.56
3:A:309:FAD:H8A	5:A:1528:HOH:O	2.05	0.56
1:A:69:SER:HB2	1:A:145:VAL:HG13	1.87	0.55
1:B:795:ARG:HD2	1:B:799:LYS:HZ1	1.71	0.55
1:B:792:GLU:O	1:B:796:THR:HG23	2.07	0.55
1:B:528:PRO:HD2	1:B:529:LYS:HZ3	1.72	0.55
1:A:269:LYS:HD2	5:A:1510:HOH:O	2.05	0.55
1:B:616:ASN:HD21	1:B:620:GLU:HB2	1.72	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:517:GLN:HE22	1:B:677:LYS:NZ	2.06	0.54
1:B:742:GLN:HE22	4:B:810:NAP:H2A	1.71	0.54
1:B:768:LEU:O	1:B:771:MET:HG2	2.08	0.54
1:A:29:LYS:HD3	1:A:29:LYS:N	2.24	0.53
1:B:546:ASP:HB2	5:B:1600:HOH:O	2.08	0.53
1:A:249:ALA:HB1	1:A:282:LEU:HD21	1.91	0.53
1:A:281:SER:O	1:A:285:LYS:HG2	2.11	0.51
1:B:740:TYR:HB3	1:B:742:GLN:NE2	2.25	0.51
1:A:212:LYS:O	1:A:216:LYS:HG3	2.11	0.50
1:A:78:ASP:HB2	5:A:1393:HOH:O	2.11	0.50
1:B:704:LEU:HD13	1:B:707:LYS:HD2	1.92	0.50
1:A:27:LYS:HB3	1:A:29:LYS:HZ2	1.76	0.50
1:A:283:ALA:HB1	1:A:288:ILE:HB	1.94	0.50
1:A:291:ILE:HD12	1:A:291:ILE:N	2.27	0.50
1:A:115:THR:HA	1:A:120:GLU:O	2.11	0.50
1:A:183:HIS:HB2	1:A:186:TYR:O	2.11	0.50
1:B:795:ARG:NH2	1:B:796:THR:HG22	2.26	0.49
1:B:683:HIS:H	1:B:687:GLN:HE21	1.57	0.49
1:A:295:ARG:HE	1:A:299:LYS:CE	2.26	0.49
1:B:800:ALA:O	1:B:801:GLU:HB2	2.13	0.49
1:B:529:LYS:HE2	1:B:529:LYS:H	1.78	0.49
1:A:208:GLU:O	1:A:212:LYS:HG3	2.14	0.48
1:A:190:GLY:O	1:A:220:ASN:HB2	2.14	0.48
1:B:527:LYS:HB3	1:B:529:LYS:NZ	2.29	0.48
1:A:152:PRO:HD3	1:A:264:TYR:OH	2.13	0.48
1:B:744:ARG:HH11	1:B:747:GLN:HE22	1.59	0.47
1:B:682:LYS:HA	1:B:687:GLN:HE22	1.79	0.47
1:B:586:LEU:C	1:B:586:LEU:HD12	2.36	0.46
1:B:744:ARG:HH11	1:B:747:GLN:NE2	2.14	0.45
1:A:30:GLU:HG3	5:A:1091:HOH:O	2.16	0.45
1:B:728:SER:O	1:B:738:LYS:HG2	2.16	0.45
1:B:732:VAL:HG13	1:B:736:GLY:HA2	1.99	0.45
1:A:28:PRO:HG2	1:A:29:LYS:HD3	1.99	0.45
1:B:570:ILE:HG23	1:B:591:ILE:HD11	1.97	0.45
1:A:28:PRO:HD2	1:A:29:LYS:HZ2	1.81	0.45
1:A:29:LYS:HD3	1:A:29:LYS:H	1.82	0.44
1:A:215:GLU:HB2	5:A:1618:HOH:O	2.17	0.44
1:B:527:LYS:N	1:B:527:LYS:HD3	2.32	0.44
1:B:802:GLN:NE2	5:B:1494:HOH:O	2.48	0.44
1:A:280:VAL:HG23	1:A:290:TRP:HB3	1.99	0.44
1:A:17:GLN:HG3	1:A:151:MET:O	2.18	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:LYS:HD2	1:A:27:LYS:N	2.33	0.44
1:A:233:ASN:HD21	1:A:237:GLU:CD	2.21	0.44
1:B:716:LYS:HE3	1:B:716:LYS:HB2	1.89	0.43
1:B:690:GLY:O	1:B:720:ASN:HB2	2.18	0.43
1:B:517:GLN:HE22	1:B:677:LYS:HZ2	1.65	0.43
1:A:21:ILE:HD12	1:A:150:LEU:HD11	1.99	0.43
1:A:5:GLU:HG3	1:A:5:GLU:O	2.19	0.43
1:A:219:GLU:HB2	5:A:1395:HOH:O	2.19	0.42
1:A:216:LYS:CB	1:A:216:LYS:NZ	2.82	0.42
1:A:58:THR:O	1:A:59:GLU:HG2	2.19	0.42
1:B:795:ARG:HH21	1:B:796:THR:HG22	1.83	0.42
1:B:527:LYS:HB3	1:B:529:LYS:HE2	2.02	0.41
1:A:46:ASP:N	1:A:46:ASP:OD1	2.48	0.41
1:A:178:MET:HE2	1:A:178:MET:HB2	1.82	0.41
1:B:742:GLN:NE2	1:B:742:GLN:H	2.18	0.41
1:B:740:TYR:CB	1:B:742:GLN:HE21	2.30	0.41
1:B:771:MET:HA	4:B:810:NAP:N1A	2.36	0.41
1:B:655:PRO:HB2	1:B:687:GLN:O	2.21	0.41
1:B:795:ARG:CZ	1:B:799:LYS:HD2	2.51	0.40
1:A:112:LEU:HD11	3:A:309:FAD:N7A	2.36	0.40
1:A:156:ASN:ND2	1:A:189:ASN:HB3	2.37	0.40
1:B:558:THR:O	1:B:561:GLU:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/308 (99%)	298 (97%)	8 (3%)	0	100	100
1	B	293/308 (95%)	285 (97%)	8 (3%)	0	100	100
All	All	599/616 (97%)	583 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/267 (100%)	255 (96%)	12 (4%)	34	16
1	B	256/267 (96%)	246 (96%)	10 (4%)	39	21
All	All	523/534 (98%)	501 (96%)	22 (4%)	36	18

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	GLU
1	A	40	THR
1	A	46	ASP
1	A	110	LYS
1	A	115	THR
1	A	184	GLU
1	A	189	ASN
1	A	232	VAL
1	A	258	LYS
1	A	278	ILE
1	A	280	VAL
1	A	292	GLU
1	B	527	LYS
1	B	530	GLU
1	B	538	LEU
1	B	540	THR
1	B	561	GLU
1	B	577	ILE
1	B	617	ASP
1	B	639	VAL
1	B	742	GLN
1	B	795	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	156	ASN
1	A	183	HIS
1	A	187	GLN
1	A	247	GLN
1	B	517	GLN
1	B	656	ASN
1	B	683	HIS
1	B	687	GLN
1	B	742	GLN
1	B	747	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	FAD	A	309	-	48,58,58	1.25	4 (8%)	54,89,89	1.49	4 (7%)
4	NAP	A	310	-	42,52,52	1.39	6 (14%)	54,80,80	2.09	10 (18%)
2	SO4	A	311	-	4,4,4	0.31	0	6,6,6	0.20	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FAD	B	809	-	48,58,58	1.34	6 (12%)	54,89,89	1.54	4 (7%)
4	NAP	B	810	-	42,52,52	1.39	4 (9%)	54,80,80	2.12	9 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	309	-	-	0/30/50/50	0/6/6/6
4	NAP	A	310	-	-	0/27/67/67	0/5/5/5
2	SO4	A	311	-	-	0/0/0/0	0/0/0/0
3	FAD	B	809	-	-	0/30/50/50	0/6/6/6
4	NAP	B	810	-	-	0/27/67/67	0/5/5/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	810	NAP	C3N-C7N	-3.69	1.44	1.50
3	A	309	FAD	P-O2P	-2.53	1.44	1.54
4	A	310	NAP	C3N-C7N	-2.40	1.46	1.50
3	B	809	FAD	C8A-N7A	-2.30	1.30	1.34
3	B	809	FAD	PA-O2A	-2.26	1.45	1.54
3	A	309	FAD	PA-O2A	-2.24	1.45	1.54
4	A	310	NAP	C4A-N3A	-2.14	1.32	1.35
4	A	310	NAP	P2B-O3X	-2.11	1.47	1.54
4	B	810	NAP	C5A-C4A	-2.09	1.35	1.40
3	B	809	FAD	C4X-N5	2.02	1.36	1.33
3	B	809	FAD	O5'-C5'	2.03	1.53	1.44
4	A	310	NAP	O4D-C1D	2.27	1.44	1.41
4	A	310	NAP	O4B-C4B	2.53	1.50	1.45
3	A	309	FAD	O4B-C1B	2.67	1.44	1.41
4	B	810	NAP	O4D-C1D	2.98	1.45	1.41
3	A	309	FAD	C4-N3	3.61	1.39	1.33
3	B	809	FAD	O4B-C1B	3.78	1.46	1.41
4	A	310	NAP	O4B-C1B	3.93	1.46	1.41
3	B	809	FAD	C4-N3	4.02	1.40	1.33
4	B	810	NAP	O4B-C1B	4.09	1.46	1.41

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	810	NAP	N3A-C2A-N1A	-10.55	120.82	128.89
4	A	310	NAP	N3A-C2A-N1A	-10.00	121.24	128.89
3	B	809	FAD	C4X-C4-N3	-5.20	116.48	123.59
3	A	309	FAD	C4X-C4-N3	-5.03	116.72	123.59
4	A	310	NAP	O2A-PA-O3	-4.20	86.02	105.09
4	A	310	NAP	C4B-O4B-C1B	-3.79	105.55	109.72
4	B	810	NAP	C4B-O4B-C1B	-3.47	105.91	109.72
4	A	310	NAP	O3-PN-O5D	-3.22	94.39	102.94
3	B	809	FAD	C2B-C1B-N9A	-3.18	109.44	114.29
4	B	810	NAP	O3-PN-O5D	-2.85	95.36	102.94
4	A	310	NAP	C1B-N9A-C4A	-2.73	122.83	126.94
4	B	810	NAP	C1B-N9A-C4A	-2.63	122.97	126.94
4	B	810	NAP	C3N-C7N-N7N	-2.36	115.23	117.82
3	A	309	FAD	C2B-C1B-N9A	-2.35	110.70	114.29
4	A	310	NAP	C3N-C7N-N7N	-2.33	115.27	117.82
3	B	809	FAD	C4X-C10-N10	-2.22	119.21	120.52
3	A	309	FAD	C4X-C10-N10	-2.01	119.34	120.52
4	A	310	NAP	O2A-PA-O1A	2.06	123.67	112.53
4	B	810	NAP	O2N-PN-O3	2.37	115.86	105.09
4	A	310	NAP	O4B-C1B-N9A	2.43	113.18	108.10
4	B	810	NAP	O4B-C1B-N9A	2.66	113.67	108.10
4	A	310	NAP	O3-PA-O5B	3.43	112.04	102.94
4	B	810	NAP	O4D-C1D-N1N	3.82	112.33	108.13
4	A	310	NAP	O4D-C1D-N1N	4.29	112.84	108.13
4	B	810	NAP	O3-PA-O5B	5.17	116.65	102.94
3	A	309	FAD	C4-N3-C2	7.50	121.73	115.25
3	B	809	FAD	C4-N3-C2	7.61	121.83	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	309	FAD	2	0
4	A	310	NAP	1	0
4	B	810	NAP	6	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	308/308 (100%)	0.11	16 (5%) 31 25	13, 26, 54, 72	0
1	B	295/308 (95%)	0.11	14 (4%) 35 29	16, 28, 59, 72	0
All	All	603/616 (97%)	0.11	30 (4%) 32 27	13, 27, 56, 72	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	13	HIS	6.6
1	A	1	GLN	4.9
1	A	12	LYS	4.3
1	A	11	VAL	4.1
1	B	617	ASP	4.0
1	A	119	GLY	3.9
1	A	117	ASP	3.8
1	A	118	ALA	3.3
1	B	795	ARG	3.1
1	B	546	ASP	3.0
1	B	799	LYS	2.8
1	A	116	ASN	2.8
1	A	299	LYS	2.8
1	B	530	GLU	2.7
1	A	234	ASP	2.6
1	B	718	PRO	2.6
1	A	29	LYS	2.3
1	B	579	LYS	2.3
1	B	788	ILE	2.3
1	B	616	ASN	2.3
1	B	529	LYS	2.2
1	A	256	LEU	2.2
1	A	182	LYS	2.1
1	A	2	VAL	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	784	ALA	2.1
1	B	577	ILE	2.1
1	B	775	ILE	2.1
1	A	215	GLU	2.1
1	A	160	ILE	2.0
1	B	618	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAP	B	810	48/48	0.94	0.11	0.04	25,40,48,58	0
3	FAD	B	809	53/53	0.95	0.10	-0.09	17,21,67,70	0
3	FAD	A	309	53/53	0.97	0.08	-0.49	14,17,54,58	0
4	NAP	A	310	48/48	0.97	0.08	-0.66	17,24,29,37	0
2	SO4	A	311	5/5	0.97	0.10	-	39,41,44,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.