



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:56 AM GMT

PDB ID : 3QF3
Title : Crystal structure of EspR transcription factor from mycobacterium tuberculosis
Authors : Blasco, B.; Pojer, F.; Cole, S.T.
Deposited on : 2011-01-21
Resolution : 2.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3386 (2.44-2.40)
Clashscore	102246	3897 (2.44-2.40)
Ramachandran outliers	100387	3837 (2.44-2.40)
Sidechain outliers	100360	3838 (2.44-2.40)
RSRZ outliers	91569	3396 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	133	 73% 26% .
1	B	133	 75% 22% . .
1	C	133	 80% 17% . .
1	D	133	 81% 14% . .
1	E	133	 80% 17% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	133	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	LMR	A	134	-	-	-	X
3	MLT	B	134	-	-	-	X
3	MLT	E	134	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ESX-1 secretion-associated regulator EspR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	131	Total	C	N	O	S	0	1	0
			1028	641	189	193	5			
1	D	128	Total	C	N	O	S	0	0	0
			1006	629	186	187	4			
1	B	131	Total	C	N	O	S	0	1	0
			1028	641	189	194	4			
1	C	129	Total	C	N	O	S	0	0	0
			1012	632	187	189	4			
1	E	131	Total	C	N	O	S	0	1	0
			1033	644	192	193	4			
1	F	130	Total	C	N	O	S	0	1	0
			1028	641	191	192	4			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	EXPRESSION TAG	UNP P96228
A	2	SER	-	EXPRESSION TAG	UNP P96228
D	1	GLY	-	EXPRESSION TAG	UNP P96228
D	2	SER	-	EXPRESSION TAG	UNP P96228
B	1	GLY	-	EXPRESSION TAG	UNP P96228
B	2	SER	-	EXPRESSION TAG	UNP P96228
C	1	GLY	-	EXPRESSION TAG	UNP P96228
C	2	SER	-	EXPRESSION TAG	UNP P96228
E	1	GLY	-	EXPRESSION TAG	UNP P96228
E	2	SER	-	EXPRESSION TAG	UNP P96228
F	1	GLY	-	EXPRESSION TAG	UNP P96228
F	2	SER	-	EXPRESSION TAG	UNP P96228

- Molecule 2 is (2S)-2-HYDROXYBUTANEDIOIC ACID (three-letter code: LMR) (formula: C₄H₆O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			9	4	5		

- Molecule 3 is D-MALATE (three-letter code: MLT) (formula: $C_4H_6O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			9	4	5		
3	E	1	Total	C	O	0	0
			9	4	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	19	Total 19	O 19	0	0
4	D	38	Total 38	O 38	0	0
4	B	46	Total 46	O 46	0	0
4	C	43	Total 43	O 43	0	0
4	E	40	Total 40	O 40	0	0
4	F	46	Total 46	O 46	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ESX-1 secretion-associated regulator EspR




- Molecule 1: ESX-1 secretion-associated regulator EspR




- Molecule 1: ESX-1 secretion-associated regulator EspR




- Molecule 1: ESX-1 secretion-associated regulator EspR



- Molecule 1: ESX-1 secretion-associated regulator EspR



- Molecule 1: ESX-1 secretion-associated regulator EspR



GLY	SER	S3	T4	R9	P19	R22	S27	T31	L34	N41	S42	A43	P44	Y45	L46	S47	Q48	R54	Y75	L85	N91	R101	Q105	R126	R127	D132	ALA
-----	-----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.21 Å 81.70 Å 124.80 Å 90.00° 95.80° 90.00°	Depositor
Resolution (Å)	36.92 – 2.41 36.92 – 2.41	Depositor EDS
% Data completeness (in resolution range)	99.6 (36.92-2.41) 99.6 (36.92-2.41)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 2.42 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.184 , 0.243 0.183 , 0.238	Depositor DCC
R_{free} test set	2013 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	44.2	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 50.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 40268 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6394	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LMR, MLT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1050	0.51	0/1419
1	B	0.40	0/1050	0.52	0/1419
1	C	0.43	0/1031	0.57	0/1393
1	D	0.40	0/1025	0.53	0/1385
1	E	0.40	0/1055	0.55	0/1425
1	F	0.40	0/1050	0.53	0/1418
All	All	0.40	0/6261	0.54	0/8459

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1028	0	1019	24	0
1	B	1028	0	1019	31	0
1	C	1012	0	1005	18	0
1	D	1006	0	1000	28	0
1	E	1033	0	1027	23	0
1	F	1028	0	1022	20	0
2	A	9	0	4	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	9	0	4	1	0
3	E	9	0	4	0	0
4	A	19	0	0	0	0
4	B	46	0	0	2	0
4	C	43	0	0	1	0
4	D	38	0	0	1	0
4	E	40	0	0	0	0
4	F	46	0	0	0	0
All	All	6394	0	6104	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:55:THR:HG22	1:E:56:ASN:N	1.72	1.05
1:E:55:THR:HG22	1:E:56:ASN:H	1.20	1.03
1:A:55:THR:HG22	1:A:56:ASN:H	1.28	0.96
1:E:35:LYS:HA	1:E:39:ILE:O	1.72	0.89
1:F:126[B]:ARG:NH2	1:F:132:ASP:HA	1.88	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	130/133 (98%)	126 (97%)	4 (3%)	0	100	100
1	B	130/133 (98%)	129 (99%)	1 (1%)	0	100	100
1	C	127/133 (96%)	127 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	126/133 (95%)	121 (96%)	5 (4%)	0	100	100
1	E	130/133 (98%)	128 (98%)	2 (2%)	0	100	100
1	F	129/133 (97%)	124 (96%)	5 (4%)	0	100	100
All	All	772/798 (97%)	755 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/105 (100%)	101 (96%)	4 (4%)	40	59
1	B	105/105 (100%)	103 (98%)	2 (2%)	65	82
1	C	103/105 (98%)	101 (98%)	2 (2%)	65	82
1	D	102/105 (97%)	96 (94%)	6 (6%)	24	37
1	E	105/105 (100%)	102 (97%)	3 (3%)	50	70
1	F	105/105 (100%)	103 (98%)	2 (2%)	65	82
All	All	625/630 (99%)	606 (97%)	19 (3%)	50	69

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	126	ARG
1	B	5	THR
1	E	37	GLU
1	D	55	THR
1	E	54	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LMR	A	134	-	1,8,8	0.65	0	2,10,10	1.50	1 (50%)
3	MLT	B	134	-	1,8,8	0.31	0	2,10,10	0.65	0
3	MLT	E	134	-	1,8,8	0.15	0	2,10,10	1.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LMR	A	134	-	-	0/2/8/8	0/0/0/0
3	MLT	B	134	-	-	0/2/8/8	0/0/0/0
3	MLT	E	134	-	-	0/2/8/8	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	134	LMR	C3-C2-C1	-2.12	108.16	111.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	134	LMR	1	0
3	B	134	MLT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	131/133 (98%)	-0.34	0 100 100	37, 63, 92, 111	0
1	B	131/133 (98%)	-0.26	0 100 100	26, 59, 89, 110	0
1	C	129/133 (96%)	-0.26	2 (1%) 74 74	22, 47, 79, 106	0
1	D	128/133 (96%)	-0.36	1 (0%) 87 87	31, 51, 85, 126	0
1	E	131/133 (98%)	-0.10	3 (2%) 64 63	30, 55, 102, 130	0
1	F	130/133 (97%)	-0.23	0 100 100	34, 51, 83, 108	0
All	All	780/798 (97%)	-0.26	6 (0%) 87 87	22, 55, 92, 130	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	131	ILE	2.9
1	E	37	GLU	2.7
1	E	39	ILE	2.5
1	C	130	GLY	2.5
1	E	93	CYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	LMR	A	134	9/9	0.77	0.21	5.16	77,95,100,103	0
3	MLT	E	134	9/9	0.82	0.26	3.04	72,81,88,91	0
3	MLT	B	134	9/9	0.91	0.19	2.34	48,67,84,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.