



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:56 PM GMT

PDB ID : 4QFG
Title : Structure of AMPK in complex with STAUROSPORINE inhibitor and in the absence of a synthetic activator
Authors : Calabrese, M.F.; Kurumbail, R.G.
Deposited on : 2014-05-20
Resolution : 3.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 3.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1000 (3.56-3.36)
Clashscore	102246	1090 (3.56-3.36)
Ramachandran outliers	100387	1057 (3.56-3.36)
Sidechain outliers	100360	1058 (3.56-3.36)
RSRZ outliers	91569	1005 (3.56-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	503	
2	B	204	
3	C	330	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SO4	C	402	-	-	-	X
6	SO4	C	403	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6593 atoms, of which 26 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5'-AMP-activated protein kinase catalytic subunit alpha-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	366	Total	C	N	O	P	S	0	0	0
			2970	1905	518	528	1	18			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP P54645
A	517	ALA	-	SEE REMARK 999	UNP P54645
A	518	SER	-	SEE REMARK 999	UNP P54645
A	519	GLY	-	SEE REMARK 999	UNP P54645
A	520	GLY	-	SEE REMARK 999	UNP P54645
A	521	PRO	-	SEE REMARK 999	UNP P54645
A	522	GLY	-	SEE REMARK 999	UNP P54645
A	523	GLY	-	SEE REMARK 999	UNP P54645
A	524	SER	-	SEE REMARK 999	UNP P54645

- Molecule 2 is a protein called 5'-AMP-activated protein kinase subunit beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	161	Total	C	N	O	S	0	0	0
			1271	828	215	225	3			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	67	MET	-	EXPRESSION TAG	UNP P80386
B	108	ASP	SER	ENGINEERED MUTATION	UNP P80386

- Molecule 3 is a protein called 5'-AMP-activated protein kinase subunit gamma-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	287	Total	C	N	O	S	0	0	0
			2250	1463	376	404	7			

- Molecule 4 is STAUROSPORINE (three-letter code: STU) (formula: $C_{28}H_{26}N_4O_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	0	0
			61	28	26	4	3		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total 3 Cl 3	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: $C_{10}H_{14}N_5O_7P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-AMP-activated protein kinase catalytic subunit alpha-1




- Molecule 2: 5'-AMP-activated protein kinase subunit beta-1



- Molecule 3: 5'-AMP-activated protein kinase subunit gamma-1





4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	124.37Å 124.37Å 404.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.86 – 3.46 29.86 – 3.46	Depositor EDS
% Data completeness (in resolution range)	97.0 (29.86-3.46) 96.3 (29.86-3.46)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.42 (at 3.47Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.222 , 0.267 0.217 , 0.266	Depositor DCC
R_{free} test set	1208 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	79.2	Xtriage
Anisotropy	0.448	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 85.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 24416 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6593	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, STU, CL, SO4, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/3026	0.76	0/4081
2	B	0.48	0/1306	0.73	0/1779
3	C	0.47	0/2294	0.66	0/3119
All	All	0.48	0/6626	0.72	0/8979

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2970	0	2996	48	0
2	B	1271	0	1268	19	0
3	C	2250	0	2283	24	0
4	A	35	26	26	6	0
5	A	3	0	0	0	0
6	A	5	0	0	0	0
6	C	10	0	0	0	0
7	C	23	0	12	0	0
All	All	6567	26	6585	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:79:PRO:HB2	2:B:115:ILE:HG22	1.44	0.99
4:A:601:STU:H16	4:A:601:STU:H261	1.60	0.82
2:B:120:GLU:HA	2:B:155:VAL:HG22	1.72	0.70
1:A:51:LYS:O	1:A:54:SER:HB3	1.96	0.65
1:A:141:LYS:HG3	1:A:143:GLU:HB2	1.79	0.64
1:A:185:ILE:HD11	1:A:227:CYS:SG	2.38	0.62
1:A:77:ILE:HG21	4:A:601:STU:H91	1.81	0.62
1:A:401:ARG:O	1:A:548:GLN:HB3	2.00	0.62
3:C:69:ARG:HE	3:C:151:ARG:HH12	1.48	0.60
1:A:12:LYS:HB3	1:A:17:ILE:HA	1.85	0.59
3:C:263:LYS:HA	3:C:266:GLN:HE21	1.69	0.57
1:A:540:ALA:HB2	2:B:251:LEU:HD11	1.87	0.56
1:A:160:LEU:HD13	1:A:174:CYS:HB2	1.88	0.56
3:C:104:GLN:HA	3:C:255:ASN:HD21	1.72	0.54
3:C:184:LYS:H	3:C:184:LYS:HE3	1.72	0.54
1:A:218:HIS:CE1	1:A:220:PRO:HD2	2.43	0.54
3:C:73:LEU:HD21	3:C:85:LEU:HB2	1.90	0.53
2:B:145:GLN:H	2:B:145:GLN:CD	2.11	0.53
3:C:42:PRO:HG2	3:C:45:SER:HB3	1.91	0.53
2:B:264:THR:HG23	3:C:47:LEU:HD23	1.91	0.53
1:A:193:PRO:O	1:A:197:ILE:HG12	2.09	0.53
1:A:49:ARG:HH21	1:A:86:PRO:HA	1.72	0.53
1:A:398:LEU:HD22	2:B:210:LEU:HB3	1.91	0.52
1:A:12:LYS:HD3	1:A:17:ILE:HG22	1.92	0.51
1:A:171:ARG:HH22	2:B:204:PRO:HG3	1.75	0.51
3:C:248:LEU:HA	3:C:253:THR:HG23	1.92	0.51
3:C:243:PHE:O	3:C:246:ILE:HG13	2.10	0.51
3:C:179:ILE:HD11	3:C:184:LYS:HA	1.93	0.50
1:A:258:THR:H	1:A:261:ASP:HB2	1.76	0.50
4:A:601:STU:H272	4:A:601:STU:N4	2.26	0.50
1:A:127:VAL:CG2	1:A:140:LEU:HD21	2.42	0.50
1:A:273:PRO:HD2	1:A:276:LEU:HD12	1.93	0.49
1:A:237:LEU:HB3	1:A:242:ILE:HD11	1.94	0.49
1:A:247:HIS:CG	1:A:257:ALA:HB2	2.48	0.48
3:C:212:VAL:O	3:C:216:LEU:HG	2.13	0.48
1:A:530:SER:HB2	3:C:161:ASN:HB3	1.96	0.48
1:A:414:CYS:HA	1:A:417:ILE:HD12	1.96	0.48
1:A:179:TYR:CD2	1:A:202:VAL:CG2	2.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:ARG:O	1:A:172:TPO:C	2.61	0.48
2:B:242:LEU:HB2	2:B:250:VAL:HB	1.96	0.48
3:C:302:VAL:HA	3:C:308:VAL:HA	1.96	0.48
1:A:77:ILE:HG12	1:A:154:LYS:HB3	1.95	0.47
1:A:465:SER:H	2:B:237:ASN:HB3	1.79	0.47
1:A:463:PHE:HB2	2:B:239:LEU:HB3	1.96	0.47
1:A:218:HIS:HD2	1:A:221:THR:HG23	1.79	0.47
1:A:143:GLU:HB3	4:A:601:STU:H281	1.96	0.47
1:A:254:MET:CG	2:B:208:PRO:HB2	2.45	0.47
3:C:216:LEU:O	3:C:220:VAL:HG23	2.15	0.46
2:B:84:TRP:HB3	2:B:112:PHE:HB2	1.97	0.46
3:C:187:PHE:HA	3:C:190:LYS:HD2	1.96	0.46
1:A:179:TYR:CD2	1:A:202:VAL:HG23	2.51	0.45
4:A:601:STU:C16	4:A:601:STU:H261	2.29	0.45
1:A:78:LYS:H	1:A:94:GLU:HG2	1.81	0.45
3:C:89:ASP:OD2	3:C:116:TRP:HZ3	1.99	0.45
1:A:55:LEU:HB2	1:A:57:VAL:HG22	1.99	0.45
3:C:27:VAL:HG11	3:C:182:PHE:CZ	2.51	0.45
3:C:278:CYS:HB3	3:C:299:LEU:HD13	1.99	0.45
2:B:145:GLN:NE2	2:B:145:GLN:H	2.14	0.45
3:C:144:LEU:HD23	3:C:149:ILE:HG23	1.98	0.44
1:A:433:ARG:HE	1:A:466:ILE:HD13	1.82	0.44
3:C:72:PRO:HD3	3:C:165:ILE:HD11	1.98	0.44
1:A:61:ILE:HD11	2:B:166:LEU:HD11	2.00	0.44
3:C:205:MET:HA	3:C:228:PRO:HD2	1.99	0.44
3:C:41:ILE:HD11	3:C:154:VAL:HG11	2.00	0.44
3:C:48:VAL:O	3:C:71:ALA:HB1	2.18	0.43
1:A:434:VAL:HG13	1:A:447:MET:HG3	2.01	0.43
4:A:601:STU:C16	4:A:601:STU:C26	2.95	0.43
2:B:166:LEU:HA	2:B:166:LEU:HD12	1.91	0.43
1:A:132:ARG:HH21	1:A:278:PRO:HG2	1.84	0.42
1:A:48:ASN:HA	1:A:88:ASP:OD2	2.19	0.42
2:B:110:ASN:ND2	2:B:110:ASN:H	2.17	0.42
1:A:179:TYR:HD2	1:A:202:VAL:CG2	2.31	0.42
3:C:88:THR:HA	3:C:91:ILE:HD12	2.02	0.42
1:A:132:ARG:NH2	1:A:278:PRO:HG2	2.35	0.42
1:A:35:HIS:CE1	1:A:37:LEU:HB2	2.55	0.42
1:A:49:ARG:O	1:A:53:ARG:HB3	2.20	0.41
1:A:179:TYR:HD2	1:A:202:VAL:HG23	1.85	0.41
1:A:71:PHE:CD1	1:A:129:TYR:HE1	2.38	0.41
2:B:93:LEU:HB2	2:B:105:LEU:HD21	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:GLY:O	1:A:41:LYS:HA	2.21	0.41
1:A:539:CYS:HB3	2:B:239:LEU:HD22	2.03	0.41
2:B:79:PRO:HA	2:B:117:ASP:HA	2.03	0.41
1:A:127:VAL:HG22	1:A:140:LEU:HD21	2.02	0.41
1:A:138:ARG:O	1:A:139:ASP:HB3	2.21	0.41
1:A:137:HIS:HE1	1:A:139:ASP:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/503 (71%)	333 (93%)	22 (6%)	4 (1%)	17	62
2	B	155/204 (76%)	143 (92%)	10 (6%)	2 (1%)	15	58
3	C	281/330 (85%)	265 (94%)	15 (5%)	1 (0%)	39	80
All	All	795/1037 (77%)	741 (93%)	47 (6%)	7 (1%)	21	67

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	108	ASP
3	C	303	ASP
1	A	139	ASP
2	B	223	CYS
1	A	399	GLY
1	A	151	MET
1	A	427	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/448 (73%)	288 (88%)	40 (12%)	6	28
2	B	140/185 (76%)	121 (86%)	19 (14%)	5	24
3	C	247/299 (83%)	223 (90%)	24 (10%)	10	41
All	All	715/932 (77%)	632 (88%)	83 (12%)	7	31

All (83) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	ASP
1	A	22	LEU
1	A	29	LYS
1	A	32	VAL
1	A	52	ILE
1	A	53	ARG
1	A	70	LEU
1	A	85	THR
1	A	107	LYS
1	A	110	ARG
1	A	112	ASP
1	A	138	ARG
1	A	147	LEU
1	A	170	LEU
1	A	178	ASN
1	A	185	ILE
1	A	189	LEU
1	A	207	LEU
1	A	215	ASP
1	A	219	VAL
1	A	222	LEU
1	A	224	LYS
1	A	233	THR
1	A	251	VAL
1	A	272	LEU
1	A	401	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	402	SER
1	A	403	GLN
1	A	405	ARG
1	A	418	LYS
1	A	432	LEU
1	A	433	ARG
1	A	434	VAL
1	A	436	ARG
1	A	457	ARG
1	A	458	THR
1	A	537	GLU
1	A	538	MET
1	A	545	ILE
1	A	548	GLN
2	B	90	GLU
2	B	94	SER
2	B	110	ASN
2	B	111	ASN
2	B	116	LEU
2	B	117	ASP
2	B	134	THR
2	B	145	GLN
2	B	155	VAL
2	B	158	THR
2	B	166	LEU
2	B	167	MET
2	B	170	SER
2	B	205	ILE
2	B	230	GLU
2	B	234	VAL
2	B	262	VAL
2	B	263	THR
2	B	264	THR
3	C	56	VAL
3	C	84	MET
3	C	88	THR
3	C	108	LEU
3	C	111	HIS
3	C	121	LEU
3	C	128	LEU
3	C	137	LEU
3	C	145	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	161	ASN
3	C	180	THR
3	C	184	LYS
3	C	189	SER
3	C	192	LEU
3	C	207	ARG
3	C	208	THR
3	C	225	SER
3	C	238	ASP
3	C	246	ILE
3	C	255	ASN
3	C	284	LEU
3	C	287	ILE
3	C	291	LEU
3	C	318	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	218	HIS
2	B	109	GLN
2	B	110	ASN
2	B	145	GLN
2	B	232	ASN
2	B	233	HIS
3	C	255	ASN
3	C	266	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	172	1	8,10,11	1.30	1 (12%)	7,14,16	2.10	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	172	1	-	0/8/11/13	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	172	TPO	CG2-CB	2.01	1.56	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	172	TPO	O3P-P-O2P	-3.28	94.91	107.38
1	A	172	TPO	OG1-P-O1P	-2.22	101.55	107.11
1	A	172	TPO	CG2-CB-CA	2.79	118.83	113.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	172	TPO	1	0

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	STU	A	601	-	27,42,42	2.31	9 (33%)	23,68,68	1.82	8 (34%)
6	SO4	A	605	-	4,4,4	0.22	0	6,6,6	0.15	0
7	AMP	C	401	-	20,25,25	0.62	0	22,38,38	0.75	0
6	SO4	C	402	-	4,4,4	0.22	0	6,6,6	0.28	0
6	SO4	C	403	-	4,4,4	0.19	0	6,6,6	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	STU	A	601	-	-	0/4/42/42	0/0/8/8
6	SO4	A	605	-	-	0/0/0/0	0/0/0/0
7	AMP	C	401	-	-	0/6/26/26	0/3/3/3
6	SO4	C	402	-	-	0/0/0/0	0/0/0/0
6	SO4	C	403	-	-	0/0/0/0	0/0/0/0

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	601	STU	O5-C8	2.79	1.28	1.23
4	A	601	STU	C7-C10	2.87	1.47	1.40
4	A	601	STU	C6-C19	3.16	1.46	1.42
4	A	601	STU	C19-C18	3.30	1.49	1.41
4	A	601	STU	C7-C6	3.65	1.49	1.43
4	A	601	STU	C11-C18	3.95	1.47	1.42
4	A	601	STU	C10-C11	4.43	1.49	1.42
4	A	601	STU	C5-C20	4.49	1.48	1.41
4	A	601	STU	C12-C17	4.61	1.49	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	601	STU	C1-C20-C5	-2.85	116.61	120.73
4	A	601	STU	C16-C17-C12	-2.64	116.92	120.73
4	A	601	STU	O5-C8-C7	-2.20	124.95	128.62
4	A	601	STU	C3-C4-C5	-2.01	117.31	120.79
4	A	601	STU	C1-C20-N3	2.04	134.64	132.18
4	A	601	STU	C13-C12-C17	2.64	122.73	119.39
4	A	601	STU	C4-C5-C20	2.98	123.16	119.39
4	A	601	STU	C16-C17-N2	4.49	137.70	132.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	601	STU	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	365/503 (72%)	0.09	10 (2%) 58 51	38, 70, 125, 160	0
2	B	161/204 (78%)	0.33	8 (4%) 32 27	52, 80, 116, 144	0
3	C	287/330 (86%)	0.08	8 (2%) 56 50	65, 95, 126, 173	0
All	All	813/1037 (78%)	0.13	26 (3%) 51 45	38, 84, 125, 173	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	469	GLU	4.7
1	A	278	PRO	3.8
2	B	222	SER	3.6
1	A	444	PHE	3.3
2	B	201	LYS	3.3
1	A	468	ASP	3.2
3	C	237	VAL	3.1
2	B	223	CYS	2.9
3	C	122	GLN	2.9
3	C	186	GLU	2.8
1	A	9	GLY	2.7
2	B	224	ASP	2.5
3	C	196	GLN	2.5
1	A	441	THR	2.4
1	A	223	PHE	2.4
3	C	267	HIS	2.4
3	C	25	SER	2.4
2	B	270	ILE	2.4
1	A	70	LEU	2.3
2	B	216	ASN	2.3
1	A	54	SER	2.3
2	B	217	LYS	2.2
3	C	306	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	234	GLY	2.2
2	B	123	HIS	2.1
1	A	185	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	A	172	11/12	0.93	0.18	-	74,76,84,87	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	SO4	C	403	5/5	0.69	0.42	6.02	145,145,145,146	0
6	SO4	C	402	5/5	0.83	0.48	5.08	144,145,145,146	0
7	AMP	C	401	23/23	0.79	0.32	1.96	139,151,154,156	0
6	SO4	A	605	5/5	0.96	0.21	1.27	76,78,79,79	0
4	STU	A	601	35/35	0.96	0.24	0.70	41,50,59,61	0
5	CL	A	602	1/1	0.95	0.18	-	65,65,65,65	0
5	CL	A	604	1/1	0.90	0.19	-	52,52,52,52	0
5	CL	A	603	1/1	0.63	0.16	-	95,95,95,95	0

6.5 Other polymers

There are no such residues in this entry.