



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:47 PM GMT

PDB ID : 1QNA
Title : CRYSTAL STRUCTURE OF THE T(-30) ADENOVIRUS MAJOR LATE PROMOTER TATA BOX VARIANT BOUND TO WILD-TYPE TBP (ARABIDOPSIS THALIANA TBP ISOFORM 2). TATA ELEMENT RECOGNITION BY THE TATA BOX-BINDING PROTEIN HAS BEEN CONSERVED THROUGHOUT EVOLUTION.
Authors : Patikoglou, G.A.; Kim, J.L.; Sun, L.; Yang, S.-H.; Kodadek, T.; Burley, S.K.
Deposited on : 1999-10-14
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	200	
1	B	200	
2	C	14	
2	E	14	
3	D	14	
3	F	14	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4514 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTION INITIATION FACTOR TFIID-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	1	0
			1431	935	241	247	8			
1	B	186	Total	C	N	O	S	0	1	0
			1471	959	250	254	8			

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*CP*TP*TP*TP*AP*AP*AP*AP*GP*GP*GP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	14	Total	C	N	O	P	0	0	0
			286	137	57	79	13			
2	E	14	Total	C	N	O	P	0	0	0
			288	138	57	80	13			

- Molecule 3 is a DNA chain called DNA (5'-D(*TP*GP*CP*CP*CP*TP*TP*TP*TP*AP*AP*AP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	14	Total	C	N	O	P	0	0	0
			280	136	47	84	13			
3	F	14	Total	C	N	O	P	0	0	0
			280	136	47	84	13			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	146	Total	O	0	0
			146	146		
4	B	158	Total	O	0	0
			158	158		
4	C	44	Total	O	0	0
			44	44		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	40	Total 40	O 40	0	0
4	E	43	Total 43	O 43	0	0
4	F	47	Total 47	O 47	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: TRANSCRIPTION INITIATION FACTOR TFIIID-1

Chain A: 



- Molecule 1: TRANSCRIPTION INITIATION FACTOR TFIIID-1

Chain B: 



- Molecule 2: DNA (5'-D(*GP*CP*TP*TP*TP*AP*AP*AP*AP*GP*GP*GP*CP*A)-3')

Chain C: 



- Molecule 2: DNA (5'-D(*GP*CP*TP*TP*TP*AP*AP*AP*AP*GP*GP*GP*CP*A)-3')

Chain E: 



- Molecule 3: DNA (5'-D(*TP*GP*CP*CP*CP*TP*TP*TP*TP*AP*AP*AP*GP*C)-3')

Chain D: 



- Molecule 3: DNA (5'-D(*TP*GP*CP*CP*CP*TP*TP*TP*TP*AP*AP*AP*GP*C)-3')

Chain F:  36% 43% 21%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.80 Å 146.70 Å 57.40 Å 90.00° 90.50° 90.00°	Depositor
Resolution (Å)	6.00 – 1.80	Depositor
% Data completeness (in resolution range)	98.1 (6.00-1.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.193 , 0.239	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4514	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/1465	0.66	0/1975
1	B	0.46	0/1505	0.66	0/2026
2	C	1.06	2/322 (0.6%)	1.58	6/496 (1.2%)
2	E	1.06	1/324 (0.3%)	1.58	5/499 (1.0%)
3	D	1.24	1/312 (0.3%)	1.85	13/479 (2.7%)
3	F	1.22	1/312 (0.3%)	1.84	16/479 (3.3%)
All	All	0.73	5/4240 (0.1%)	1.12	40/5954 (0.7%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	222	DT	C5-C7	6.71	1.54	1.50
2	C	204	DT	C5-C7	5.95	1.53	1.50
2	E	204	DT	C5-C7	5.34	1.53	1.50
3	D	221	DT	C5-C7	5.20	1.53	1.50
2	C	205	DT	C5-C7	5.03	1.53	1.50

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	217	DC	O4'-C1'-N1	9.34	114.54	108.00
2	C	204	DT	C6-C5-C7	-7.44	118.43	122.90
3	F	216	DG	O4'-C4'-C3'	-7.14	101.64	104.50
3	D	216	DG	P-O3'-C3'	6.77	127.82	119.70
3	F	222	DT	C6-C5-C7	-6.67	118.90	122.90
3	D	220	DT	C6-C5-C7	-6.60	118.94	122.90
2	E	204	DT	C6-C5-C7	-6.39	119.06	122.90
3	D	223	DT	O4'-C4'-C3'	-6.38	101.95	104.50
3	F	215	DT	C6-C5-C7	-6.37	119.08	122.90
3	D	217	DC	C1'-O4'-C4'	-6.32	103.78	110.10
3	D	222	DT	C6-C5-C7	-6.29	119.12	122.90
3	F	218	DC	O4'-C1'-N1	6.22	112.36	108.00
2	E	203	DT	C6-C5-C7	-6.15	119.21	122.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	205	DT	C6-C5-C7	-6.13	119.22	122.90
3	F	223	DT	C6-C5-C7	-6.10	119.24	122.90
3	D	223	DT	C6-C5-C7	-6.09	119.24	122.90
3	F	221	DT	C6-C5-C7	-6.07	119.26	122.90
3	F	220	DT	C6-C5-C7	-6.06	119.26	122.90
3	D	221	DT	C6-C5-C7	-6.00	119.30	122.90
3	F	215	DT	C1'-O4'-C4'	-5.88	104.22	110.10
2	E	205	DT	C6-C5-C7	-5.85	119.39	122.90
3	F	216	DG	O4'-C1'-N9	5.85	112.09	108.00
3	F	217	DC	O4'-C1'-N1	5.77	112.04	108.00
2	E	213	DC	O4'-C1'-C2'	-5.73	101.32	105.90
2	C	203	DT	C6-C5-C7	-5.66	119.50	122.90
3	D	215	DT	C6-C5-C7	-5.62	119.53	122.90
3	D	217	DC	O4'-C1'-C2'	-5.35	101.62	105.90
2	E	214	DA	O4'-C1'-C2'	-5.24	101.71	105.90
3	F	215	DT	C4-C5-C6	5.24	121.14	118.00
3	F	225	DA	O4'-C1'-N9	5.24	111.67	108.00
3	D	223	DT	C4-C5-C6	5.20	121.12	118.00
2	C	209	DA	O4'-C1'-N9	5.17	111.62	108.00
3	F	215	DT	P-O3'-C3'	5.17	125.90	119.70
3	F	216	DG	P-O3'-C3'	5.16	125.90	119.70
3	D	218	DC	O4'-C1'-N1	5.14	111.60	108.00
3	D	215	DT	O4'-C4'-C3'	-5.13	102.45	104.50
3	F	223	DT	C4-C5-C6	5.10	121.06	118.00
2	C	213	DC	O4'-C1'-C2'	-5.06	101.85	105.90
2	C	213	DC	O4'-C1'-N1	5.05	111.54	108.00
3	F	220	DT	C4-C5-C6	5.04	121.02	118.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1431	0	1499	13	0
1	B	1471	0	1552	24	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	286	0	155	3	0
2	E	288	0	159	4	0
3	D	280	0	161	5	0
3	F	280	0	161	2	0
4	A	146	0	0	2	0
4	B	158	0	0	4	0
4	C	44	0	0	0	0
4	D	40	0	0	2	0
4	E	43	0	0	1	0
4	F	47	0	0	1	0
All	All	4514	0	3687	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:13:ASP:HB3	1:B:16:LYS:HE2	1.51	0.93
1:B:49:ASN:N	4:B:2033:HOH:O	1.95	0.90
1:A:158:PRO:HD3	1:A:183:LYS:HD2	1.63	0.81
1:B:129:ARG:HG3	1:B:197:LYS:HB3	1.66	0.77
1:B:99:ARG:HD2	1:B:103:LYS:HE2	1.66	0.75
1:B:166[A]:VAL:HG22	4:F:2036:HOH:O	1.91	0.69
1:B:49:ASN:ND2	4:B:2030:HOH:O	2.25	0.69
1:B:129:ARG:CG	1:B:197:LYS:HB3	2.22	0.68
1:B:46:GLN:NE2	4:B:2024:HOH:O	2.26	0.66
3:D:215:DT:H5''	3:D:215:DT:H6	1.65	0.61
1:A:166[B]:VAL:HG23	4:A:2123:HOH:O	2.02	0.60
1:A:102:GLN:HG3	1:A:108:ALA:HB3	1.87	0.57
1:B:34:LEU:HD12	1:B:38:LEU:HD11	1.89	0.55
1:A:49:ASN:O	1:A:62:MET:HA	2.08	0.54
1:B:161:VAL:HG21	2:E:206:DA:H5'	1.90	0.53
1:A:59:ALA:HB2	1:A:74:PHE:CE2	2.45	0.52
1:A:51:GLU:HG2	1:A:61:ILE:HB	1.92	0.50
1:A:49:ASN:ND2	1:A:49:ASN:H	2.09	0.50
3:D:216:DG:H4'	3:D:217:DC:OP1	2.12	0.50
4:A:2123:HOH:O	3:D:226:DA:H5''	2.13	0.49
1:B:47:ALA:HB2	1:B:104:LEU:HD11	1.95	0.48
1:A:99:ARG:O	1:A:103:LYS:HG3	2.13	0.48
1:B:127:PRO:HB2	1:B:197:LYS:HG3	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:203:DT:C6	2:C:204:DT:H72	2.49	0.47
1:B:13:ASP:CB	1:B:16:LYS:HE2	2.35	0.47
1:B:166[B]:VAL:HG13	1:B:197:LYS:HD3	1.95	0.47
1:B:23:PRO:HA	1:B:122:CYS:HB3	1.96	0.46
1:B:37:LYS:HA	1:B:37:LYS:HD3	1.68	0.45
3:D:215:DT:H5''	3:D:215:DT:C6	2.50	0.45
3:F:216:DG:H2''	3:F:217:DC:O5'	2.18	0.44
1:B:66:GLU:O	1:B:96:LYS:HE2	2.17	0.44
1:A:169:LYS:HE3	4:D:2034:HOH:O	2.19	0.43
2:E:213:DC:H2''	2:E:214:DA:C8	2.54	0.43
1:A:51:GLU:CG	1:A:61:ILE:HB	2.48	0.43
1:B:59:ALA:HB2	1:B:74:PHE:CE2	2.53	0.43
2:C:210:DG:N3	2:C:210:DG:H2'	2.34	0.43
1:B:129:ARG:HG2	1:B:197:LYS:HB3	2.00	0.42
2:E:214:DA:H1'	4:E:2043:HOH:O	2.19	0.42
2:C:202:DC:H2'	2:C:203:DT:C6	2.55	0.42
1:A:19:SER:HA	1:B:14:LEU:HB2	2.02	0.42
1:B:35:ASP:O	1:B:36:CYS:HB3	2.20	0.42
1:B:31:THR:HG23	1:B:116:GLN:HG3	2.02	0.42
1:B:48:ARG:HA	4:B:2033:HOH:O	2.19	0.41
1:A:183:LYS:NZ	1:A:187:ASN:HD21	2.17	0.41
2:E:210:DG:H2'	2:E:210:DG:N3	2.36	0.41
3:D:227:DG:H1'	4:D:2039:HOH:O	2.21	0.41
3:F:215:DT:H2''	3:F:216:DG:O5'	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	181/200 (90%)	176 (97%)	5 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	185/200 (92%)	179 (97%)	5 (3%)	1 (0%)	34	17
All	All	366/400 (92%)	355 (97%)	10 (3%)	1 (0%)	46	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	58	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/171 (90%)	152 (99%)	2 (1%)	76	68
1	B	160/171 (94%)	157 (98%)	3 (2%)	65	52
All	All	314/342 (92%)	309 (98%)	5 (2%)	70	59

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	ASN
1	A	104	LEU
1	B	49	ASN
1	B	125	LYS
1	B	197	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	ASN
1	A	187	ASN
1	B	53	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.