



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:47 PM GMT

PDB ID : 1QNL  
Title : AMIDE RECEPTOR/NEGATIVE REGULATOR OF THE AMIDASE  
OPERON OF PSEUDOMONAS AERUGINOSA (AMIC) COMPLEXED  
WITH BUTYRAMIDE  
Authors : Pearl, L.H.; O'Hara, B.P.; Roe, S.M.  
Deposited on : 1999-10-19  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	<div> <div>7%</div> <div>59%</div> <div>26%</div> <div>8%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BMD	A	1001	-	X	X	-

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2963 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ALIPHATIC AMIDASE EXPRESSION-REGULATING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	259	0	0
			2914	1835	526	545	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	GLN	HIS	CONFLICT	UNP P27017
A	28	ARG	ALA	CONFLICT	UNP P27017

- Molecule 2 is BUTYRAMIDE (three-letter code: BMD) (formula:  $C_4H_9NO$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			6	4	1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	43	Total 43	O 43	5	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.15Å 104.15Å 65.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.70 26.84 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.0 (15.00-2.70) 97.1 (26.84-2.70)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.08 (at 2.72Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.269 , 0.314 0.242 , 0.295	Depositor DCC
$R_{free}$ test set	477 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.1	Xtriage
Anisotropy	0.170	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 57.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 10077 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	2963	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.86	14/2989 (0.5%)	1.99	77/4074 (1.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	13

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	261	ASP	C-N	-15.77	0.97	1.34
1	A	221	TYR	C-N	-13.55	1.08	1.33
1	A	307	ARG	C-N	11.81	1.61	1.34
1	A	67	ALA	C-N	10.73	1.58	1.34
1	A	317	ASP	C-N	-9.62	1.11	1.34
1	A	72	ARG	C-N	9.46	1.55	1.34
1	A	50	GLU	C-N	-9.30	1.12	1.34
1	A	328	ARG	C-N	-8.29	1.15	1.34
1	A	327	VAL	C-N	-7.37	1.17	1.34
1	A	68	GLU	C-N	-7.29	1.17	1.34
1	A	37	GLN	CB-CG	-7.29	1.32	1.52
1	A	358	PRO	C-N	-5.88	1.20	1.34
1	A	136	ILE	C-N	-5.77	1.20	1.34
1	A	71	ILE	C-N	5.37	1.46	1.34

All (77) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	221	TYR	O-C-N	-28.43	74.87	123.20
1	A	328	ARG	O-C-N	-26.59	80.16	122.70
1	A	317	ASP	O-C-N	-22.25	87.10	122.70
1	A	39	ASN	O-C-N	-20.12	90.51	122.70
1	A	328	ARG	C-N-CA	19.57	170.62	121.70
1	A	266	ARG	NE-CZ-NH2	-18.16	111.22	120.30
1	A	346	ASP	O-C-N	-17.74	94.31	122.70
1	A	74	ARG	NE-CZ-NH1	-17.52	111.54	120.30
1	A	47	ARG	NE-CZ-NH1	17.39	129.00	120.30
1	A	328	ARG	CA-C-N	15.74	151.82	117.20
1	A	348	ARG	NE-CZ-NH2	15.73	128.16	120.30
1	A	260	ILE	O-C-N	-14.27	99.87	122.70
1	A	261	ASP	C-N-CA	13.96	156.60	121.70
1	A	261	ASP	CB-CG-OD1	13.44	130.40	118.30
1	A	193	ILE	O-C-N	-12.55	102.62	122.70
1	A	50	GLU	O-C-N	-12.17	103.23	122.70
1	A	50	GLU	C-N-CA	11.77	151.13	121.70
1	A	68	GLU	O-C-N	-10.65	105.65	122.70
1	A	337	ARG	NE-CZ-NH2	-9.79	115.40	120.30
1	A	50	GLU	OE1-CD-OE2	-9.79	111.55	123.30
1	A	165	ARG	CD-NE-CZ	9.75	137.25	123.60
1	A	328	ARG	CD-NE-CZ	9.62	137.07	123.60
1	A	266	ARG	NE-CZ-NH1	9.59	125.09	120.30
1	A	328	ARG	NE-CZ-NH2	9.54	125.07	120.30
1	A	47	ARG	N-CA-CB	9.01	126.81	110.60
1	A	68	GLU	C-N-CA	8.85	143.83	121.70
1	A	47	ARG	CD-NE-CZ	8.71	135.79	123.60
1	A	61	ASP	CB-CG-OD1	8.69	126.12	118.30
1	A	261	ASP	O-C-N	-8.61	108.93	122.70
1	A	348	ARG	NE-CZ-NH1	-8.42	116.09	120.30
1	A	215	ARG	NE-CZ-NH1	8.38	124.49	120.30
1	A	137	ARG	CD-NE-CZ	8.31	135.24	123.60
1	A	41	GLU	CG-CD-OE2	8.29	134.88	118.30
1	A	327	VAL	O-C-N	-8.23	109.53	122.70
1	A	319	ASP	O-C-N	7.89	135.32	122.70
1	A	62	ARG	NE-CZ-NH2	-7.63	116.48	120.30
1	A	307	ARG	C-N-CA	-7.54	102.84	121.70
1	A	266	ARG	CD-NE-CZ	7.43	134.00	123.60
1	A	328	ARG	NH1-CZ-NH2	-7.35	111.31	119.40
1	A	331	ARG	CD-NE-CZ	7.31	133.83	123.60
1	A	221	TYR	CA-C-N	7.22	130.65	116.20
1	A	154	ARG	NE-CZ-NH1	7.14	123.87	120.30
1	A	50	GLU	CA-C-N	7.06	132.73	117.20

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	161	ARG	NE-CZ-NH1	-7.05	116.78	120.30
1	A	74	ARG	NH1-CZ-NH2	7.02	127.12	119.40
1	A	41	GLU	OE1-CD-OE2	-7.01	114.89	123.30
1	A	68	GLU	OE1-CD-OE2	6.99	131.69	123.30
1	A	72	ARG	CB-CG-CD	6.91	129.57	111.60
1	A	261	ASP	CA-C-N	6.86	132.30	117.20
1	A	29	TYR	CA-CB-CG	-6.68	100.71	113.40
1	A	225	ARG	NE-CZ-NH2	-6.67	116.96	120.30
1	A	328	ARG	NE-CZ-NH1	6.56	123.58	120.30
1	A	72	ARG	NE-CZ-NH2	6.55	123.58	120.30
1	A	197	ARG	NE-CZ-NH1	6.54	123.57	120.30
1	A	319	ASP	CB-CG-OD2	-6.51	112.44	118.30
1	A	40	ARG	NE-CZ-NH1	-6.05	117.28	120.30
1	A	47	ARG	NH1-CZ-NH2	-6.03	112.76	119.40
1	A	319	ASP	C-N-CA	-5.93	106.86	121.70
1	A	341	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	A	261	ASP	CB-CG-OD2	-5.77	113.11	118.30
1	A	137	ARG	NE-CZ-NH1	-5.73	117.44	120.30
1	A	193	ILE	CA-C-N	5.73	129.80	117.20
1	A	307	ARG	CD-NE-CZ	5.71	131.59	123.60
1	A	307	ARG	CB-CA-C	5.68	121.76	110.40
1	A	88	ARG	NE-CZ-NH2	5.67	123.14	120.30
1	A	226	ARG	O-C-N	5.62	131.77	121.10
1	A	306	TRP	CA-CB-CG	-5.53	103.19	113.70
1	A	319	ASP	CB-CG-OD1	5.48	123.24	118.30
1	A	359	GLU	OE1-CD-OE2	-5.44	116.78	123.30
1	A	97	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	291	TRP	CA-CB-CG	5.28	123.73	113.70
1	A	77	ARG	NE-CZ-NH2	5.15	122.87	120.30
1	A	195	GLN	CG-CD-OE1	5.14	131.87	121.60
1	A	48	PRO	O-C-N	5.13	130.90	122.70
1	A	319	ASP	CA-C-O	-5.12	109.36	120.10
1	A	165	ARG	CB-CG-CD	5.02	124.66	111.60
1	A	59	ASP	CB-CG-OD1	5.02	122.82	118.30

There are no chirality outliers.

All (13) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	136	ILE	Mainchain
1	A	193	ILE	Mainchain
1	A	221	TYR	Mainchain

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	A	260	ILE	Mainchain
1	A	261	ASP	Peptide
1	A	306	TRP	Mainchain
1	A	317	ASP	Mainchain
1	A	319	ASP	Mainchain
1	A	327	VAL	Mainchain
1	A	328	ARG	Mainchain,Peptide
1	A	346	ASP	Mainchain
1	A	39	ASN	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2914	0	2801	63	0
2	A	6	0	4	12	0
3	A	43	0	0	0	0
All	All	2963	0	2805	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1001:BMD:CA	2:A:1001:BMD:NA	1.69	1.51
2:A:1001:BMD:CA	2:A:1001:BMD:C1	1.89	1.50
2:A:1001:BMD:C3	2:A:1001:BMD:C1	2.15	1.24
2:A:1001:BMD:C2	2:A:1001:BMD:C3	0.90	0.90
1:A:9:LEU:HD11	1:A:52:LEU:HD13	1.54	0.87
1:A:141:GLU:HB2	1:A:169:GLY:HA2	1.60	0.82
2:A:1001:BMD:C3	2:A:1001:BMD:CA	2.62	0.77
2:A:1001:BMD:C2	2:A:1001:BMD:C1	2.66	0.74
1:A:346:ASP:HB2	1:A:350:VAL:H	1.54	0.72
1:A:92:MET:HG3	1:A:117:ILE:HD11	1.71	0.71
1:A:337:ARG:HH21	1:A:362:ARG:HH11	1.43	0.66

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:LEU:HD11	1:A:67:ALA:HB2	1.79	0.64
1:A:134:TYR:O	1:A:138:HIS:HB2	2.00	0.62
1:A:136:ILE:HA	1:A:140:GLY:O	2.00	0.61
2:A:1001:BMD:OA	2:A:1001:BMD:NA	2.32	0.59
1:A:81:GLY:O	1:A:82:CYS:HB2	2.03	0.58
1:A:236:GLU:HG3	1:A:351:PHE:CE1	2.39	0.58
1:A:337:ARG:HG2	1:A:362:ARG:HA	1.87	0.57
1:A:139:TYR:CE1	1:A:228:PRO:HG3	2.40	0.57
1:A:213:LEU:O	1:A:217:ILE:HG13	2.05	0.56
1:A:92:MET:HG3	1:A:117:ILE:CD1	2.34	0.56
1:A:346:ASP:HB2	1:A:350:VAL:N	2.21	0.56
1:A:84:MET:HA	2:A:1001:BMD:OA	2.06	0.55
1:A:172:LEU:HD12	1:A:198:ALA:HB2	1.90	0.53
1:A:103:CYS:O	1:A:105:PRO:HD3	2.08	0.53
1:A:298:ARG:HD2	1:A:298:ARG:H	1.74	0.52
1:A:134:TYR:OH	1:A:354:ARG:HG3	2.10	0.51
1:A:320:ILE:HG22	1:A:321:ASP:N	2.26	0.51
1:A:298:ARG:N	1:A:298:ARG:HD2	2.25	0.51
1:A:59:ASP:HB3	1:A:62:ARG:HD3	1.93	0.50
1:A:312:GLN:O	1:A:313:ARG:C	2.49	0.49
1:A:184:ASP:HB3	1:A:188:ARG:NH1	2.28	0.49
1:A:218:ALA:CB	1:A:246:VAL:HG22	2.41	0.49
1:A:369:VAL:O	1:A:372:LEU:HB2	2.13	0.49
1:A:78:PHE:HD2	1:A:296:LEU:HD11	1.78	0.48
1:A:83:TYR:CE2	2:A:1001:BMD:C3	2.97	0.48
1:A:343:ALA:HA	1:A:353:VAL:HA	1.95	0.48
1:A:100:ALA:O	1:A:116:ASN:HB3	2.14	0.47
1:A:159:VAL:HG11	1:A:367:VAL:HG11	1.96	0.47
1:A:291:TRP:HZ3	1:A:295:LEU:HD22	1.80	0.47
1:A:14:PHE:CE1	1:A:81:GLY:HA2	2.50	0.47
1:A:10:ILE:O	1:A:51:THR:HA	2.14	0.47
1:A:86:HIS:H	1:A:86:HIS:CD2	2.32	0.46
1:A:236:GLU:HG3	1:A:351:PHE:HE1	1.79	0.46
1:A:152:TYR:HB3	1:A:153:PRO:HD3	1.98	0.46
1:A:187:GLN:NE2	1:A:187:GLN:HA	2.31	0.46
1:A:205:VAL:HG21	1:A:213:LEU:HD22	1.98	0.45
1:A:17:THR:O	1:A:57:GLY:HA2	2.17	0.45
1:A:299:ALA:HB3	1:A:311:VAL:HG13	1.99	0.45
1:A:129:ALA:HB3	1:A:130:PRO:CD	2.46	0.45
1:A:63:TYR:CD1	1:A:87:THR:HB	2.52	0.45
1:A:142:ARG:NH1	1:A:197:ARG:O	2.50	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:LEU:O	1:A:181:PRO:HD3	2.17	0.44
1:A:77:ARG:HG2	1:A:77:ARG:HH11	1.83	0.43
1:A:129:ALA:HB3	1:A:130:PRO:HD3	2.00	0.43
1:A:357:SER:HA	1:A:358:PRO:HD3	1.86	0.43
1:A:152:TYR:CG	2:A:1001:BMD:CA	3.02	0.43
1:A:184:ASP:HB3	1:A:188:ARG:HH12	1.84	0.43
1:A:10:ILE:HD13	1:A:10:ILE:HG21	1.87	0.42
1:A:266:ARG:O	1:A:270:GLN:HG2	2.19	0.42
2:A:1001:BMD:NA	2:A:1001:BMD:C1	2.80	0.42
1:A:163:LEU:HD13	1:A:369:VAL:HG22	2.00	0.42
1:A:9:LEU:HG	1:A:76:VAL:HG22	2.02	0.42
1:A:331:ARG:HB3	1:A:331:ARG:HE	1.68	0.41
1:A:119:TYR:HB2	1:A:335:HIS:HA	2.02	0.41
1:A:92:MET:HB3	1:A:93:PRO:HD3	2.03	0.41
1:A:250:GLN:O	1:A:345:ILE:HG13	2.20	0.41
1:A:12:LEU:HD13	1:A:28:ARG:HG3	2.01	0.41
1:A:186:LEU:HD11	1:A:212:GLU:HB3	2.02	0.41
1:A:199:ASP:O	1:A:228:PRO:HD2	2.21	0.41
2:A:1001:BMD:C2	2:A:1001:BMD:CA	2.99	0.41
1:A:166:GLN:OE1	1:A:166:GLN:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	366/385 (95%)	335 (92%)	26 (7%)	5 (1%)	14 35

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	138	HIS

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	261	ASP
1	A	329	VAL
1	A	225	ARG
1	A	41	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/315 (96%)	274 (90%)	29 (10%)	10	24

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	LEU
1	A	40	ARG
1	A	47	ARG
1	A	48	PRO
1	A	65	LEU
1	A	72	ARG
1	A	74	ARG
1	A	86	HIS
1	A	108	TYR
1	A	138	HIS
1	A	163	LEU
1	A	165	ARG
1	A	173	GLU
1	A	187	GLN
1	A	225	ARG
1	A	226	ARG
1	A	244	SER
1	A	261	ASP
1	A	265	SER
1	A	270	GLN
1	A	291	TRP
1	A	298	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	315	LEU
1	A	319	ASP
1	A	321	ASP
1	A	328	ARG
1	A	369	VAL
1	A	373	ASP
1	A	374	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	158	HIS
1	A	187	GLN
1	A	270	GLN
1	A	312	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BMD	A	1001	-	5,5,5	12.12	5 (100%)	5,5,5	11.52	5 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BMD	A	1001	-	-	0/3/3/3	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	BMD	C3-C2	-6.89	0.90	1.49
2	A	1001	BMD	OA-CA	6.43	1.44	1.24
2	A	1001	BMD	C1-CA	9.37	1.89	1.51
2	A	1001	BMD	CA-NA	11.44	1.69	1.32
2	A	1001	BMD	C2-C1	20.68	2.66	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	BMD	C3-C2-C1	-15.29	46.71	112.64
2	A	1001	BMD	C2-C1-CA	-12.96	80.08	112.80
2	A	1001	BMD	OA-CA-NA	-9.47	95.27	122.46
2	A	1001	BMD	C1-CA-NA	-4.38	102.76	116.53
2	A	1001	BMD	OA-CA-C1	12.35	157.03	121.05

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	BMD	12	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	338/385 (87%)	0.52	26 (7%) 16 14	10, 29, 51, 64	3 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	244	SER	5.3
1	A	303	ALA	4.2
1	A	374	ASP	4.0
1	A	246	VAL	3.5
1	A	136	ILE	3.5
1	A	245	ASP	3.5
1	A	270	GLN	3.4
1	A	248	GLU	3.2
1	A	317	ASP	2.8
1	A	243	GLU	2.8
1	A	279	ASN	2.6
1	A	306	TRP	2.6
1	A	168	GLY	2.5
1	A	141	GLU	2.5
1	A	232	LEU	2.4
1	A	102	LEU	2.4
1	A	278	GLU	2.4
1	A	219	ARG	2.4
1	A	345	ILE	2.2
1	A	310	ASP	2.2
1	A	105	PRO	2.2
1	A	358	PRO	2.1
1	A	187	GLN	2.1
1	A	75	GLY	2.1
1	A	197	ARG	2.1
1	A	354	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BMD	A	1001	6/6	0.95	0.27	1.37	3,4,12,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.