



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 07:03 PM GMT

PDB ID : 4QQQ
Title : Crystal structure of pneumolysin from *Streptococcus pneumoniae*, in complex with mannose as a component of cell membrane
Authors : Park, S.A.; Lee, K.S.
Deposited on : 2014-06-27
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	A	501	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	A	502	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3967 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pneumolysin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	483	3822	2408	656	751	7	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	HIS	-	EXPRESSION TAG	UNP Q2XU26
A	-11	HIS	-	EXPRESSION TAG	UNP Q2XU26
A	-10	HIS	-	EXPRESSION TAG	UNP Q2XU26
A	-9	SER	-	EXPRESSION TAG	UNP Q2XU26
A	-8	SER	-	EXPRESSION TAG	UNP Q2XU26
A	-7	GLY	-	EXPRESSION TAG	UNP Q2XU26
A	-6	LEU	-	EXPRESSION TAG	UNP Q2XU26
A	-5	VAL	-	EXPRESSION TAG	UNP Q2XU26
A	-4	PRO	-	EXPRESSION TAG	UNP Q2XU26
A	-3	ARG	-	EXPRESSION TAG	UNP Q2XU26
A	-2	GLY	-	EXPRESSION TAG	UNP Q2XU26
A	-1	SER	-	EXPRESSION TAG	UNP Q2XU26
A	0	HIS	-	EXPRESSION TAG	UNP Q2XU26

- Molecule 2 is SUGAR (ALPHA-D-MANNOSE) (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 12 6 6	0	0
2	A	1	Total C O 12 6 6	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	121	Total O 121 121	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pneumolysin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	24.68Å 157.45Å 211.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.82 – 2.50 40.82 – 2.49	Depositor EDS
% Data completeness (in resolution range)	88.7 (40.82-2.50) 88.4 (40.82-2.49)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.48Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.212 , 0.271 0.211 , 0.270	Depositor DCC
R_{free} test set	1334 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	33.0	Xtrriage
Anisotropy	0.223	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.5	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Outliers	0 of 26779 reflections	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3967	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/3900	0.65	1/5300 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	234	LEU	CB-CG-CD1	-5.40	101.81	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3822	0	3764	118	0
2	A	24	0	24	6	0
3	A	121	0	0	5	0
All	All	3967	0	3788	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (118) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:LYS:HA	1:A:50:LYS:HZ2	1.38	0.86
1:A:399:ARG:NH1	1:A:402:GLN:OE1	2.08	0.86
1:A:399:ARG:HH11	1:A:399:ARG:HG2	1.42	0.83
1:A:245:GLN:HB3	1:A:329:ILE:HD12	1.61	0.82
1:A:145:PRO:HB3	1:A:319:ARG:HH11	1.44	0.81
1:A:103:LEU:HD12	1:A:104:PRO:HD2	1.64	0.80
1:A:39:LEU:HB2	1:A:40:PRO:HD2	1.65	0.77
1:A:276:THR:OG1	1:A:277:GLU:N	2.15	0.76
1:A:433:TRP:NE1	3:A:703:HOH:O	2.18	0.76
1:A:403:ASP:OD2	2:A:502:MAN:H1	1.86	0.75
1:A:32:PHE:HZ	1:A:50:LYS:HD2	1.52	0.74
1:A:449:ARG:NH2	1:A:470:ASN:O	2.20	0.73
1:A:33:ILE:HD11	1:A:387:GLN:H	1.55	0.71
1:A:153:ILE:HG12	1:A:164:LYS:HG2	1.73	0.70
1:A:244:ARG:HG3	1:A:293:GLY:HA2	1.74	0.69
1:A:79:ASP:HB3	1:A:81:THR:H	1.59	0.68
1:A:259:VAL:HG13	1:A:281:ILE:HD11	1.79	0.63
1:A:20:LEU:HD13	1:A:78:VAL:HG21	1.81	0.62
1:A:196:LYS:HD3	1:A:329:ILE:HG22	1.82	0.62
1:A:247:TYR:HB2	1:A:290:VAL:HG13	1.80	0.62
1:A:324:HIS:O	1:A:324:HIS:ND1	2.35	0.60
1:A:148:MET:HE1	1:A:195:PHE:HD1	1.65	0.60
1:A:193:VAL:HG13	1:A:248:LEU:HB3	1.85	0.59
1:A:50:LYS:NZ	1:A:50:LYS:HA	2.15	0.58
1:A:386:HIS:O	1:A:387:GLN:HB2	2.04	0.58
1:A:378:THR:HG22	1:A:422:SER:HB2	1.84	0.58
1:A:256:SER:O	1:A:258:GLU:N	2.37	0.57
1:A:303:VAL:C	1:A:305:GLY:H	2.08	0.57
1:A:455:ILE:HG22	1:A:464:VAL:HB	1.86	0.56
1:A:26:GLU:OE1	1:A:348:THR:OG1	2.24	0.56
1:A:406:ALA:O	2:A:501:MAN:H3	2.07	0.55
1:A:286:GLU:OE2	1:A:306:LYS:NZ	2.30	0.55
1:A:51:ARG:NH1	3:A:698:HOH:O	2.31	0.55
1:A:88:THR:HG21	3:A:676:HOH:O	2.06	0.55
1:A:190:ILE:HG12	1:A:251:GLU:HG3	1.89	0.55
1:A:415:LYS:O	1:A:418:VAL:HG13	2.07	0.55
1:A:275:GLN:O	1:A:275:GLN:NE2	2.40	0.55
1:A:294:GLY:HA2	1:A:326:GLY:HA2	1.89	0.54
1:A:45:VAL:HG13	1:A:356:THR:HB	1.90	0.54
1:A:176:LEU:HD21	1:A:290:VAL:HG12	1.89	0.54
1:A:137:ASP:N	1:A:137:ASP:OD1	2.40	0.54
1:A:148:MET:HE1	1:A:195:PHE:CD1	2.43	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:LYS:HA	1:A:303:VAL:HA	1.90	0.53
1:A:51:ARG:NH1	1:A:352:GLU:OE1	2.38	0.53
1:A:435:TRP:HE1	2:A:502:MAN:H62	1.74	0.53
1:A:110:ASP:N	1:A:110:ASP:OD1	2.42	0.53
1:A:26:GLU:N	1:A:26:GLU:OE1	2.42	0.52
1:A:398:ASP:OD1	1:A:398:ASP:N	2.30	0.52
1:A:266:LEU:HD21	1:A:307:VAL:HB	1.92	0.52
1:A:234:LEU:HD11	1:A:336:LEU:HD22	1.92	0.51
1:A:440:TYR:OH	1:A:442:LYS:HD2	2.11	0.51
1:A:105:GLY:HA3	1:A:138:TYR:CZ	2.46	0.51
1:A:406:ALA:O	2:A:501:MAN:O1	2.22	0.51
1:A:98:THR:OG1	1:A:114:GLN:OE1	2.18	0.51
1:A:371:TYR:HB3	1:A:459:THR:HG22	1.93	0.51
1:A:10:ILE:HD13	1:A:336:LEU:HD23	1.92	0.51
1:A:295:ASP:HB3	1:A:296:PRO:HD2	1.92	0.51
1:A:222:ASP:OD1	3:A:658:HOH:O	2.20	0.50
1:A:256:SER:OG	1:A:256:SER:O	2.30	0.49
1:A:203:SER:HB3	1:A:239:SER:HB3	1.93	0.49
1:A:353:THR:O	1:A:353:THR:OG1	2.31	0.49
1:A:379:TRP:HE1	1:A:381:GLU:HG3	1.76	0.49
1:A:234:LEU:HD11	1:A:336:LEU:CD2	2.43	0.49
1:A:153:ILE:HG23	1:A:164:LYS:HE2	1.95	0.48
1:A:81:THR:OG1	1:A:88:THR:HG22	2.14	0.48
1:A:440:TYR:CZ	1:A:442:LYS:HD2	2.48	0.48
1:A:7:ASN:OD1	1:A:209:ASN:HB3	2.14	0.48
1:A:29:GLU:HG2	1:A:50:LYS:HD3	1.95	0.48
1:A:135:HIS:CE1	1:A:140:GLN:HG2	2.49	0.48
1:A:321:THR:OG1	1:A:323:ASP:OD1	2.33	0.47
1:A:43:PHE:CD1	1:A:417:ASN:HB3	2.49	0.47
1:A:142:ASN:OD1	1:A:142:ASN:N	2.46	0.47
1:A:80:GLU:HG2	1:A:81:THR:N	2.30	0.47
1:A:50:LYS:HZ2	1:A:51:ARG:H	1.63	0.47
1:A:11:LEU:HD13	1:A:210:PRO:HG2	1.96	0.47
1:A:103:LEU:HA	1:A:104:PRO:HD2	1.82	0.47
1:A:244:ARG:HD3	1:A:317:GLY:HA3	1.96	0.47
1:A:103:LEU:HG	1:A:134:TRP:CE2	2.49	0.46
1:A:237:ILE:HD11	1:A:240:VAL:CG2	2.45	0.46
1:A:311:GLU:O	1:A:315:GLN:HG2	2.15	0.46
1:A:393:THR:HA	1:A:394:PRO:HD3	1.82	0.46
1:A:196:LYS:NZ	1:A:245:GLN:OE1	2.48	0.46
1:A:39:LEU:HD11	1:A:42:GLU:OE2	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:VAL:HG13	1:A:242:TYR:HE1	1.81	0.46
1:A:338:ASP:OD1	1:A:340:VAL:HG13	2.16	0.46
1:A:29:GLU:HA	1:A:50:LYS:HD3	1.98	0.46
1:A:39:LEU:HD11	1:A:42:GLU:HG2	1.98	0.45
1:A:248:LEU:HD12	1:A:248:LEU:HA	1.73	0.45
1:A:107:ALA:HB3	1:A:110:ASP:OD2	2.16	0.45
1:A:435:TRP:NE1	2:A:502:MAN:O4	2.49	0.45
1:A:406:ALA:C	2:A:501:MAN:H3	2.36	0.45
1:A:29:GLU:HG3	1:A:349:ASP:OD2	2.16	0.45
1:A:303:VAL:C	1:A:305:GLY:N	2.70	0.45
1:A:78:VAL:HG22	1:A:233:PRO:HD2	1.99	0.44
1:A:17:LYS:HE3	1:A:340:VAL:HG21	2.00	0.44
1:A:20:LEU:HD13	1:A:78:VAL:CG2	2.46	0.44
1:A:256:SER:O	1:A:257:ASP:C	2.56	0.44
1:A:90:LEU:HD13	1:A:228:ILE:HD11	1.99	0.43
1:A:150:TYR:CE1	1:A:193:VAL:HG21	2.54	0.43
1:A:56:ASN:HA	1:A:344:PHE:O	2.19	0.43
1:A:32:PHE:HB2	1:A:48:ARG:HB2	2.00	0.43
1:A:105:GLY:HA3	1:A:138:TYR:CE2	2.54	0.43
1:A:19:LYS:HE3	1:A:19:LYS:HB2	1.60	0.43
1:A:450:LYS:HB3	1:A:469:GLU:HB2	2.02	0.42
1:A:360:ASN:HB3	3:A:706:HOH:O	2.19	0.42
1:A:27:SER:O	1:A:29:GLU:N	2.52	0.42
1:A:303:VAL:HG13	1:A:303:VAL:O	2.20	0.42
1:A:55:THR:HG22	1:A:346:ASN:HB2	2.02	0.42
1:A:306:LYS:HD3	1:A:306:LYS:HA	1.67	0.42
1:A:399:ARG:NH1	1:A:399:ARG:HG2	2.17	0.41
1:A:329:ILE:HD13	1:A:329:ILE:HA	1.88	0.41
1:A:209:ASN:HB2	1:A:212:ASP:OD1	2.20	0.40
1:A:202:VAL:HG13	1:A:242:TYR:CE1	2.55	0.40
1:A:75:LEU:HD23	1:A:236:TYR:HB3	2.03	0.40
1:A:361:GLY:HA3	1:A:448:VAL:O	2.22	0.40
1:A:106:LEU:HD12	1:A:111:SER:HB3	2.02	0.40
1:A:399:ARG:CG	1:A:399:ARG:HH11	2.24	0.40
1:A:234:LEU:HD12	1:A:234:LEU:C	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	481/484 (99%)	437 (91%)	30 (6%)	14 (3%)	6 8

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	28	ILE
1	A	145	PRO
1	A	257	ASP
1	A	25	GLY
1	A	107	ALA
1	A	138	TYR
1	A	256	SER
1	A	297	SER
1	A	30	ASN
1	A	304	THR
1	A	303	VAL
1	A	387	GLN
1	A	305	GLY
1	A	40	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	426/427 (100%)	367 (86%)	59 (14%)	4 8

All (59) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-11	HIS
1	A	1	MET
1	A	11	LEU
1	A	19	LYS
1	A	26	GLU
1	A	30	ASN
1	A	32	PHE
1	A	33	ILE
1	A	41	ASP
1	A	42	GLU
1	A	47	GLU
1	A	50	LYS
1	A	51	ARG
1	A	52	SER
1	A	55	THR
1	A	70	LEU
1	A	79	ASP
1	A	80	GLU
1	A	88	THR
1	A	106	LEU
1	A	110	ASP
1	A	137	ASP
1	A	138	TYR
1	A	142	ASN
1	A	143	ASN
1	A	153	ILE
1	A	160	GLN
1	A	175	SER
1	A	191	GLN
1	A	193	VAL
1	A	202	VAL
1	A	205	ASP
1	A	223	LEU
1	A	237	ILE
1	A	245	GLN
1	A	253	THR
1	A	254	SER
1	A	255	LYS
1	A	271	LYS
1	A	275	GLN
1	A	276	THR
1	A	290	VAL
1	A	304	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	307	VAL
1	A	308	ASP
1	A	311	GLU
1	A	324	HIS
1	A	327	LEU
1	A	330	SER
1	A	351	VAL
1	A	353	THR
1	A	367	HIS
1	A	378	THR
1	A	386	HIS
1	A	398	ASP
1	A	399	ARG
1	A	418	VAL
1	A	421	LEU
1	A	464	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	MAN	A	501	-	12,12,12	0.92	0	17,17,17	1.82	4 (23%)
2	MAN	A	502	-	12,12,12	0.76	0	17,17,17	1.95	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	A	501	-	-	0/2/22/22	0/1/1/1
2	MAN	A	502	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	MAN	C1-C2-C3	-4.43	103.42	110.68
2	A	502	MAN	C4-C3-C2	-3.94	103.53	110.79
2	A	502	MAN	C3-C4-C5	-3.37	104.22	110.23
2	A	502	MAN	O2-C2-C3	-2.96	103.69	110.36
2	A	502	MAN	C1-O5-C5	-2.38	109.00	113.54
2	A	501	MAN	C1-O5-C5	-2.17	109.39	113.54
2	A	501	MAN	C4-C3-C2	2.11	114.67	110.79
2	A	502	MAN	O5-C1-C2	2.37	114.15	110.00
2	A	502	MAN	O5-C5-C6	2.50	112.85	106.38
2	A	501	MAN	O2-C2-C3	3.60	118.47	110.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	MAN	3	0
2	A	502	MAN	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	483/484 (99%)	0.09	33 (6%) 20 23	11, 42, 80, 114	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	273	ALA	9.0
1	A	272	VAL	8.8
1	A	274	PRO	6.4
1	A	32	PHE	5.5
1	A	31	ARG	5.3
1	A	140	GLN	4.0
1	A	270	VAL	3.7
1	A	33	ILE	3.6
1	A	298	SER	3.6
1	A	29	GLU	3.6
1	A	28	ILE	3.5
1	A	30	ASN	3.5
1	A	276	THR	3.4
1	A	50	LYS	3.1
1	A	27	SER	3.1
1	A	304	THR	3.0
1	A	271	LYS	2.9
1	A	106	LEU	2.8
1	A	25	GLY	2.8
1	A	386	HIS	2.8
1	A	309	MET	2.7
1	A	141	VAL	2.7
1	A	105	GLY	2.5
1	A	301	ARG	2.4
1	A	387	GLN	2.4
1	A	139	GLY	2.3
1	A	297	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	388	GLY	2.2
1	A	136	GLN	2.2
1	A	315	GLN	2.2
1	A	24	GLN	2.1
1	A	308	ASP	2.1
1	A	18	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MAN	A	502	12/12	0.82	0.42	10.71	27,41,58,58	0
2	MAN	A	501	12/12	0.90	0.24	4.13	20,31,39,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.