



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2016 – 05:26 PM EDT

PDB ID : 4QRB
Title : Structure and specificity of L-D-Transpeptidase from Mycobacterium tuberculosis and antibiotic resistance: Calcium binding promotes dimer formation
Authors : Gokulan, K.; Varughese, K.I.
Deposited on : 2014-06-30
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1953 (1.66-1.62)
Clashscore	102246	2091 (1.66-1.62)
Ramachandran outliers	100387	2052 (1.66-1.62)
Sidechain outliers	100360	2052 (1.66-1.62)
RSRZ outliers	91569	1955 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MLD	A	501	-	-	-	X

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2812 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L,d-transpeptidase LdtB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	0	0	0
			2554	1601	447	498	8			

- Molecule 2 is GLCNAC(BETA1-4)-MURNAC(1,6-ANHYDRO)-L-ALA-GAMMA-D-GLU-MESO-A2PM-D-ALA (three-letter code: MLD) (formula: C₃₇H₅₉N₇O₂₀).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			49	28	7	14		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	209	Total	O	0	0
			209	209		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L,d-transpeptidase LdtB



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	57.33 Å 66.46 Å 206.87 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.52 – 1.64 42.48 – 1.64	Depositor EDS
% Data completeness (in resolution range)	98.9 (42.52-1.64) 98.9 (42.48-1.64)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.34 (at 1.64 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.214 , 0.236 0.219 , 0.245	Depositor DCC
R_{free} test set	2431 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.338	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2812	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MLD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	2/2615 (0.1%)	1.22	18/3578 (0.5%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	197	GLU	CD-OE2	-5.47	1.19	1.25
1	A	113	SER	CB-OG	-5.44	1.35	1.42

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ASP	CB-CG-OD1	10.89	128.10	118.30
1	A	402	ARG	NE-CZ-NH1	9.81	125.20	120.30
1	A	297	ARG	NE-CZ-NH1	8.51	124.56	120.30
1	A	237	MET	CG-SD-CE	-8.37	86.81	100.20
1	A	211	ARG	NE-CZ-NH2	-8.28	116.16	120.30
1	A	370	LYS	CD-CE-NZ	8.00	130.09	111.70
1	A	160	ASP	CB-CG-OD1	7.79	125.31	118.30
1	A	68	ASP	CB-CG-OD1	7.74	125.26	118.30
1	A	232	ASP	CB-CG-OD2	-7.51	111.54	118.30
1	A	267	ARG	NE-CZ-NH1	7.13	123.86	120.30
1	A	96	ASP	CB-CG-OD2	-6.86	112.13	118.30
1	A	297	ARG	NE-CZ-NH2	-6.62	116.99	120.30
1	A	123	ARG	NE-CZ-NH2	-6.29	117.16	120.30
1	A	206	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	A	211	ARG	NE-CZ-NH1	5.67	123.14	120.30
1	A	141	LEU	CB-CG-CD1	5.23	119.89	111.00
1	A	393	ASP	CB-CG-OD1	5.14	122.93	118.30
1	A	200	PHE	CB-CG-CD2	5.08	124.36	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2554	0	2459	22	0
2	A	49	0	38	13	0
3	A	209	0	0	2	0
All	All	2812	0	2497	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:ARG:HD2	2:A:501:MLD:HBG	1.39	1.04
2:A:501:MLD:HA1	2:A:501:MLD:CAF	1.96	0.94
2:A:501:MLD:HBP2	2:A:501:MLD:CBT	1.99	0.92
2:A:501:MLD:HA1	2:A:501:MLD:HAF3	1.50	0.91
1:A:127:ASN:HD21	1:A:140:GLN:HE21	1.23	0.87
2:A:501:MLD:C2	2:A:501:MLD:HA61	2.06	0.85
1:A:291:ILE:H	1:A:395:ASN:HD21	1.27	0.81
1:A:139:ARG:CD	2:A:501:MLD:HBG	2.11	0.80
2:A:501:MLD:HBP2	2:A:501:MLD:OBU	1.87	0.73
2:A:501:MLD:N2	2:A:501:MLD:HA61	2.04	0.71
1:A:177:ASN:HD22	1:A:205:ASN:HB3	1.60	0.67
1:A:223:ASP:OD1	1:A:246:HIS:HD2	1.79	0.65
1:A:289:ASN:HD21	1:A:380:THR:H	1.48	0.61
1:A:246:HIS:HE1	3:A:633:HOH:O	1.84	0.60
1:A:177:ASN:ND2	1:A:205:ASN:HB3	2.17	0.59
1:A:302:ILE:O	1:A:302:ILE:HG23	2.02	0.58
1:A:120:TYR:O	1:A:121:ASN:HB2	2.04	0.57
2:A:501:MLD:CA1	2:A:501:MLD:HAF3	2.33	0.56
1:A:296:SER:OG	1:A:371:ARG:NH2	2.38	0.55
2:A:501:MLD:N2	2:A:501:MLD:CA6	2.73	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:SER:OG	1:A:350:THR:HG22	2.09	0.52
1:A:243:VAL:HG12	1:A:244:GLN:N	2.26	0.49
2:A:501:MLD:HAF2	2:A:501:MLD:HA1	1.90	0.49
2:A:501:MLD:C2	2:A:501:MLD:CA6	2.85	0.48
1:A:127:ASN:ND2	1:A:140:GLN:HE21	2.02	0.47
1:A:297:ARG:HB3	1:A:297:ARG:CZ	2.43	0.46
1:A:177:ASN:HD22	1:A:205:ASN:CB	2.27	0.45
1:A:265:THR:HG23	3:A:714:HOH:O	2.20	0.42
1:A:289:ASN:HD21	1:A:380:THR:N	2.15	0.42
1:A:301:ILE:CG1	1:A:302:ILE:N	2.84	0.41
1:A:64:ALA:O	2:A:501:MLD:HAI	2.20	0.41
1:A:224:VAL:HB	1:A:245:THR:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	335/352 (95%)	321 (96%)	13 (4%)	1 (0%)	46 23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	303	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	271/282 (96%)	262 (97%)	9 (3%)	45 14

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	LYS
1	A	131	LEU
1	A	133	LEU
1	A	177	ASN
1	A	284	SER
1	A	297	ARG
1	A	303	MET
1	A	350	THR
1	A	393	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	127	ASN
1	A	177	ASN
1	A	246	HIS
1	A	289	ASN
1	A	327	GLN
1	A	361	ASN
1	A	363	GLN
1	A	395	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MLD	A	501	-	44,50,66	1.76	11 (25%)	52,69,93	3.55	27 (51%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLD	A	501	-	-	0/46/82/114	0/0/2/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	MLD	CAF-CAE	-3.84	1.42	1.50
2	A	501	MLD	OBI-CBH	-2.67	1.18	1.23
2	A	501	MLD	OAM-CAL	-2.62	1.18	1.23
2	A	501	MLD	OA5-CA5	-2.13	1.40	1.45
2	A	501	MLD	CA2-NAD	2.10	1.49	1.45
2	A	501	MLD	O1-C1	2.32	1.48	1.41
2	A	501	MLD	OA5-CA1	2.41	1.46	1.41
2	A	501	MLD	CA3-CA4	3.18	1.59	1.52
2	A	501	MLD	CBH-NBJ	3.23	1.39	1.32
2	A	501	MLD	CA3-CA2	3.82	1.59	1.53
2	A	501	MLD	CA1-CA2	4.09	1.60	1.53

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	MLD	OA5-CA1-OA6	-12.64	95.55	105.88
2	A	501	MLD	OA5-CA5-CA6	-7.00	94.25	101.51
2	A	501	MLD	CBP-CBG-NBF	-6.55	98.45	110.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	MLD	OBI-CBH-CBG	-6.04	113.23	120.24
2	A	501	MLD	CBB-CBC-CBD	-5.12	101.55	113.26
2	A	501	MLD	CA4-CA3-CA2	-5.00	100.52	110.26
2	A	501	MLD	OAG-CAE-CAF	-4.72	113.38	122.07
2	A	501	MLD	OBE-CBD-NBF	-4.27	115.78	122.96
2	A	501	MLD	OAM-CAL-CAI	-2.85	113.35	119.54
2	A	501	MLD	CB-CA-C	-2.24	105.71	110.20
2	A	501	MLD	OAM-CAL-N	-2.22	118.59	122.91
2	A	501	MLD	CAJ-CAI-CAL	-2.02	107.21	110.91
2	A	501	MLD	CA6-OA6-CA1	2.07	109.70	106.81
2	A	501	MLD	CB-CA-N	2.28	114.62	110.27
2	A	501	MLD	CBC-CBB-CAO	2.41	117.64	113.05
2	A	501	MLD	CAI-CAL-N	2.98	127.97	117.85
2	A	501	MLD	CA2-NAD-CAE	3.04	131.20	123.21
2	A	501	MLD	OBE-CBD-CBC	3.20	127.52	121.97
2	A	501	MLD	OAH-CAI-CAJ	3.29	117.01	107.55
2	A	501	MLD	OAG-CAE-NAD	3.31	128.59	121.84
2	A	501	MLD	OA5-CA5-CA4	3.66	114.30	108.66
2	A	501	MLD	CBG-CBH-NBJ	3.71	123.54	116.93
2	A	501	MLD	OAH-CA3-CA4	3.93	117.74	107.48
2	A	501	MLD	CAP-CAO-NAN	4.16	121.24	112.93
2	A	501	MLD	OAH-CAI-CAL	4.42	121.38	111.01
2	A	501	MLD	CA3-CA2-NAD	5.73	120.87	111.06
2	A	501	MLD	O1-CA4-CA3	6.83	125.30	107.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	MLD	13	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	339/352 (96%)	0.17	13 (3%) 44 41	13, 24, 44, 61	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	58	LEU	5.8
1	A	302	ILE	5.7
1	A	320	THR	5.7
1	A	57	LEU	5.3
1	A	133	LEU	5.2
1	A	340	TRP	3.4
1	A	303	MET	2.8
1	A	338	ALA	2.8
1	A	359	PRO	2.7
1	A	304	ASP	2.4
1	A	342	VAL	2.3
1	A	265	THR	2.3
1	A	301	ILE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MLD	A	501	49/64	0.54	0.25	16.04	39,61,85,96	0

6.5 Other polymers [i](#)

There are no such residues in this entry.