



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:10 PM GMT

PDB ID : 4QVW  
Title : yCP beta5-A49S-mutant in complex with bortezomib  
Authors : Huber, E.M.; Heinemeyer, W.; Groll, M.  
Deposited on : 2014-07-16  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	<div> <div>98%</div> <div> <div></div> <div>98%</div> <div>5%</div> <div>5%</div> </div> </div>
1	O	250	<div> <div>98%</div> <div> <div></div> <div>98%</div> <div>5%</div> <div>5%</div> </div> </div>
2	B	258	<div> <div>89%</div> <div> <div></div> <div>89%</div> <div>6%</div> <div>5%</div> </div> </div>
2	P	258	<div> <div>88%</div> <div> <div></div> <div>88%</div> <div>6%</div> <div>5%</div> </div> </div>
3	C	254	<div> <div>87%</div> <div> <div></div> <div>87%</div> <div>6%</div> <div>6%</div> </div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	254	
4	D	260	
4	R	260	
5	E	234	
5	S	234	
6	F	288	
6	T	288	
7	G	252	
7	U	252	
8	H	232	
8	V	232	
9	I	205	
9	W	205	
10	J	198	
10	X	198	
11	K	212	
11	Y	212	
12	L	222	
12	Z	222	
13	M	246	
13	a	246	
14	N	196	
14	b	196	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
15	MG	I	301	-	-	-	X
15	MG	J	201	-	-	-	X
15	MG	Z	301	-	-	-	X
17	BO2	N	201	-	-	-	X
17	BO2	V	301	-	-	-	X
17	BO2	Y	301	-	-	-	X
17	BO2	b	201	-	-	-	X

## 2 Entry composition

There are 18 unique types of molecules in this entry. The entry contains 49726 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			
2	P	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			

- Molecule 3 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			
3	Q	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			

- Molecule 4 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			
4	R	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			

- Molecule 5 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			
5	S	231	Total	C	N	O	S	0	0	0
			1773	1114	307	348	4			

- Molecule 6 is a protein called Probable proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			
6	T	243	Total	C	N	O	S	0	0	0
			1892	1203	329	356	4			

- Molecule 7 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			
7	U	241	Total	C	N	O	S	0	0	0
			1907	1214	320	365	8			

- Molecule 8 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	226	Total	C	N	O	S	0	0	0
			1719	1082	298	332	7			
8	V	226	Total	C	N	O	S	0	0	0
			1719	1082	298	332	7			

- Molecule 9 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

- Molecule 11 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1645	1045	280	313	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1645	1045	280	313	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
K	49	SER	ALA	ENGINEERED MUTATION	UNP P30656
Y	49	SER	ALA	ENGINEERED MUTATION	UNP P30656

- Molecule 12 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	a	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	b	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
15	G	1	Total Mg 1 1	0	0
15	J	1	Total Mg 1 1	0	0
15	K	1	Total Mg 1 1	0	0
15	I	1	Total Mg 1 1	0	0
15	V	1	Total Mg 1 1	0	0
15	Z	1	Total Mg 1 1	0	0
15	N	1	Total Mg 1 1	0	0
15	Y	1	Total Mg 1 1	0	0

- Molecule 16 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	G	1	Total Cl 1 1	0	0
16	b	1	Total Cl 1 1	0	0
16	N	1	Total Cl 1 1	0	0
16	U	1	Total Cl 1 1	0	0

- Molecule 17 is N-[(1R)-1-(DIHYDROXYBORYL)-3-METHYLBUTYL]-N-(PYRAZIN-2-YLCARBONYL)-L-PHENYLALANINAMIDE (three-letter code: BO2) (formula: C<sub>19</sub>H<sub>25</sub>BN<sub>4</sub>O<sub>4</sub>).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
17	H	1	Total	B	C	N	O	0	0
			28	1	19	4	4		
17	K	1	Total	B	C	N	O	0	0
			28	1	19	4	4		
17	N	1	Total	B	C	N	O	0	0
			28	1	19	4	4		
17	V	1	Total	B	C	N	O	0	0
			28	1	19	4	4		
17	Y	1	Total	B	C	N	O	0	0
			28	1	19	4	4		
17	b	1	Total	B	C	N	O	0	0
			28	1	19	4	4		

- Molecule 18 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
18	A	6	Total	O	0	0
			6	6		
18	B	7	Total	O	0	0
			7	7		
18	C	5	Total	O	0	0
			5	5		
18	D	3	Total	O	0	0
			3	3		
18	E	3	Total	O	0	0
			3	3		
18	F	5	Total	O	0	0
			5	5		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
18	G	6	Total 6	O 6	0	0
18	H	7	Total 7	O 7	0	0
18	I	3	Total 3	O 3	0	0
18	J	6	Total 6	O 6	0	0
18	K	7	Total 7	O 7	0	0
18	L	11	Total 11	O 11	0	0
18	M	9	Total 9	O 9	0	0
18	N	4	Total 4	O 4	0	0
18	O	3	Total 3	O 3	0	0
18	P	3	Total 3	O 3	0	0
18	Q	9	Total 9	O 9	0	0
18	R	4	Total 4	O 4	0	0
18	S	1	Total 1	O 1	0	0
18	T	5	Total 5	O 5	0	0
18	U	8	Total 8	O 8	0	0
18	V	8	Total 8	O 8	0	0
18	W	6	Total 6	O 6	0	0
18	X	12	Total 12	O 12	0	0
18	Y	10	Total 10	O 10	0	0
18	Z	8	Total 8	O 8	0	0
18	a	11	Total 11	O 11	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
18	b	8	Total	O	0	0
			8	8		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit alpha type-2




- Molecule 1: Proteasome subunit alpha type-2




- Molecule 2: Proteasome subunit alpha type-3




- Molecule 2: Proteasome subunit alpha type-3




- Molecule 3: Proteasome subunit alpha type-4




- Molecule 3: Proteasome subunit alpha type-4




- Molecule 4: Proteasome subunit alpha type-5




- Molecule 4: Proteasome subunit alpha type-5




- Molecule 5: Proteasome subunit alpha type-6




- Molecule 5: Proteasome subunit alpha type-6



- Molecule 6: Probable proteasome subunit alpha type-7




- Molecule 6: Probable proteasome subunit alpha type-7




GLN  
GLU  
GLY  
ASP  
ILE  
HIS  
LEU  
GLU

• Molecule 7: Proteasome subunit alpha type-1

Chain G:  90% 6% •


MET SER GLY ALA ALA ALA SER ALA ALA G2 F23 T26 V73 I78 P79 L115 R122 M125 T133 D149 P150 E208 R235 L236 Q242 ASP

• Molecule 7: Proteasome subunit alpha type-1

Chain U:  89% 6% •


MET SER GLY ALA ALA ALA SER ALA ALA G2 P12 F23 T26 V73 I78 P79 N83 L115 R122 M125 T133 D149 P150 E208 D222 R235 L236 Q242 ASP

• Molecule 8: Proteasome subunit beta type-2

Chain H:  92% 5% • •

T1 N30 R35 T36 L68 D104 P105 I113 H114 T119 S126 L127 R196 D222 E225 GLN VAL ASP ILE THR ALA

• Molecule 8: Proteasome subunit beta type-2

Chain V:  91% 6% • •

T1 N30 R35 T36 L68 K34 Q35 D104 P105 I113 T119 S126 L127 E139 R196 C221 D222 I223 Q224 E225 E226 GLN VAL ASP ILE THR ALA

• Molecule 9: Proteasome subunit beta type-3

Chain I:  94% 5%


MET S1 G9 I10 V20 S36 N37 K38 K41 P118 A141 E150 L171 D204


• Molecule 9: Proteasome subunit beta type-3

Chain W:  94% 5%


MET S1 G9 I10 V20 S36 N37 K38 K41 P118 A141 E150 L171 D204

• Molecule 10: Proteasome subunit beta type-4



Chain J:  90% 7% • •



- Molecule 10: Proteasome subunit beta type-4



- Molecule 11: Proteasome subunit beta type-5



- Molecule 11: Proteasome subunit beta type-5



- Molecule 12: Proteasome subunit beta type-6



- Molecule 12: Proteasome subunit beta type-6




- Molecule 13: Proteasome subunit beta type-7




- Molecule 13: Proteasome subunit beta type-7






● Molecule 14: Proteasome subunit beta type-1



● Molecule 14: Proteasome subunit beta type-1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.75Å 300.43Å 145.37Å 90.00° 112.99° 90.00°	Depositor
Resolution (Å)	15.00 – 3.00 15.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	98.6 (15.00-3.00) 98.7 (15.00-3.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.84 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.176 , 0.209 0.180 , 0.210	Depositor DCC
$R_{free}$ test set	10435 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	64.3	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 43.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 208712 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	49726	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BO2, MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/1952	0.46	0/2642
1	O	0.27	0/1952	0.46	0/2642
2	B	0.27	0/1934	0.49	0/2618
2	P	0.27	0/1934	0.49	0/2618
3	C	0.27	0/1910	0.50	0/2586
3	Q	0.27	0/1910	0.50	0/2586
4	D	0.26	0/1837	0.47	0/2475
4	R	0.26	0/1837	0.47	0/2475
5	E	0.27	0/1800	0.47	0/2433
5	S	0.27	0/1800	0.47	0/2433
6	F	0.27	0/1932	0.45	0/2609
6	T	0.27	0/1932	0.45	0/2609
7	G	0.27	0/1945	0.46	0/2634
7	U	0.27	0/1945	0.46	0/2634
8	H	0.25	0/1750	0.46	0/2373
8	V	0.25	0/1750	0.46	0/2373
9	I	0.27	0/1611	0.48	0/2174
9	W	0.27	0/1611	0.48	0/2174
10	J	0.43	1/1589 (0.1%)	0.53	1/2142 (0.0%)
10	X	0.40	0/1589	0.53	1/2142 (0.0%)
11	K	0.26	0/1682	0.51	0/2275
11	Y	0.26	0/1682	0.51	0/2275
12	L	0.27	0/1795	0.48	0/2420
12	Z	0.27	0/1795	0.47	0/2420
13	M	0.27	0/1855	0.50	0/2514
13	a	0.27	0/1855	0.50	0/2514
14	N	0.25	0/1541	0.47	0/2087
14	b	0.25	0/1541	0.47	0/2087
All	All	0.28	1/50266 (0.0%)	0.48	2/67964 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	J	150	PRO	N-CD	5.01	1.54	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	X	149	ARG	C-N-CD	5.87	140.72	128.40
10	J	149	ARG	C-N-CD	5.68	140.33	128.40

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1929	0	0
1	O	1915	0	1929	2	0
2	B	1904	0	1904	7	0
2	P	1904	0	1904	6	0
3	C	1881	0	1895	5	0
3	Q	1881	0	1895	5	0
4	D	1813	0	1797	2	0
4	R	1813	0	1797	2	0
5	E	1773	0	1775	2	0
5	S	1773	0	1775	3	0
6	F	1892	0	1883	2	0
6	T	1892	0	1883	2	0
7	G	1907	0	1901	4	0
7	U	1907	0	1901	5	0
8	H	1719	0	1718	12	0
8	V	1719	0	1718	11	0
9	I	1581	0	1574	6	0
9	W	1581	0	1574	6	0
10	J	1561	0	1569	11	0
10	X	1561	0	1569	21	0
11	K	1645	0	1594	16	0
11	Y	1645	0	1594	13	0
12	L	1757	0	1711	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
12	Z	1757	0	1711	2	0
13	M	1824	0	1832	3	0
13	a	1824	0	1832	0	0
14	N	1512	0	1480	3	0
14	b	1512	0	1480	0	0
15	G	1	0	0	0	0
15	I	1	0	0	0	0
15	J	1	0	0	0	0
15	K	1	0	0	0	0
15	N	1	0	0	0	0
15	V	1	0	0	0	0
15	Y	1	0	0	0	0
15	Z	1	0	0	0	0
16	G	1	0	0	0	0
16	N	1	0	0	0	0
16	U	1	0	0	0	0
16	b	1	0	0	0	0
17	H	28	0	25	1	0
17	K	28	0	25	1	0
17	N	28	0	25	4	0
17	V	28	0	25	1	0
17	Y	28	0	25	1	0
17	b	28	0	25	0	0
18	A	6	0	0	0	0
18	B	7	0	0	1	0
18	C	5	0	0	0	0
18	D	3	0	0	0	0
18	E	3	0	0	0	0
18	F	5	0	0	0	0
18	G	6	0	0	0	0
18	H	7	0	0	0	0
18	I	3	0	0	0	0
18	J	6	0	0	0	0
18	K	7	0	0	0	0
18	L	11	0	0	0	0
18	M	9	0	0	0	0
18	N	4	0	0	0	0
18	O	3	0	0	0	0
18	P	3	0	0	0	0
18	Q	9	0	0	0	0
18	R	4	0	0	0	0
18	S	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
18	T	5	0	0	0	0
18	U	8	0	0	0	0
18	V	8	0	0	0	0
18	W	6	0	0	0	0
18	X	12	0	0	0	0
18	Y	10	0	0	0	0
18	Z	8	0	0	0	0
18	a	11	0	0	0	0
18	b	8	0	0	0	0
All	All	49726	0	49274	135	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 135 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:X:152:MET:HE3	10:X:156:GLU:HB3	1.54	0.89
10:J:149:ARG:O	10:J:152:MET:HG3	1.74	0.87
11:Y:104:TYR:CZ	11:Y:182:GLU:HG3	2.24	0.72
11:K:104:TYR:CZ	11:K:182:GLU:HG3	2.27	0.69
8:H:114:HIS:CD2	17:N:201:BO2:H5	2.29	0.68

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	248/250 (99%)	239 (96%)	8 (3%)	1 (0%)	39 80
1	O	248/250 (99%)	239 (96%)	8 (3%)	1 (0%)	39 80

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	242/258 (94%)	233 (96%)	5 (2%)	4 (2%)	11	46
2	P	242/258 (94%)	234 (97%)	4 (2%)	4 (2%)	11	46
3	C	238/254 (94%)	233 (98%)	2 (1%)	3 (1%)	15	53
3	Q	238/254 (94%)	233 (98%)	2 (1%)	3 (1%)	15	53
4	D	231/260 (89%)	228 (99%)	3 (1%)	0	100	100
4	R	231/260 (89%)	228 (99%)	3 (1%)	0	100	100
5	E	229/234 (98%)	221 (96%)	8 (4%)	0	100	100
5	S	229/234 (98%)	221 (96%)	8 (4%)	0	100	100
6	F	241/288 (84%)	238 (99%)	3 (1%)	0	100	100
6	T	241/288 (84%)	238 (99%)	3 (1%)	0	100	100
7	G	239/252 (95%)	237 (99%)	2 (1%)	0	100	100
7	U	239/252 (95%)	237 (99%)	2 (1%)	0	100	100
8	H	224/232 (97%)	219 (98%)	5 (2%)	0	100	100
8	V	224/232 (97%)	219 (98%)	5 (2%)	0	100	100
9	I	202/205 (98%)	194 (96%)	8 (4%)	0	100	100
9	W	202/205 (98%)	194 (96%)	8 (4%)	0	100	100
10	J	193/198 (98%)	188 (97%)	4 (2%)	1 (0%)	34	76
10	X	193/198 (98%)	189 (98%)	4 (2%)	0	100	100
11	K	210/212 (99%)	206 (98%)	4 (2%)	0	100	100
11	Y	210/212 (99%)	206 (98%)	4 (2%)	0	100	100
12	L	220/222 (99%)	215 (98%)	5 (2%)	0	100	100
12	Z	220/222 (99%)	215 (98%)	5 (2%)	0	100	100
13	M	231/246 (94%)	224 (97%)	7 (3%)	0	100	100
13	a	231/246 (94%)	224 (97%)	7 (3%)	0	100	100
14	N	194/196 (99%)	189 (97%)	5 (3%)	0	100	100
14	b	194/196 (99%)	189 (97%)	5 (3%)	0	100	100
All	All	6284/6614 (95%)	6130 (98%)	137 (2%)	17 (0%)	46	84

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	51	VAL
2	B	222	GLY

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
3	C	202	GLN
2	P	51	VAL
2	P	222	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	206 (99%)	3 (1%)	74	93
1	O	209/209 (100%)	206 (99%)	3 (1%)	74	93
2	B	203/216 (94%)	200 (98%)	3 (2%)	72	92
2	P	203/216 (94%)	199 (98%)	4 (2%)	63	89
3	C	212/226 (94%)	202 (95%)	10 (5%)	32	72
3	Q	212/226 (94%)	202 (95%)	10 (5%)	32	72
4	D	194/215 (90%)	186 (96%)	8 (4%)	37	76
4	R	194/215 (90%)	186 (96%)	8 (4%)	37	76
5	E	190/193 (98%)	183 (96%)	7 (4%)	41	79
5	S	190/193 (98%)	183 (96%)	7 (4%)	41	79
6	F	201/239 (84%)	192 (96%)	9 (4%)	34	74
6	T	201/239 (84%)	192 (96%)	9 (4%)	34	74
7	G	206/210 (98%)	200 (97%)	6 (3%)	50	84
7	U	206/210 (98%)	199 (97%)	7 (3%)	44	81
8	H	185/190 (97%)	181 (98%)	4 (2%)	60	88
8	V	185/190 (97%)	181 (98%)	4 (2%)	60	88
9	I	172/173 (99%)	170 (99%)	2 (1%)	78	94
9	W	172/173 (99%)	170 (99%)	2 (1%)	78	94
10	J	173/175 (99%)	168 (97%)	5 (3%)	50	84
10	X	173/175 (99%)	167 (96%)	6 (4%)	43	80
11	K	170/170 (100%)	162 (95%)	8 (5%)	32	72

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
11	Y	170/170 (100%)	162 (95%)	8 (5%)	32	72
12	L	185/185 (100%)	182 (98%)	3 (2%)	70	92
12	Z	185/185 (100%)	181 (98%)	4 (2%)	60	88
13	M	199/208 (96%)	193 (97%)	6 (3%)	48	83
13	a	199/208 (96%)	193 (97%)	6 (3%)	48	83
14	N	162/162 (100%)	157 (97%)	5 (3%)	47	83
14	b	162/162 (100%)	157 (97%)	5 (3%)	47	83
All	All	5322/5542 (96%)	5160 (97%)	162 (3%)	48	83

5 of 162 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
13	M	187	ARG
3	Q	147	GLN
12	Z	136	CYS
14	N	39	ASP
2	P	55	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 88 such sidechains are listed below:

Mol	Chain	Res	Type
12	L	3	ASN
2	P	95	GLN
12	Z	3	ASN
12	L	79	HIS
13	M	194	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
17	BO2	H	301	8	28,29,29	1.55	5 (17%)	31,38,38	1.30	6 (19%)
17	BO2	K	301	11	28,29,29	1.53	5 (17%)	31,38,38	1.25	5 (16%)
17	BO2	N	201	14	28,29,29	1.51	4 (14%)	31,38,38	1.11	3 (9%)
17	BO2	V	301	8	28,29,29	1.54	5 (17%)	31,38,38	1.28	6 (19%)
17	BO2	Y	301	11	28,29,29	1.54	5 (17%)	31,38,38	1.25	4 (12%)
17	BO2	b	201	14	28,29,29	1.50	4 (14%)	31,38,38	1.11	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
17	BO2	H	301	8	-	0/22/28/28	0/2/2/2
17	BO2	K	301	11	-	0/22/28/28	0/2/2/2
17	BO2	N	201	14	-	0/22/28/28	0/2/2/2
17	BO2	V	301	8	-	0/22/28/28	0/2/2/2
17	BO2	Y	301	11	-	0/22/28/28	0/2/2/2
17	BO2	b	201	14	-	0/22/28/28	0/2/2/2

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
17	N	201	BO2	C2-C7	-5.24	1.38	1.50
17	b	201	BO2	C2-C7	-5.22	1.38	1.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
17	H	301	BO2	C2-C7	-4.77	1.39	1.50
17	Y	301	BO2	C2-C7	-4.72	1.39	1.50
17	V	301	BO2	C2-C7	-4.70	1.39	1.50

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
17	b	201	BO2	C11-C10-N9	-2.91	104.70	110.80
17	N	201	BO2	C11-C10-N9	-2.72	105.11	110.80
17	H	301	BO2	B26-C21-C22	-2.53	106.23	112.79
17	V	301	BO2	B26-C21-C22	-2.50	106.31	112.79
17	Y	301	BO2	C18-C10-N9	-2.49	104.23	111.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
17	H	301	BO2	1	0
17	K	301	BO2	1	0
17	N	201	BO2	4	0
17	V	301	BO2	1	0
17	Y	301	BO2	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/250 (100%)	-0.61	2 (0%) 87 67	47, 61, 98, 130	0
1	O	250/250 (100%)	-0.60	2 (0%) 87 67	49, 67, 109, 143	0
2	B	244/258 (94%)	-0.59	3 (1%) 81 55	46, 66, 110, 160	0
2	P	244/258 (94%)	-0.53	5 (2%) 68 39	51, 70, 110, 159	0
3	C	240/254 (94%)	-0.45	6 (2%) 61 30	46, 70, 131, 156	0
3	Q	240/254 (94%)	-0.33	8 (3%) 50 22	53, 83, 160, 181	0
4	D	235/260 (90%)	-0.62	0 100 100	50, 72, 103, 144	0
4	R	235/260 (90%)	-0.50	3 (1%) 79 53	57, 78, 119, 148	0
5	E	231/234 (98%)	-0.50	2 (0%) 85 64	53, 76, 116, 159	0
5	S	231/234 (98%)	-0.48	1 (0%) 93 80	54, 81, 124, 162	0
6	F	243/288 (84%)	-0.61	1 (0%) 93 80	49, 70, 119, 142	0
6	T	243/288 (84%)	-0.58	1 (0%) 93 80	46, 76, 135, 162	0
7	G	241/252 (95%)	-0.66	0 100 100	45, 64, 103, 153	0
7	U	241/252 (95%)	-0.61	2 (0%) 87 67	48, 64, 98, 145	0
8	H	226/232 (97%)	-0.64	2 (0%) 85 64	41, 56, 94, 148	0
8	V	226/232 (97%)	-0.63	5 (2%) 65 35	43, 60, 94, 170	0
9	I	204/205 (99%)	-0.82	0 100 100	40, 56, 86, 110	0
9	W	204/205 (99%)	-0.80	1 (0%) 91 76	43, 57, 85, 114	0
10	J	195/198 (98%)	-0.70	1 (0%) 91 76	43, 59, 88, 129	0
10	X	195/198 (98%)	-0.68	2 (1%) 84 60	46, 62, 93, 145	0
11	K	212/212 (100%)	-0.59	5 (2%) 62 32	41, 61, 100, 114	0
11	Y	212/212 (100%)	-0.64	1 (0%) 91 76	42, 59, 97, 114	0
12	L	222/222 (100%)	-0.65	3 (1%) 78 51	45, 61, 110, 142	0
12	Z	222/222 (100%)	-0.65	4 (1%) 71 43	33, 60, 110, 142	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
13	M	233/246 (94%)	-0.75	1 (0%) 93 80	43, 60, 80, 99	0
13	a	233/246 (94%)	-0.77	1 (0%) 93 80	39, 58, 80, 97	0
14	N	196/196 (100%)	-0.81	1 (0%) 91 76	42, 53, 83, 109	0
14	b	196/196 (100%)	-0.76	1 (0%) 91 76	42, 55, 86, 116	0
All	All	6344/6614 (95%)	-0.62	64 (1%) 84 60	33, 65, 112, 181	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	Q	49	THR	5.5
11	Y	212	GLY	4.6
12	Z	165	ASN	4.1
3	Q	50	LEU	4.1
2	P	219	ALA	4.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
15	MG	I	301	1/1	0.96	0.34	7.56	67,67,67,67	0
17	BO2	N	201	28/28	0.90	0.25	4.07	40,54,66,68	0
15	MG	J	201	1/1	0.96	0.20	3.78	38,38,38,38	0
15	MG	Z	301	1/1	0.96	0.28	3.62	62,62,62,62	0
17	BO2	b	201	28/28	0.90	0.23	3.02	48,56,67,67	0

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
17	BO2	V	301	28/28	0.92	0.22	2.17	51,58,85,87	0
17	BO2	Y	301	28/28	0.95	0.19	2.04	49,60,68,69	0
17	BO2	H	301	28/28	0.93	0.20	1.56	47,54,72,76	0
17	BO2	K	301	28/28	0.94	0.17	1.48	43,56,63,65	0
16	CL	b	202	1/1	0.97	0.13	-0.04	56,56,56,56	0
16	CL	N	203	1/1	0.99	0.11	-0.36	47,47,47,47	0
15	MG	V	302	1/1	0.96	0.10	-1.21	64,64,64,64	0
15	MG	Y	302	1/1	0.97	0.09	-1.49	54,54,54,54	0
15	MG	N	202	1/1	0.98	0.06	-2.02	47,47,47,47	0
15	MG	K	302	1/1	0.98	0.07	-2.06	62,62,62,62	0
15	MG	G	301	1/1	0.98	0.04	-2.67	59,59,59,59	0
16	CL	G	302	1/1	0.97	0.16	-	53,53,53,53	0
16	CL	U	301	1/1	0.97	0.23	-	55,55,55,55	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.