



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:30 AM GMT

PDB ID : 2QZC  
Title : Crystal structure of a putative tena-like thiaminase (tena-1, sso2206) from  
sulfolobus solfataricus p2 at 1.50 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-08-16  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	<div> <div>5%</div> <div>92%</div> <div>5% ..</div> </div>
1	B	214	<div> <div>23%</div> <div>87%</div> <div>7% 5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	219	-	-	-	X
3	GOL	B	216	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3856 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Transcriptional activator TenA-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	209	Total	C	N	O	S	Se	0	19	0
			1805	1198	276	326	2	3			
1	B	204	Total	C	N	O	S	Se	0	10	0
			1687	1118	267	297	2	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP Q97WL0
B	0	GLY	-	LEADER SEQUENCE	UNP Q97WL0

- Molecule 2 is IMIDAZOLE (three-letter code: IMD) (formula: C<sub>3</sub>H<sub>5</sub>N<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			5	3	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	209	Total	O	0	0
			209	209		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	103	Total 103	O 103	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.86 Å 88.86 Å 135.76 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.15 – 1.50 38.14 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.15-1.50) 99.8 (38.14-1.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 1.50 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.165 , 0.184 0.175 , 0.192	Depositor DCC
$R_{free}$ test set	4372 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.8	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 55.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 87231 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3856	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.77	2/1908 (0.1%)	0.84	4/2584 (0.2%)
1	B	0.67	0/1758	0.77	4/2389 (0.2%)
All	All	0.72	2/3666 (0.1%)	0.81	8/4973 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	136[A]	CYS	CB-SG	-6.23	1.71	1.82
1	A	136[B]	CYS	CB-SG	-6.23	1.71	1.82

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	196	ARG	NE-CZ-NH2	-6.01	117.30	120.30
1	B	207	ARG	NE-CZ-NH1	5.64	123.12	120.30
1	A	114	TYR	CB-CG-CD1	5.31	124.19	121.00
1	B	116[A]	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	B	116[B]	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	A	202	TRP	CA-CB-CG	5.14	123.46	113.70
1	B	53	ASP	CB-CG-OD2	-5.11	113.70	118.30
1	A	207	ARG	NE-CZ-NH1	5.05	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1805	0	1828	7	0
1	B	1687	0	1664	16	0
2	A	5	0	5	0	0
2	B	5	0	5	0	0
3	A	30	0	40	3	0
3	B	12	0	16	1	0
4	A	209	0	0	1	0
4	B	103	0	0	1	0
All	All	3856	0	3558	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110[B]:ILE:CD1	1:B:204:ALA:HB2	2.05	0.87
1:B:110[B]:ILE:HD11	1:B:204:ALA:CB	2.07	0.84
1:B:27:ILE:HD11	1:B:202[B]:TRP:CZ3	2.15	0.82
1:B:110[B]:ILE:HD12	1:B:200:GLY:O	1.79	0.82
1:B:110[B]:ILE:HD13	1:B:204:ALA:HB2	1.67	0.77
1:B:37:LEU:HD11	1:B:95:LEU:HD21	1.68	0.76
1:A:55[B]:LEU:HD11	1:A:78:PHE:CZ	2.22	0.75
1:B:110[B]:ILE:HD11	1:B:204:ALA:HB3	1.68	0.74
1:B:110[B]:ILE:CD1	1:B:204:ALA:CB	2.70	0.63
1:B:110[B]:ILE:CD1	1:B:200:GLY:O	2.50	0.60
1:B:27:ILE:HD11	1:B:202[B]:TRP:CH2	2.40	0.55
1:B:110[B]:ILE:HD12	1:B:200:GLY:C	2.28	0.53
1:A:136[A]:CYS:HG	3:A:215:GOL:HO2	1.55	0.53
1:A:136[A]:CYS:SG	3:A:215:GOL:O2	2.68	0.51
1:A:51[A]:VAL:HG21	4:A:320:HOH:O	2.12	0.50
1:B:11:ILE:O	1:B:14:VAL:HG22	2.14	0.47
1:A:38[B]:ASP:OD2	1:A:208:LYS:NZ	2.33	0.46
1:B:27:ILE:HD13	1:B:143:VAL:HG22	1.98	0.45
1:B:51[B]:VAL:HG21	4:B:281:HOH:O	2.15	0.45
1:B:14:VAL:HG12	1:B:192:ILE:HG12	1.99	0.45
1:A:3:ILE:HD12	1:A:11:ILE:HD12	2.00	0.44
1:B:136[B]:CYS:SG	3:B:215:GOL:C3	3.06	0.43
1:A:51[A]:VAL:HG22	3:A:215:GOL:H32	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/214 (106%)	224 (99%)	2 (1%)	0	100	100
1	B	210/214 (98%)	207 (99%)	3 (1%)	0	100	100
All	All	436/428 (102%)	431 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/187 (107%)	199 (100%)	1 (0%)	92	81
1	B	179/187 (96%)	176 (98%)	3 (2%)	68	37
All	All	379/374 (101%)	375 (99%)	4 (1%)	84	58

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ILE
1	B	114	TYR
1	B	116[A]	ARG
1	B	116[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	IMD	A	214	-	3,5,5	0.55	0	4,5,5	1.28	0
3	GOL	A	215	-	5,5,5	0.57	0	5,5,5	1.49	1 (20%)
3	GOL	A	216	-	5,5,5	0.85	0	5,5,5	1.34	0
3	GOL	A	217	-	5,5,5	0.24	0	5,5,5	0.51	0
3	GOL	A	218	-	5,5,5	0.39	0	5,5,5	0.29	0
3	GOL	A	219	-	5,5,5	0.48	0	5,5,5	0.51	0
2	IMD	B	214	-	3,5,5	0.70	0	4,5,5	1.02	0
3	GOL	B	215	-	5,5,5	0.40	0	5,5,5	0.33	0
3	GOL	B	216	-	5,5,5	0.35	0	5,5,5	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMD	A	214	-	-	0/0/0/0	0/1/1/1
3	GOL	A	215	-	-	0/4/4/4	0/0/0/0
3	GOL	A	216	-	-	0/4/4/4	0/0/0/0
3	GOL	A	217	-	-	0/4/4/4	0/0/0/0
3	GOL	A	218	-	-	0/4/4/4	0/0/0/0
3	GOL	A	219	-	-	0/4/4/4	0/0/0/0
2	IMD	B	214	-	-	0/0/0/0	0/1/1/1
3	GOL	B	215	-	-	0/4/4/4	0/0/0/0
3	GOL	B	216	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	215	GOL	C3-C2-C1	2.30	120.14	111.12

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	215	GOL	3	0
3	B	215	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	206/214 (96%)	0.35	11 (5%) 30 31	13, 19, 38, 65	0
1	B	201/214 (93%)	1.37	49 (24%) 1 1	19, 33, 52, 73	0
All	All	407/428 (95%)	0.86	60 (14%) 3 3	13, 25, 50, 73	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	85	LEU	9.6
1	B	87	THR	7.9
1	A	3	ILE	6.4
1	B	95	LEU	6.3
1	B	92	TYR	6.3
1	B	161	PHE	5.8
1	B	88	HIS	5.7
1	B	89	GLY	5.5
1	A	80	SER	5.3
1	A	84	GLY	5.2
1	B	80	SER	4.8
1	B	93	SER	4.8
1	A	79	SER	4.8
1	B	12[A]	ASN	4.2
1	B	79	SER	4.1
1	B	110[A]	ILE	4.1
1	B	97	ILE	3.9
1	B	100	ASP	3.7
1	B	98	SER	3.6
1	B	165[A]	THR	3.6
1	B	163	ALA	3.5
1	B	101	VAL	3.3
1	B	86	GLU	3.3
1	B	114	TYR	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	176	TYR	3.3
1	B	9	ASN	3.2
1	B	118	LEU	3.1
1	B	112	TYR	3.1
1	B	157	THR	3.1
1	B	113	ALA	3.1
1	B	96	ASP	3.1
1	B	94	LYS	3.1
1	B	90	LYS	3.0
1	B	60	ILE	2.9
1	B	27	ILE	2.9
1	B	91	LEU	2.9
1	B	119	TYR	2.9
1	B	5	GLY	2.7
1	B	59	LEU	2.7
1	B	124	LEU	2.7
1	B	120	TYR	2.6
1	B	37	LEU	2.6
1	B	146	TYR	2.6
1	B	148	ILE	2.5
1	B	28	LEU	2.4
1	B	99	ARG	2.4
1	A	4	VAL	2.4
1	B	117	HIS	2.4
1	B	15	GLY	2.4
1	A	121	TYR	2.3
1	A	124	LEU	2.2
1	A	122	ALA	2.2
1	B	154	VAL	2.2
1	B	115	THR	2.2
1	B	138	PHE	2.1
1	B	14	VAL	2.1
1	B	121	TYR	2.1
1	A	129	PHE	2.0
1	B	19	ASN	2.0
1	A	120	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	219	6/6	0.85	0.24	3.93	43,59,67,71	0
3	GOL	B	216	6/6	0.76	0.24	2.31	60,66,67,75	0
3	GOL	A	215	6/6	0.86	0.14	1.69	39,41,45,48	0
3	GOL	A	216	6/6	0.74	0.18	0.94	28,33,38,48	0
3	GOL	A	217	6/6	0.90	0.15	0.83	34,47,51,59	0
3	GOL	B	215	6/6	0.78	0.15	0.66	67,69,70,71	0
2	IMD	B	214	5/5	0.94	0.11	-0.35	26,29,32,34	0
2	IMD	A	214	5/5	0.98	0.05	-2.07	20,20,21,22	0
3	GOL	A	218	6/6	0.74	0.28	-	62,68,68,70	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.