



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:31 AM GMT

PDB ID : 2R0L  
Title : Short Form HGFA with Inhibitory Fab75  
Authors : Eigenbrot, C.; Shia, S.  
Deposited on : 2007-08-20  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	214	<div> <div>89%</div> <div>10% •</div> </div>
2	H	220	<div> <div>5%</div> <div>87%</div> <div>11% •</div> </div>
3	A	248	<div> <div>88%</div> <div>8% • •</div> </div>
4	B	35	<div> <div>3%</div> <div>26%</div> <div>71%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	A	741	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called antibody light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	214	Total	C	N	O	S	7	0	0
			1636	1024	271	335	6			

- Molecule 2 is a protein called antibody heavy chain, Fab portion only.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	220	Total	C	N	O	S	0	1	0
			1649	1045	277	321	6			

- Molecule 3 is a protein called Hepatocyte growth factor activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	239	Total	C	N	O	S	0	0	0
			1824	1156	316	339	13			

- Molecule 4 is a protein called Hepatocyte growth factor activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	10	Total	C	N	O	S	0	0	0
			78	44	22	11	1			

- Molecule 5 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	66	Total	O	0	0
			66	66		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	4	Total 4	O 4	0	0
6	H	63	Total 63	O 63	0	0
6	L	66	Total 66	O 66	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: antibody light chain

Chain L: 




- Molecule 2: antibody heavy chain, Fab portion only

Chain H: 




- Molecule 3: Hepatocyte growth factor activator

Chain A: 



- Molecule 4: Hepatocyte growth factor activator

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.82Å 48.23Å 97.05Å 98.39° 96.24° 100.77°	Depositor
Resolution (Å)	47.51 – 2.20 46.69 – 2.20	Depositor EDS
% Data completeness (in resolution range)	94.4 (47.51-2.20) 90.2 (46.69-2.20)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.89 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.1.07	Depositor
R, $R_{free}$	0.193 , 0.248 0.191 , 0.242	Depositor DCC
$R_{free}$ test set	1535 reflections (4.95%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.6	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 40.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 32514 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5425	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.61	2/1672 (0.1%)	0.87	6/2273 (0.3%)
2	H	0.41	0/1693	0.69	2/2311 (0.1%)
3	A	0.42	0/1878	0.70	5/2558 (0.2%)
4	B	0.37	0/79	0.57	0/102
All	All	0.48	2/5322 (0.0%)	0.75	13/7244 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	24	ARG	CD-NE	-14.76	1.21	1.46
1	L	142	ARG	NE-CZ	10.42	1.46	1.33

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	142	ARG	NE-CZ-NH1	-15.97	112.31	120.30
1	L	142	ARG	NE-CZ-NH2	14.10	127.35	120.30
1	L	24	ARG	CD-NE-CZ	-8.68	111.44	123.60
1	L	142	ARG	CD-NE-CZ	6.34	132.48	123.60
2	H	58	ASP	CB-CG-OD2	5.77	123.49	118.30
3	A	186	ASP	CB-CG-OD2	5.59	123.33	118.30
3	A	145	ASP	CB-CG-OD2	5.54	123.29	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	185	ASP	CB-CG-OD2	5.44	123.20	118.30
3	A	102	ASP	CB-CG-OD2	5.37	123.14	118.30
3	A	39	ASP	CB-CG-OD2	5.35	123.12	118.30
3	A	175	ASP	CB-CG-OD2	5.30	123.07	118.30
2	H	208	ASP	CB-CG-OD2	5.19	122.97	118.30
1	L	82	ASP	CB-CG-OD2	5.16	122.94	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	L	142	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1636	0	1586	11	0
2	H	1649	0	1605	16	0
3	A	1824	0	1737	9	0
4	B	78	0	76	0	0
5	A	39	0	34	0	0
6	A	66	0	0	0	0
6	B	4	0	0	0	0
6	H	63	0	0	0	0
6	L	66	0	0	0	0
All	All	5425	0	5038	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:211:ARG:HG2	1:L:211:ARG:HH11	1.51	0.74

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:97:TRP:CZ3	2:H:98:ARG:HD2	2.30	0.66
2:H:126:PRO:O	2:H:127:SER:HB2	1.97	0.63
1:L:198:HIS:CD2	1:L:200:GLY:H	2.17	0.63
3:A:170:SER:HB3	3:A:170(B):GLU:OE2	2.00	0.61
2:H:36:TRP:CD1	2:H:69[B]:ILE:HD12	2.36	0.60
2:H:82:MET:HE2	2:H:82(C):LEU:HD21	1.86	0.58
2:H:36:TRP:HD1	2:H:69[B]:ILE:HD12	1.70	0.56
1:L:211:ARG:NH1	1:L:211:ARG:HG2	2.20	0.56
3:A:51:TRP:CE3	3:A:105:LEU:HG	2.41	0.55
3:A:181:LEU:HD23	3:A:230:ARG:HG3	1.89	0.54
2:H:97:TRP:HZ3	2:H:98:ARG:HD2	1.70	0.54
1:L:151:ASP:OD2	1:L:189:HIS:HD2	1.91	0.54
1:L:140:TYR:CG	1:L:141:PRO:HA	2.45	0.52
2:H:11:LEU:HB2	2:H:147:PRO:HG3	1.94	0.49
3:A:17:ILE:HD11	3:A:221(A):GLY:HA3	1.94	0.49
2:H:131:THR:OG1	2:H:132:SER:N	2.46	0.48
2:H:69[A]:ILE:HG23	2:H:80:LEU:HD13	1.97	0.47
1:L:125:LEU:HG	1:L:183:LYS:HE3	1.97	0.47
1:L:141:PRO:O	1:L:198:HIS:HE1	2.00	0.45
2:H:35:HIS:O	2:H:92:CYS:HA	2.17	0.45
2:H:51:ILE:HG22	2:H:69[B]:ILE:HD11	1.98	0.44
3:A:144:LEU:HD21	3:A:156:ARG:HD2	1.99	0.44
3:A:211:GLY:HA2	3:A:229:THR:O	2.17	0.44
2:H:82:MET:HB3	2:H:82(C):LEU:HD21	1.99	0.43
2:H:12:VAL:HG21	2:H:18:LEU:HB2	2.00	0.43
2:H:40:ALA:HB3	2:H:43:LYS:HG2	2.01	0.42
3:A:144:LEU:CD2	3:A:156:ARG:HD2	2.48	0.42
1:L:151:ASP:OD2	1:L:189:HIS:CD2	2.71	0.42
2:H:69[A]:ILE:HG22	2:H:79:TYR:O	2.20	0.41
1:L:50:SER:HB3	1:L:53:PHE:CD2	2.55	0.41
3:A:61:ARG:HD3	3:A:87:LYS:HA	2.03	0.41
3:A:184:TYR:CD1	3:A:188:LYS:HB3	2.56	0.41
1:L:16:GLY:HA2	1:L:77:SER:OG	2.21	0.41
2:H:59:TYR:CE1	2:H:69[A]:ILE:HG12	2.56	0.40
1:L:137:ASN:HA	1:L:137:ASN:HD22	1.72	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	212/214 (99%)	208 (98%)	4 (2%)	0	100	100
2	H	219/220 (100%)	205 (94%)	12 (6%)	2 (1%)	21	19
3	A	237/248 (96%)	231 (98%)	4 (2%)	2 (1%)	24	22
4	B	8/35 (23%)	6 (75%)	1 (12%)	1 (12%)	0	0
All	All	676/717 (94%)	650 (96%)	21 (3%)	5 (1%)	26	25

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	127	SER
2	H	215	SER
3	A	188	LYS
4	B	389	GLY
3	A	171	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	188/188 (100%)	182 (97%)	6 (3%)	46	57
2	H	182/181 (101%)	174 (96%)	8 (4%)	35	42
3	A	199/207 (96%)	198 (100%)	1 (0%)	92	96
4	B	7/30 (23%)	7 (100%)	0	100	100
All	All	576/606 (95%)	561 (97%)	15 (3%)	57	66

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	83	PHE
1	L	135	LEU
1	L	137	ASN
1	L	181	LEU
1	L	182	SER
1	L	201	LEU
2	H	19	ARG
2	H	43	LYS
2	H	69[A]	ILE
2	H	69[B]	ILE
2	H	97	TRP
2	H	128	SER
2	H	150	VAL
2	H	199	ASN
3	A	170(B)	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	3	GLN
1	L	137	ASN
1	L	138	ASN
1	L	189	HIS
1	L	198	HIS
2	H	199	ASN
3	A	60(A)	HIS
3	A	70	GLN
3	A	116	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

3 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	NAG	A	741	3,5	14,14,15	0.55	0	15,19,21	0.92	1 (6%)
5	NAG	A	742	5	14,14,15	0.48	0	15,19,21	2.27	5 (33%)
5	BMA	A	743	5	11,11,12	0.58	0	14,15,17	1.43	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	741	3,5	-	0/6/23/26	0/1/1/1
5	NAG	A	742	5	-	0/6/23/26	0/1/1/1
5	BMA	A	743	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	742	NAG	C4-C3-C2	-5.76	102.28	111.23
5	A	742	NAG	C2-N2-C7	-3.14	119.01	123.04
5	A	741	NAG	C2-N2-C7	-2.28	120.11	123.04
5	A	742	NAG	O4-C4-C3	2.01	114.86	110.34
5	A	742	NAG	O4-C4-C5	2.34	115.44	109.24
5	A	743	BMA	C1-O5-C5	2.43	115.33	112.25
5	A	743	BMA	C1-C2-C3	4.00	114.27	109.54
5	A	742	NAG	C1-O5-C5	4.32	117.73	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	214/214 (100%)	-0.37	1 (0%) 91 91	10, 21, 39, 79	2 (0%)
2	H	220/220 (100%)	0.04	12 (5%) 29 28	6, 20, 46, 103	0
3	A	239/248 (96%)	-0.32	1 (0%) 93 93	8, 18, 43, 71	1 (0%)
4	B	10/35 (28%)	0.56	1 (10%) 9 8	37, 75, 91, 103	0
All	All	683/717 (95%)	-0.21	15 (2%) 65 64	6, 20, 45, 103	3 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	133	GLY	12.7
2	H	130	SER	12.7
2	H	129	LYS	11.4
2	H	216	CYS	9.4
2	H	131	THR	8.4
2	H	132	SER	7.3
1	L	214	CYS	5.7
2	H	127	SER	3.3
2	H	134	GLY	3.2
3	A	243	ARG	3.2
2	H	215	SER	3.0
2	H	187	SER	2.9
2	H	128	SER	2.9
4	B	388	PRO	2.2
2	H	97	TRP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	A	741	14/15	0.74	0.19	3.92	53,63,68,68	0
5	NAG	A	742	14/15	0.75	0.30	-	70,75,83,83	0
5	BMA	A	743	11/12	0.71	0.34	-	77,78,80,80	0

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.