



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:23 PM GMT

PDB ID : 3R2A  
Title : Crystal structure of RXRalpha ligand-binding domain complexed with corepressor SMRT2 and antagonist rhein  
Authors : Zhang, H.; Chen, L.; Chen, J.; Jiang, H.; Shen, X.  
Deposited on : 2011-03-14  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	 2% 43% 39% 6% 12%
1	B	240	 2% 33% 45% 11% 10%
1	C	240	 % 41% 36% 8% 15%
1	D	240	 2% 31% 46% 13% 9%
2	E	16	 38% 6% 56%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	16	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	RHN	A	1	-	-	-	X
3	RHN	C	2	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6940 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Retinoic acid receptor RXR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1666	1069	287	301	9			
1	B	217	Total	C	N	O	S	0	0	0
			1717	1099	296	311	11			
1	C	203	Total	C	N	O	S	0	0	0
			1607	1034	276	288	9			
1	D	219	Total	C	N	O	S	0	0	0
			1733	1111	298	313	11			

- Molecule 2 is a protein called Nuclear receptor corepressor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	7	Total	C	N	O	S	0	0	0
			50	31	8	10	1			
2	F	9	Total	C	N	O	S	0	0	0
			65	41	10	13	1			

- Molecule 3 is 4,5-DIHYDROXY-9,10-DIOXO-9,10-DIHYDROANTHRACENE-2-CARBOXYLIC ACID (three-letter code: RHN) (formula: C<sub>15</sub>H<sub>8</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			21	15	6		
3	C	1	Total	C	O	0	0
			21	15	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	17	Total	O	0	0
			17	17		
4	B	15	Total	O	0	0
			15	15		
4	C	13	Total	O	0	0
			13	13		
4	D	12	Total	O	0	0
			12	12		
4	E	2	Total	O	0	0
			2	2		
4	F	1	Total	O	0	0
			1	1		







• Molecule 1: Retinoic acid receptor RXR-alpha



• Molecule 2: Nuclear receptor corepressor 2



• Molecule 2: Nuclear receptor corepressor 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.03Å 117.58Å 117.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.52 – 3.00 44.70 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.52-3.00) 93.6 (44.70-3.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.76 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.254 , 0.335 0.268 , 0.278	Depositor DCC
$R_{free}$ test set	1191 reflections (5.50%)	DCC
Wilson B-factor (Å <sup>2</sup> )	73.9	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 47.6	EDS
Estimated twinning fraction	0.447 for -h,l,k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 23832 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6940	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RHN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.38	2/1699 (0.1%)	0.93	3/2297 (0.1%)
1	B	1.25	4/1751 (0.2%)	1.09	12/2367 (0.5%)
1	C	1.40	3/1639 (0.2%)	1.21	5/2214 (0.2%)
1	D	1.18	1/1768 (0.1%)	1.02	9/2389 (0.4%)
2	E	0.87	0/49	1.04	1/64 (1.6%)
2	F	0.89	0/64	1.06	0/85
All	All	1.30	10/6970 (0.1%)	1.06	30/9416 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	371	ARG	CZ-NH1	-10.83	1.19	1.33
1	B	371	ARG	CG-CD	-8.97	1.29	1.51
1	B	371	ARG	CD-NE	-7.02	1.34	1.46
1	C	366	GLU	CD-OE2	6.94	1.33	1.25
1	A	369	CYS	CB-SG	-6.76	1.70	1.82

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	366	GLU	OE1-CD-OE2	-27.93	89.78	123.30
1	C	366	GLU	CG-CD-OE2	17.80	153.90	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	371	ARG	CG-CD-NE	12.84	138.77	111.80
1	B	371	ARG	NE-CZ-NH1	-8.72	115.94	120.30
1	B	371	ARG	CD-NE-CZ	-8.32	111.95	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	443	GLY	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1666	0	1701	142	0
1	B	1717	0	1736	262	3
1	C	1607	0	1639	145	1
1	D	1733	0	1764	270	2
2	E	50	0	50	14	0
2	F	65	0	68	17	0
3	A	21	0	5	3	0
3	C	21	0	6	3	0
4	A	17	0	0	3	0
4	B	15	0	0	0	0
4	C	13	0	0	1	0
4	D	12	0	0	3	0
4	E	2	0	0	0	0
4	F	1	0	0	0	0
All	All	6940	0	6969	803	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 58.

The worst 5 of 803 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:324:ILE:HG13	1:B:346:PHE:CE1	1.44	1.52
1:D:236:LEU:O	1:D:240:LEU:CD1	1.65	1.43
1:D:385:ASN:CG	1:D:388:GLU:HB2	1.38	1.43
1:D:305:TRP:N	1:D:308:LEU:HD23	1.30	1.38
1:B:381:LYS:CE	1:C:445:THR:OG1	1.63	1.37

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:285:ARG:NH1	1:D:237:GLU:OE1[1_655]	2.00	0.20
1:B:285:ARG:NH1	1:D:237:GLU:OE2[1_655]	2.15	0.05
1:B:321:LYS:NZ	1:C:273:ASP:OD1[3_454]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/240 (86%)	199 (96%)	6 (3%)	2 (1%)	19	61
1	B	213/240 (89%)	203 (95%)	8 (4%)	2 (1%)	21	64
1	C	199/240 (83%)	189 (95%)	6 (3%)	4 (2%)	9	41
1	D	215/240 (90%)	208 (97%)	5 (2%)	2 (1%)	21	64
2	E	5/16 (31%)	4 (80%)	1 (20%)	0	100	100
2	F	7/16 (44%)	6 (86%)	1 (14%)	0	100	100
All	All	846/992 (85%)	809 (96%)	27 (3%)	10 (1%)	16	56

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	445	THR
1	C	445	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	305	TRP
1	D	447	ILE
1	A	446	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/207 (87%)	166 (92%)	15 (8%)	14	46
1	B	187/207 (90%)	153 (82%)	34 (18%)	2	11
1	C	174/207 (84%)	156 (90%)	18 (10%)	9	33
1	D	189/207 (91%)	155 (82%)	34 (18%)	2	11
2	E	5/12 (42%)	3 (60%)	2 (40%)	0	0
2	F	7/12 (58%)	4 (57%)	3 (43%)	0	0
All	All	743/852 (87%)	637 (86%)	106 (14%)	4	19

5 of 106 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	459	HIS
1	C	390	GLU
1	D	455	LEU
1	C	229	ASP
1	C	318	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	411	GLN
1	C	385	ASN
1	D	411	GLN
1	B	361	GLN
1	D	338	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	RHN	A	1	-	20,23,23	1.55	4 (20%)	30,35,35	0.84	1 (3%)
3	RHN	C	2	-	20,23,23	1.56	4 (20%)	30,35,35	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RHN	A	1	-	-	0/0/20/20	0/3/3/3
3	RHN	C	2	-	-	0/0/20/20	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2	RHN	CAR-CAP	-4.02	1.39	1.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1	RHN	CAR-CAP	-3.78	1.40	1.48
3	C	2	RHN	CAS-CAP	-3.69	1.40	1.48
3	A	1	RHN	CAS-CAP	-3.69	1.40	1.48
3	A	1	RHN	CAT-CAQ	-2.95	1.40	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1	RHN	OAE-CAN-CAT	-2.24	116.76	121.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1	RHN	3	0
3	C	2	RHN	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	211/240 (87%)	0.09	4 (1%) 70 41	20, 62, 94, 103	0
1	B	217/240 (90%)	0.24	4 (1%) 71 43	20, 75, 112, 122	0
1	C	203/240 (84%)	0.13	3 (1%) 76 49	20, 61, 86, 99	0
1	D	219/240 (91%)	0.30	5 (2%) 64 33	20, 75, 104, 116	0
2	E	7/16 (43%)	0.52	0 100 100	101, 102, 103, 104	0
2	F	9/16 (56%)	0.28	0 100 100	99, 100, 100, 101	0
All	All	866/992 (87%)	0.19	16 (1%) 71 43	20, 68, 100, 122	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	267	ASN	5.7
1	B	341	GLY	4.2
1	A	322	ASP	3.4
1	C	445	THR	3.0
1	D	271	ALA	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	RHN	C	2	21/21	0.71	0.92	11.60	138,139,139,139	0
3	RHN	A	1	21/21	0.75	0.62	4.95	129,129,129,129	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.