



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:53 PM GMT

PDB ID : 1R38  
Title : Crystal structure of H114A mutant of *Candida tenuis* xylose reductase  
Authors : Kratzer, R.; Kavanagh, K.L.; Wilson, D.K.; Nidetzky, B.  
Deposited on : 2003-09-30  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	322	<div> <div>%</div> <div> <div></div> <div>93%</div> <div>5% •</div> </div> </div>
1	B	322	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>9% •</div> </div> </div>
1	C	322	<div> <div></div> <div> <div></div> <div>89%</div> <div>10% •</div> </div> </div>
1	D	322	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>7% •</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11127 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called xylose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2527	1642	417	464	4			
1	B	319	Total	C	N	O	S	0	0	0
			2527	1642	417	464	4			
1	C	319	Total	C	N	O	S	0	0	0
			2527	1642	417	464	4			
1	D	319	Total	C	N	O	S	0	0	0
			2527	1642	417	464	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	114	ALA	HIS	ENGINEERED	UNP O74237
B	114	ALA	HIS	ENGINEERED	UNP O74237
C	114	ALA	HIS	ENGINEERED	UNP O74237
D	114	ALA	HIS	ENGINEERED	UNP O74237

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C<sub>21</sub>H<sub>28</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	189	Total	O	0	0
			189	189		
3	B	188	Total	O	0	0
			188	188		
3	C	206	Total	O	0	0
			206	206		
3	D	244	Total	O	0	0
			244	244		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: xylose reductase





- Molecule 1: xylose reductase



- Molecule 1: xylose reductase



- Molecule 1: xylose reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	179.98Å 129.09Å 79.87Å 90.00° 90.82° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 19.58 – 2.18	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.20) 97.2 (19.58-2.18)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.81 (at 2.19Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.185 , 0.226 0.185 , 0.226	Depositor DCC
$R_{free}$ test set	4483 reflections (4.91%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.7	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 41.8	EDS
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 91982 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11127	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/2592	0.66	0/3520
1	B	0.51	0/2592	0.66	0/3520
1	C	0.55	0/2592	0.68	0/3520
1	D	0.58	0/2592	0.69	1/3520 (0.0%)
All	All	0.55	0/10368	0.67	1/14080 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	26	LEU	CA-CB-CG	5.12	127.07	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	52	TYR	Sidechain

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	B	217	TYR	Sidechain
1	C	52	TYR	Sidechain
1	D	46	PHE	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2527	0	2523	14	0
1	B	2527	0	2523	23	0
1	C	2527	0	2523	21	0
1	D	2527	0	2523	14	0
2	A	48	0	25	4	0
2	B	48	0	25	5	0
2	C	48	0	25	3	0
2	D	48	0	25	3	0
3	A	189	0	0	0	0
3	B	188	0	0	3	0
3	C	206	0	0	2	0
3	D	244	0	0	0	0
All	All	11127	0	10192	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:PHE:HD1	2:B:951:NAP:H52A	1.40	0.86
1:C:241:ALA:HA	1:C:246:LYS:HE3	1.65	0.77
1:A:220:PHE:HD1	2:A:950:NAP:H52A	1.52	0.74
1:D:220:PHE:HD1	2:D:953:NAP:H52A	1.55	0.70
1:C:278:PRO:O	1:C:281:LEU:HB3	1.91	0.69
1:C:220:PHE:HD1	2:C:952:NAP:H52A	1.59	0.68
1:C:294:LYS:O	1:C:298:GLU:HG3	1.96	0.66
1:D:274:LYS:O	2:D:953:NAP:H8A	2.04	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:294:LYS:O	1:B:298:GLU:HG3	2.04	0.58
1:B:194:HIS:HB3	1:B:216:ALA:HB1	1.85	0.57
1:C:194:HIS:HB3	1:C:216:ALA:HB1	1.85	0.57
1:C:191:GLN:OE1	2:C:952:NAP:H2N	2.05	0.57
1:B:220:PHE:CD1	2:B:951:NAP:H52A	2.31	0.57
1:B:133:TYR:CZ	1:B:135:GLY:HA2	2.42	0.54
1:B:120:LYS:HA	1:B:143:GLU:CG	2.36	0.54
1:A:278:PRO:O	1:A:281:LEU:HB3	2.06	0.54
1:C:279:GLU:HG2	1:C:283:GLN:NE2	2.22	0.53
1:B:90:LYS:HE3	3:B:1095:HOH:O	2.10	0.52
1:C:207:ALA:HB3	1:C:214:ILE:HD11	1.90	0.52
1:A:194:HIS:HB3	1:A:216:ALA:HB1	1.92	0.52
1:C:74:GLU:HB3	3:C:1078:HOH:O	2.10	0.52
1:A:220:PHE:CD1	2:A:950:NAP:H52A	2.39	0.52
1:D:294:LYS:HE2	1:D:298:GLU:OE2	2.10	0.51
1:D:180:LEU:HD23	1:D:186:LYS:HD2	1.92	0.50
1:B:25:LYS:HE3	1:B:228:MET:SD	2.52	0.50
1:C:110:LEU:HA	1:C:165:SER:O	2.12	0.49
1:A:249:ALA:HB1	1:A:254:LYS:O	2.11	0.49
1:C:45:LEU:HD23	1:C:45:LEU:C	2.32	0.49
1:B:224:SER:HB2	2:B:951:NAP:O3	2.13	0.49
1:A:120:LYS:HA	1:A:143:GLU:CG	2.42	0.49
1:D:180:LEU:CD2	1:D:186:LYS:HD2	2.43	0.48
1:D:116:PRO:HG2	1:D:320:ILE:HD12	1.96	0.47
1:C:274:LYS:O	2:C:952:NAP:H8A	2.14	0.47
1:D:120:LYS:HA	1:D:143:GLU:CG	2.45	0.47
1:C:28:ASN:O	1:C:57:GLU:HG2	2.15	0.47
1:C:20:GLY:HA3	1:C:45:LEU:HD22	1.97	0.47
1:B:191:GLN:OE1	2:B:951:NAP:H2N	2.15	0.46
1:D:133:TYR:CZ	1:D:135:GLY:HA2	2.51	0.46
1:A:191:GLN:OE1	2:A:950:NAP:H2N	2.16	0.46
1:D:251:LYS:HE3	1:D:252:TYR:CZ	2.51	0.46
1:D:191:GLN:OE1	2:D:953:NAP:H2N	2.16	0.45
1:B:277:LEU:HG	1:B:279:GLU:HG2	1.98	0.45
1:B:56:LYS:HE3	3:B:993:HOH:O	2.16	0.45
1:B:279:GLU:O	1:B:283:GLN:HG3	2.17	0.45
1:D:180:LEU:O	1:D:186:LYS:HE3	2.17	0.44
1:C:194:HIS:HB3	1:C:216:ALA:CB	2.48	0.44
1:C:256:PRO:HD2	3:C:1149:HOH:O	2.17	0.44
1:C:277:LEU:HA	1:C:278:PRO:HD3	1.91	0.43
1:B:109:ASP:HA	1:B:164:LYS:HB2	1.99	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:274:LYS:O	2:B:951:NAP:H8A	2.18	0.43
1:B:120:LYS:HD3	3:B:994:HOH:O	2.17	0.43
1:C:116:PRO:HG2	1:C:320:ILE:HD12	2.00	0.42
1:A:294:LYS:O	1:A:298:GLU:HG3	2.20	0.42
1:B:120:LYS:HA	1:B:143:GLU:HG3	2.00	0.42
1:D:311:ASP:HA	1:D:312:PRO:HD3	1.82	0.42
1:A:285:ARG:NH1	1:A:285:ARG:HG3	2.34	0.42
1:D:303:LEU:N	1:D:303:LEU:HD23	2.34	0.42
1:A:285:ARG:HH11	1:A:285:ARG:HG3	1.85	0.41
1:A:285:ARG:O	1:A:285:ARG:HG3	2.19	0.41
1:B:255:THR:OG1	1:B:258:GLU:HG3	2.20	0.41
1:B:120:LYS:HA	1:B:143:GLU:HG2	2.02	0.41
1:D:194:HIS:HB3	1:D:216:ALA:HB1	2.03	0.41
1:A:5:ILE:HG23	1:A:19:ILE:HB	2.03	0.41
1:B:249:ALA:HB1	1:B:254:LYS:O	2.20	0.41
1:B:274:LYS:C	1:B:274:LYS:HD3	2.41	0.41
1:C:199:GLN:HG2	1:C:263:TRP:CH2	2.55	0.41
1:B:116:PRO:HG2	1:B:320:ILE:HD12	2.02	0.41
1:C:120:LYS:HA	1:C:143:GLU:CG	2.50	0.41
1:A:45:LEU:HD23	1:A:45:LEU:C	2.42	0.40
1:A:274:LYS:O	2:A:950:NAP:H8A	2.22	0.40
1:B:194:HIS:HB3	1:B:216:ALA:CB	2.49	0.40
1:C:107:TYR:CD1	1:C:107:TYR:C	2.94	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/322 (98%)	309 (98%)	8 (2%)	0	100	100
1	B	317/322 (98%)	308 (97%)	9 (3%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	317/322 (98%)	309 (98%)	8 (2%)	0	100	100
1	D	317/322 (98%)	313 (99%)	4 (1%)	0	100	100
All	All	1268/1288 (98%)	1239 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/269 (99%)	266 (100%)	1 (0%)	93	97
1	B	267/269 (99%)	265 (99%)	2 (1%)	88	94
1	C	267/269 (99%)	265 (99%)	2 (1%)	88	94
1	D	267/269 (99%)	265 (99%)	2 (1%)	88	94
All	All	1068/1076 (99%)	1061 (99%)	7 (1%)	88	94

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	285	ARG
1	B	278	PRO
1	B	319	PRO
1	C	74	GLU
1	C	319	PRO
1	D	285	ARG
1	D	319	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	317	ASN
1	B	91	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	253	ASN
1	B	276	ASN
1	B	317	ASN
1	C	34	GLN
1	C	276	ASN
1	C	283	GLN
1	C	317	ASN
1	D	91	ASN
1	D	253	ASN
1	D	283	GLN
1	D	317	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAP	A	950	-	42,52,52	1.93	7 (16%)	54,80,80	1.91	12 (22%)
2	NAP	B	951	-	42,52,52	1.83	10 (23%)	54,80,80	2.03	14 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAP	C	952	-	42,52,52	1.87	8 (19%)	54,80,80	1.96	13 (24%)
2	NAP	D	953	-	42,52,52	1.92	10 (23%)	54,80,80	1.97	12 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	950	-	-	0/27/67/67	0/5/5/5
2	NAP	B	951	-	-	0/27/67/67	0/5/5/5
2	NAP	C	952	-	-	0/27/67/67	0/5/5/5
2	NAP	D	953	-	-	0/27/67/67	0/5/5/5

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	952	NAP	PA-O1A	-2.69	1.41	1.51
2	B	951	NAP	PA-O1A	-2.68	1.41	1.51
2	D	953	NAP	PA-O1A	-2.50	1.42	1.51
2	A	950	NAP	PA-O1A	-2.14	1.43	1.51
2	B	951	NAP	PN-O2N	-2.07	1.46	1.54
2	D	953	NAP	C6N-C5N	2.01	1.43	1.38
2	D	953	NAP	C5B-C4B	2.02	1.58	1.51
2	B	951	NAP	C5B-C4B	2.17	1.58	1.51
2	D	953	NAP	C5N-C4N	2.24	1.43	1.38
2	C	952	NAP	C6N-C5N	2.24	1.43	1.38
2	B	951	NAP	C6N-C5N	2.38	1.43	1.38
2	B	951	NAP	C5N-C4N	2.42	1.43	1.38
2	A	950	NAP	C4A-N3A	2.77	1.39	1.35
2	C	952	NAP	C4A-N3A	2.79	1.39	1.35
2	D	953	NAP	C4A-N3A	2.83	1.39	1.35
2	B	951	NAP	C2A-N1A	3.32	1.40	1.33
2	B	951	NAP	C6N-N1N	3.38	1.44	1.35
2	D	953	NAP	C2A-N3A	3.64	1.38	1.32
2	D	953	NAP	C2A-N1A	3.71	1.41	1.33
2	C	952	NAP	C2A-N1A	3.71	1.41	1.33
2	A	950	NAP	C6N-N1N	3.78	1.45	1.35
2	C	952	NAP	C6N-N1N	3.79	1.45	1.35
2	A	950	NAP	C2A-N1A	3.83	1.41	1.33
2	B	951	NAP	C2A-N3A	3.90	1.39	1.32
2	A	950	NAP	C4N-C3N	4.05	1.46	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	953	NAP	C6N-N1N	4.12	1.46	1.35
2	D	953	NAP	C4N-C3N	4.15	1.46	1.39
2	C	952	NAP	C4N-C3N	4.22	1.46	1.39
2	C	952	NAP	C2A-N3A	4.33	1.39	1.32
2	A	950	NAP	C2A-N3A	4.34	1.39	1.32
2	B	951	NAP	C4N-C3N	4.47	1.46	1.39
2	B	951	NAP	O4D-C1D	4.93	1.47	1.41
2	D	953	NAP	O4D-C1D	6.05	1.48	1.41
2	C	952	NAP	O4D-C1D	6.18	1.49	1.41
2	A	950	NAP	O4D-C1D	7.18	1.50	1.41

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	952	NAP	N3A-C2A-N1A	-8.03	122.74	128.89
2	A	950	NAP	N3A-C2A-N1A	-7.82	122.90	128.89
2	B	951	NAP	N3A-C2A-N1A	-7.58	123.09	128.89
2	D	953	NAP	N3A-C2A-N1A	-7.36	123.26	128.89
2	B	951	NAP	O3-PA-O5B	-5.84	87.45	102.94
2	D	953	NAP	O3-PA-O5B	-5.59	88.12	102.94
2	A	950	NAP	PN-O3-PA	-4.24	120.82	132.73
2	D	953	NAP	C1B-N9A-C4A	-4.23	120.56	126.94
2	B	951	NAP	C1B-N9A-C4A	-3.67	121.41	126.94
2	C	952	NAP	PN-O3-PA	-3.54	122.78	132.73
2	C	952	NAP	C1B-N9A-C4A	-3.44	121.75	126.94
2	A	950	NAP	C1B-N9A-C4A	-3.09	122.27	126.94
2	A	950	NAP	O7N-C7N-C3N	-2.80	116.53	119.59
2	D	953	NAP	O5B-C5B-C4B	-2.78	98.87	109.12
2	C	952	NAP	O3-PA-O5B	-2.76	95.62	102.94
2	C	952	NAP	O7N-C7N-C3N	-2.65	116.69	119.59
2	D	953	NAP	O4B-C4B-C5B	-2.63	99.90	109.32
2	B	951	NAP	O7N-C7N-C3N	-2.61	116.73	119.59
2	B	951	NAP	C5N-C4N-C3N	-2.61	117.05	120.33
2	C	952	NAP	C5N-C4N-C3N	-2.47	117.23	120.33
2	B	951	NAP	O4B-C4B-C5B	-2.46	100.53	109.32
2	A	950	NAP	C5N-C4N-C3N	-2.39	117.33	120.33
2	C	952	NAP	O4B-C4B-C5B	-2.34	100.95	109.32
2	B	951	NAP	O5B-C5B-C4B	-2.31	100.59	109.12
2	C	952	NAP	O3B-C3B-C4B	-2.30	104.15	111.05
2	A	950	NAP	O4B-C4B-C5B	-2.25	101.26	109.32
2	A	950	NAP	O3-PA-O5B	-2.22	97.05	102.94
2	B	951	NAP	PN-O3-PA	-2.20	126.54	132.73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	953	NAP	PN-O3-PA	-2.11	126.80	132.73
2	D	953	NAP	O7N-C7N-C3N	-2.10	117.30	119.59
2	A	950	NAP	O3B-C3B-C4B	-2.06	104.87	111.05
2	A	950	NAP	C3N-C7N-N7N	2.03	120.04	117.82
2	C	952	NAP	C3N-C7N-N7N	2.04	120.06	117.82
2	D	953	NAP	C3N-C7N-N7N	2.11	120.13	117.82
2	B	951	NAP	O3X-P2B-O2X	2.25	115.94	107.38
2	C	952	NAP	O3-PN-O5D	2.27	108.95	102.94
2	A	950	NAP	O2A-PA-O1A	2.41	125.57	112.53
2	C	952	NAP	O2A-PA-O1A	2.46	125.86	112.53
2	D	953	NAP	O2A-PA-O1A	2.56	126.42	112.53
2	B	951	NAP	O2A-PA-O1A	2.56	126.42	112.53
2	B	951	NAP	C3N-C7N-N7N	2.76	120.83	117.82
2	B	951	NAP	C2N-C3N-C4N	2.99	121.62	118.29
2	D	953	NAP	C2N-C3N-C4N	3.12	121.76	118.29
2	D	953	NAP	O3-PN-O5D	3.15	111.31	102.94
2	B	951	NAP	O3-PN-O5D	3.34	111.80	102.94
2	D	953	NAP	C4B-O4B-C1B	3.43	113.49	109.72
2	B	951	NAP	C4B-O4B-C1B	3.52	113.58	109.72
2	A	950	NAP	C2N-C3N-C4N	3.73	122.44	118.29
2	C	952	NAP	C2N-C3N-C4N	3.73	122.45	118.29
2	A	950	NAP	C4B-O4B-C1B	3.90	114.00	109.72
2	C	952	NAP	C4B-O4B-C1B	4.00	114.12	109.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	950	NAP	4	0
2	B	951	NAP	5	0
2	C	952	NAP	3	0
2	D	953	NAP	3	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	319/322 (99%)	-0.44	2 (0%) 90 90	14, 22, 37, 49	0
1	B	319/322 (99%)	-0.48	3 (0%) 85 85	16, 22, 36, 47	0
1	C	319/322 (99%)	-0.45	1 (0%) 94 94	13, 21, 38, 49	0
1	D	319/322 (99%)	-0.60	4 (1%) 79 78	12, 18, 31, 43	0
All	All	1276/1288 (99%)	-0.49	10 (0%) 87 87	12, 21, 36, 49	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	SER	4.2
1	B	4	SER	3.5
1	D	29	ALA	3.2
1	D	4	SER	3.1
1	B	28	ASN	2.7
1	B	29	ALA	2.6
1	C	4	SER	2.3
1	D	138	ASN	2.3
1	A	231	GLY	2.2
1	D	28	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAP	C	952	48/48	0.94	0.10	-0.18	17,28,31,35	0
2	NAP	A	950	48/48	0.95	0.09	-0.34	15,28,33,35	0
2	NAP	B	951	48/48	0.97	0.09	-0.37	17,22,25,28	0
2	NAP	D	953	48/48	0.97	0.08	-0.91	11,17,22,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.