



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:22 PM GMT

PDB ID : 3R40  
Title : Crystal Structure of the Fluoroacetate Dehalogenase RPA1163 - Asp110Asn/apo  
Authors : Chan, P.W.Y.; Yakunin, A.F.; Edwards, E.A.; Pai, E.F.  
Deposited on : 2011-03-16  
Resolution : 1.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1077 (1.12-1.00)
Clashscore	102246	1147 (1.12-1.00)
Ramachandran outliers	100387	1086 (1.12-1.00)
Sidechain outliers	100360	1084 (1.12-1.00)
RSRZ outliers	91569	1080 (1.12-1.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	 2% 82% 12% • 5%
1	B	306	 5% 82% 11% • 6%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fluoroacetate dehalogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	291	Total	C	N	O	S	0	13	0
			2438	1572	428	425	13			
1	B	287	Total	C	N	O	S	0	8	0
			2365	1523	414	415	13			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP Q6NAM1
A	0	HIS	-	EXPRESSION TAG	UNP Q6NAM1
A	110	ASN	ASP	ENGINEERED MUTATION	UNP Q6NAM1
A	303	GLY	-	EXPRESSION TAG	UNP Q6NAM1
A	304	SER	-	EXPRESSION TAG	UNP Q6NAM1
B	-1	GLY	-	EXPRESSION TAG	UNP Q6NAM1
B	0	HIS	-	EXPRESSION TAG	UNP Q6NAM1
B	110	ASN	ASP	ENGINEERED MUTATION	UNP Q6NAM1
B	303	GLY	-	EXPRESSION TAG	UNP Q6NAM1
B	304	SER	-	EXPRESSION TAG	UNP Q6NAM1

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	4	Total	Cl	0	0
			4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	253	Total 253	O 253	0	0
4	B	144	Total 144	O 144	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fluoroacetate dehalogenase



- Molecule 1: Fluoroacetate dehalogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.93 Å 79.27 Å 85.08 Å 90.00° 103.22° 90.00°	Depositor
Resolution (Å)	41.41 – 1.05 41.41 – 1.05	Depositor EDS
% Data completeness (in resolution range)	96.8 (41.41-1.05) 96.8 (41.41-1.05)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 1.05 Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.144 , 0.168 0.152 , 0.174	Depositor DCC
$R_{free}$ test set	12237 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	9.5	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 49.8	EDS
Estimated twinning fraction	0.019 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 243419 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5206	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.22	12/2516 (0.5%)	1.22	18/3416 (0.5%)
1	B	1.29	18/2441 (0.7%)	1.15	7/3315 (0.2%)
All	All	1.25	30/4957 (0.6%)	1.18	25/6731 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
All	All	0	5

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	94	GLU	CD-OE1	-9.59	1.15	1.25
1	B	45	VAL	CB-CG2	-7.45	1.37	1.52
1	B	82	PRO	N-CA	7.25	1.59	1.47
1	B	79	GLN	C-O	7.17	1.36	1.23
1	A	50	VAL	CB-CG2	-7.12	1.38	1.52
1	B	50	VAL	CB-CG2	-7.08	1.38	1.52
1	A	269	SER	CB-OG	-6.90	1.33	1.42
1	B	56	GLU	CG-CD	6.72	1.62	1.51
1	B	149	TYR	CE1-CZ	6.32	1.46	1.38
1	B	149	TYR	CD1-CE1	6.18	1.48	1.39
1	B	141	TYR	CE2-CZ	-6.10	1.30	1.38
1	B	234	VAL	CB-CG2	-6.00	1.40	1.52
1	B	141	TYR	CB-CG	-5.77	1.43	1.51
1	B	74	PRO	CA-C	5.75	1.64	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	149	TYR	CB-CG	-5.68	1.43	1.51
1	B	141	TYR	CD2-CE2	-5.66	1.30	1.39
1	A	97[A]	GLU	CB-CG	-5.64	1.41	1.52
1	A	97[B]	GLU	CB-CG	-5.64	1.41	1.52
1	B	141	TYR	CG-CD1	-5.61	1.31	1.39
1	A	49[A]	ARG	CG-CD	-5.46	1.38	1.51
1	A	49[B]	ARG	CG-CD	-5.46	1.38	1.51
1	A	50	VAL	CA-CB	-5.42	1.43	1.54
1	B	144[A]	ARG	N-CA	5.31	1.56	1.46
1	B	144[B]	ARG	N-CA	5.31	1.56	1.46
1	A	143	GLN	CB-CG	-5.28	1.38	1.52
1	B	15	TRP	CB-CG	5.28	1.59	1.50
1	A	247	LEU	CB-CG	-5.17	1.37	1.52
1	B	141	TYR	CD1-CE1	-5.16	1.31	1.39
1	A	56	GLU	CD-OE1	-5.14	1.20	1.25
1	A	252	GLY	CA-C	5.11	1.60	1.51

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	265	ARG	NE-CZ-NH1	-9.29	115.65	120.30
1	A	244	MET	CG-SD-CE	-8.21	87.07	100.20
1	A	187	ARG	NE-CZ-NH1	-8.14	116.23	120.30
1	A	130	LEU	CB-CG-CD2	7.95	124.52	111.00
1	A	247	LEU	CA-CB-CG	7.77	133.16	115.30
1	A	77	ASP	CB-CG-OD2	7.11	124.70	118.30
1	B	220	ARG	NE-CZ-NH2	7.10	123.85	120.30
1	B	187	ARG	NE-CZ-NH2	-7.03	116.78	120.30
1	B	22	ARG	NE-CZ-NH1	-7.02	116.79	120.30
1	A	86	ARG	NE-CZ-NH2	6.81	123.70	120.30
1	A	58	PHE	CB-CG-CD2	-6.66	116.14	120.80
1	A	86	ARG	NE-CZ-NH1	-6.38	117.11	120.30
1	A	58	PHE	CB-CG-CD1	6.24	125.16	120.80
1	A	197	ARG	NE-CZ-NH2	-6.06	117.27	120.30
1	A	22	ARG	NE-CZ-NH2	5.88	123.24	120.30
1	A	220	ARG	NE-CZ-NH2	5.73	123.17	120.30
1	A	49[A]	ARG	NE-CZ-NH2	5.66	123.13	120.30
1	A	49[B]	ARG	NE-CZ-NH2	5.66	123.13	120.30
1	B	45	VAL	CG1-CB-CG2	5.44	119.61	110.90
1	A	154	TYR	CB-CG-CD2	5.36	124.22	121.00
1	A	147	ARG	NE-CZ-NH1	-5.36	117.62	120.30
1	A	126	ARG	NE-CZ-NH2	5.32	122.96	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	282	LEU	CB-CG-CD1	5.23	119.89	111.00
1	B	126	ARG	NE-CZ-NH1	-5.21	117.70	120.30
1	B	197	ARG	NE-CZ-NH2	-5.05	117.78	120.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	ARG	Sidechain
1	A	154	TYR	Sidechain
1	B	114	ARG	Sidechain
1	B	141	TYR	Sidechain
1	B	154	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2438	0	2361	39	0
1	B	2365	0	2283	15	0
2	A	1	0	0	0	0
3	A	4	0	0	0	0
3	B	1	0	0	0	0
4	A	253	0	0	4	0
4	B	144	0	0	2	0
All	All	5206	0	4644	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:TYR:CE1	1:A:153[B]:ILE:CD1	2.01	1.44
1:A:141:TYR:CE1	1:A:153[B]:ILE:HD13	1.55	1.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:TYR:CE1	1:A:153[B]:ILE:HG21	1.68	1.26
1:A:141:TYR:CE1	1:A:153[B]:ILE:HD12	1.80	1.07
1:A:141:TYR:HE1	1:A:153[B]:ILE:CD1	1.48	1.05
1:A:144[B]:ARG:O	1:A:149:TYR:CD1	2.13	1.02
1:A:168[B]:ASN:ND2	4:A:366:HOH:O	1.97	0.96
4:A:464:HOH:O	1:B:176[A]:PHE:CE2	2.20	0.95
1:A:141:TYR:CE1	1:A:153[B]:ILE:CG2	2.49	0.94
1:B:4:LEU:HD21	1:B:204:ILE:HD12	1.46	0.94
1:A:210[A]:MET:HE1	4:A:430:HOH:O	1.68	0.94
1:A:141:TYR:HE1	1:A:153[B]:ILE:HD13	0.96	0.88
1:A:141:TYR:CD1	1:A:153[B]:ILE:HD13	2.07	0.88
1:A:141:TYR:OH	1:A:153[B]:ILE:HB	1.77	0.83
4:A:464:HOH:O	1:B:176[A]:PHE:HE2	1.57	0.82
1:A:141:TYR:HE1	1:A:153[B]:ILE:CG2	1.86	0.80
1:A:153[B]:ILE:HG22	1:A:156:TRP:CD1	2.23	0.73
1:A:144[B]:ARG:O	1:A:149:TYR:HD1	1.64	0.72
1:A:141:TYR:CD1	1:A:153[B]:ILE:CD1	2.68	0.71
1:A:144[A]:ARG:HH21	1:A:144[A]:ARG:HG3	1.55	0.71
1:B:144[B]:ARG:NE	4:B:416:HOH:O	2.28	0.67
1:B:136:LEU:HB2	1:B:141:TYR:CE2	2.35	0.62
1:B:73:MET:HE1	1:B:210:MET:HB3	1.82	0.61
1:A:153[B]:ILE:CG2	1:A:156:TRP:CD1	2.84	0.60
1:A:141:TYR:CE1	1:A:153[B]:ILE:CB	2.84	0.59
1:A:141:TYR:HE1	1:A:153[B]:ILE:CB	2.15	0.59
1:B:232:ILE:HD12	4:B:383:HOH:O	2.03	0.57
1:B:136:LEU:HB2	1:B:141:TYR:HE2	1.70	0.55
1:A:152[A]:LYS:O	1:A:152[A]:LYS:HE2	2.07	0.54
1:A:141:TYR:CZ	1:A:153[B]:ILE:CG2	2.91	0.54
1:A:248:TRP:HB2	1:A:253:ILE:HG12	1.91	0.53
1:A:141:TYR:OH	1:A:153[B]:ILE:CB	2.53	0.53
1:A:141:TYR:HH	1:A:153[B]:ILE:HB	1.73	0.52
1:A:141:TYR:HE1	1:A:153[B]:ILE:HG21	1.14	0.52
1:A:263:VAL:HA	1:A:266:LYS:HE2	1.92	0.51
1:A:144[A]:ARG:NH2	1:A:144[A]:ARG:HG3	2.21	0.50
1:A:141:TYR:HE1	1:A:153[B]:ILE:CG1	2.16	0.49
1:A:139:TYR:CE2	1:A:143:GLN:HG3	2.47	0.49
1:A:46[B]:MET:SD	1:A:46[B]:MET:C	2.93	0.47
1:A:141:TYR:CD2	1:A:156:TRP:CE3	3.03	0.46
1:A:46[A]:MET:CE	1:A:283:PRO:HG2	2.46	0.46
1:B:51:ALA:N	1:B:52:PRO:CD	2.79	0.46
1:B:130:LEU:O	1:B:244[A]:MET:HA	2.15	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:277:GLU:HB2	1:B:289:GLN:HE22	1.81	0.45
1:B:46[C]:MET:C	1:B:46[C]:MET:SD	2.95	0.45
1:A:141:TYR:CZ	1:A:153[B]:ILE:HB	2.52	0.44
1:B:43:THR:O	1:B:46[B]:MET:HG3	2.19	0.42
1:A:153[B]:ILE:O	1:A:154:TYR:C	2.54	0.42
1:B:234:VAL:CG2	1:B:267:TRP:CZ2	3.03	0.42
1:A:141:TYR:OH	1:A:153[B]:ILE:CG2	2.68	0.42
1:A:141:TYR:CZ	1:A:153[B]:ILE:HG21	2.40	0.41
1:A:49[B]:ARG:NH2	1:A:193:ALA:O	2.45	0.41
1:A:144[A]:ARG:CG	1:A:144[A]:ARG:NH2	2.82	0.40
1:B:181:LYS:HD2	1:B:185:TRP:CZ2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/306 (98%)	293 (97%)	8 (3%)	0	100	100
1	B	292/306 (95%)	284 (97%)	8 (3%)	0	100	100
All	All	593/612 (97%)	577 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/242 (102%)	243 (98%)	4 (2%)	70	32
1	B	240/242 (99%)	239 (100%)	1 (0%)	93	75
All	All	487/484 (101%)	482 (99%)	5 (1%)	84	51

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	TRP
1	A	53	LYS
1	A	152[A]	LYS
1	A	152[B]	LYS
1	B	15	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	272	GLN
1	B	98	GLN
1	B	289	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	291/306 (95%)	-0.06	5 (1%) 73 65	5, 11, 21, 28	3 (1%)
1	B	287/306 (93%)	0.11	14 (4%) 33 28	8, 15, 28, 35	0
All	All	578/612 (94%)	0.03	19 (3%) 50 43	5, 12, 26, 35	3 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	5	ALA	5.9
1	B	10	GLY	4.6
1	A	300	ALA	4.5
1	B	78	GLU	3.8
1	B	4	LEU	3.5
1	B	185	TRP	3.3
1	A	253	ILE	3.0
1	B	9	PRO	2.9
1	B	75	GLU	2.7
1	B	77	ASP	2.7
1	B	261	LEU	2.6
1	B	149	TYR	2.4
1	A	144[A]	ARG	2.3
1	A	149	TYR	2.3
1	B	15	TRP	2.2
1	B	277	GLU	2.2
1	B	236	ALA	2.1
1	A	15	TRP	2.0
1	B	52	PRO	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	A	306	1/1	1.00	0.08	1.42	6,6,6,6	1
3	CL	A	307	1/1	1.00	0.06	-0.69	11,11,11,11	1
3	CL	B	305	1/1	1.00	0.04	-3.62	12,12,12,12	1
3	CL	A	309	1/1	0.99	0.09	-	13,13,13,13	1
2	CA	A	305	1/1	0.98	0.11	-	17,17,17,17	1
3	CL	A	308	1/1	1.00	0.03	-	15,15,15,15	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.