



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:17 PM GMT

PDB ID : 4RAP  
Title : Crystal structure of bacterial iron-containing dodecameric glycosyltransferase TibC from enterotoxigenic E.coli H10407  
Authors : Yao, Q.; Lu, Q.; Xu, Y.; Shao, F.  
Deposited on : 2014-09-10  
Resolution : 2.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1945 (2.90-2.86)
Clashscore	102246	2202 (2.90-2.86)
Ramachandran outliers	100387	2149 (2.90-2.86)
Sidechain outliers	100360	2152 (2.90-2.86)
RSRZ outliers	91569	1950 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	<div> <div>2%</div> <div>67% 23% 6%</div> </div>
1	B	406	<div> <div>2%</div> <div>70% 19% 5% 6%</div> </div>
1	C	406	<div> <div>2%</div> <div>70% 21% 5%</div> </div>
1	D	406	<div> <div>3%</div> <div>67% 20% 6% 6%</div> </div>
1	E	406	<div> <div>3%</div> <div>67% 24% 5%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	406	
1	G	406	
1	H	406	
1	I	406	
1	J	406	
1	K	406	
1	L	406	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FE	B	501	-	-	-	X
2	FE	C	501	-	-	-	X
2	FE	F	501	-	-	-	X
3	EDO	I	503	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 37131 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyltransferase TibC.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	381	Total	C	N	O	S	Se	0	0	0
			3069	1976	536	544	11	2			
1	B	383	Total	C	N	O	S	Se	0	0	0
			3085	1988	538	546	11	2			
1	C	389	Total	C	N	O	S	Se	0	0	0
			3124	2008	545	558	11	2			
1	D	381	Total	C	N	O	S	Se	0	0	0
			3067	1974	536	544	11	2			
1	E	390	Total	C	N	O	S	Se	0	0	0
			3132	2014	546	559	11	2			
1	F	382	Total	C	N	O	S	Se	0	0	0
			3071	1976	537	545	11	2			
1	G	389	Total	C	N	O	S	Se	0	0	0
			3124	2008	545	558	11	2			
1	H	385	Total	C	N	O	S	Se	0	0	0
			3096	1991	541	551	11	2			
1	I	389	Total	C	N	O	S	Se	0	0	0
			3124	2008	545	558	11	2			
1	J	384	Total	C	N	O	S	Se	0	0	0
			3092	1989	540	550	11	2			
1	K	389	Total	C	N	O	S	Se	0	0	0
			3124	2008	545	558	11	2			
1	L	371	Total	C	N	O	S	Se	0	0	0
			2988	1923	522	530	11	2			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
A	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
A	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
A	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
A	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
B	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
B	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
B	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
B	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
B	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
B	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
C	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
C	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
C	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
C	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
C	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
C	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
D	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
D	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
D	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
D	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
D	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
D	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
E	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
E	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
E	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
E	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
E	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
E	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
F	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
F	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
F	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
F	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
F	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
F	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
G	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
G	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
G	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
G	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
G	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
G	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
H	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
H	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
H	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
H	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
H	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
H	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
I	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
I	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
I	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
I	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
I	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
I	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
J	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
J	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
J	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
J	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
J	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
J	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
K	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
K	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
K	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
K	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
K	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
K	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
L	83	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
L	84	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
L	215	ALA	GLN	ENGINEERED MUTATION	UNP Q9S4K6
L	216	ALA	GLU	ENGINEERED MUTATION	UNP Q9S4K6
L	400	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6
L	401	ALA	LYS	ENGINEERED MUTATION	UNP Q9S4K6

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Fe 1 1	0	0
2	J	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0
2	K	1	Total Fe 1 1	0	0
2	E	1	Total Fe 1 1	0	0
2	H	1	Total Fe 1 1	0	0
2	B	1	Total Fe 1 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	I	1	Total	Fe	0	0
			1	1		
2	C	1	Total	Fe	0	0
			1	1		
2	A	1	Total	Fe	0	0
			1	1		
2	L	1	Total	Fe	0	0
			1	1		
2	F	1	Total	Fe	0	0
			1	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			4	2	2		
3	I	1	Total	C	O	0	0
			4	2	2		
3	I	1	Total	C	O	0	0
			4	2	2		
3	J	1	Total	C	O	0	0
			4	2	2		
3	L	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	O 1	0	0
4	C	1	Total 1	O 1	0	0
4	J	1	Total 1	O 1	0	0




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Glycosyltransferase TibC









• Molecule 1: Glycosyltransferase TibC




• Molecule 1: Glycosyltransferase TibC





• Molecule 1: Glycosyltransferase TibC







• Molecule 1: Glycosyltransferase TibC




• Molecule 1: Glycosyltransferase TibC



• Molecule 1: Glycosyltransferase TibC





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.79Å 314.37Å 164.53Å 90.00° 101.44° 90.00°	Depositor
Resolution (Å)	19.92 – 2.88 19.92 – 2.88	Depositor EDS
% Data completeness (in resolution range)	99.2 (19.92-2.88) 99.3 (19.92-2.88)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.74 (at 2.88Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.208 , 0.245 0.210 , 0.247	Depositor DCC
$R_{free}$ test set	9787 reflections (5.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	66.6	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 44.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	2 of 194251 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	37131	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FE, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.66	3/3164 (0.1%)	0.65	2/4310 (0.0%)
1	B	0.67	4/3182 (0.1%)	0.66	1/4336 (0.0%)
1	C	0.71	3/3220 (0.1%)	0.67	2/4388 (0.0%)
1	D	0.68	7/3162 (0.2%)	0.67	1/4307 (0.0%)
1	E	0.73	7/3228 (0.2%)	0.69	1/4399 (0.0%)
1	F	0.66	3/3164 (0.1%)	0.68	1/4305 (0.0%)
1	G	0.65	5/3220 (0.2%)	0.67	1/4388 (0.0%)
1	H	0.65	2/3192 (0.1%)	0.64	0/4349
1	I	0.66	5/3220 (0.2%)	0.64	1/4388 (0.0%)
1	J	0.70	5/3188 (0.2%)	0.68	1/4344 (0.0%)
1	K	0.74	6/3220 (0.2%)	0.74	8/4388 (0.2%)
1	L	0.69	3/3081 (0.1%)	0.68	2/4195 (0.0%)
All	All	0.68	53/38241 (0.1%)	0.67	21/52097 (0.0%)

The worst 5 of 53 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	K	29	ASN	C-O	-11.69	1.01	1.23
1	K	29	ASN	C-N	7.71	1.51	1.34
1	C	331	TRP	CD2-CE2	6.96	1.49	1.41
1	L	339	CYS	CB-SG	6.73	1.93	1.82
1	E	232	TRP	CD2-CE2	6.44	1.49	1.41

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	K	29	ASN	CA-C-O	12.00	145.29	120.10
1	K	29	ASN	O-C-N	-8.75	108.70	122.70
1	K	28	PHE	O-C-N	-8.01	109.89	122.70
1	C	339	CYS	CA-CB-SG	6.31	125.36	114.00
1	G	339	CYS	CA-CB-SG	6.30	125.34	114.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3069	0	2975	122	0
1	B	3085	0	2990	121	0
1	C	3124	0	3025	122	0
1	D	3067	0	2971	118	0
1	E	3132	0	3037	115	0
1	F	3071	0	2972	161	0
1	G	3124	0	3026	170	0
1	H	3096	0	2998	80	0
1	I	3124	0	3026	90	0
1	J	3092	0	2996	74	0
1	K	3124	0	3025	96	0
1	L	2988	0	2880	181	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	C	4	0	6	2	0
3	I	8	0	12	0	0
3	J	4	0	6	1	0
3	L	4	0	6	1	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	J	1	0	0	0	0
All	All	37131	0	35951	1447	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 1447 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:MSE:HG2	1:A:277:PHE:CE2	1.42	1.53
1:J:108:LEU:HD21	1:J:302:GLY:C	1.33	1.48
1:C:138:ILE:CD1	1:C:150:PHE:HB3	1.45	1.47
1:G:24:ILE:CG2	1:G:37:PRO:HD3	1.42	1.47
1:A:253:MSE:CG	1:A:277:PHE:CE2	2.03	1.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	377/406 (93%)	369 (98%)	8 (2%)	0	100	100
1	B	381/406 (94%)	368 (97%)	11 (3%)	2 (0%)	34	69
1	C	387/406 (95%)	373 (96%)	13 (3%)	1 (0%)	46	78
1	D	377/406 (93%)	363 (96%)	12 (3%)	2 (0%)	34	69
1	E	388/406 (96%)	369 (95%)	15 (4%)	4 (1%)	19	52
1	F	375/406 (92%)	358 (96%)	15 (4%)	2 (0%)	34	69
1	G	387/406 (95%)	370 (96%)	12 (3%)	5 (1%)	15	44
1	H	383/406 (94%)	370 (97%)	12 (3%)	1 (0%)	46	78
1	I	387/406 (95%)	373 (96%)	13 (3%)	1 (0%)	46	78
1	J	382/406 (94%)	371 (97%)	10 (3%)	1 (0%)	46	78
1	K	387/406 (95%)	376 (97%)	9 (2%)	2 (0%)	34	69
1	L	365/406 (90%)	345 (94%)	16 (4%)	4 (1%)	17	49
All	All	4576/4872 (94%)	4405 (96%)	146 (3%)	25 (0%)	34	69

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	21	PRO
1	E	23	ASN
1	E	395	VAL
1	G	33	ARG
1	I	397	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/347 (95%)	300 (91%)	31 (9%)	11	30
1	B	332/347 (96%)	308 (93%)	24 (7%)	18	44
1	C	336/347 (97%)	315 (94%)	21 (6%)	22	52
1	D	330/347 (95%)	299 (91%)	31 (9%)	11	30
1	E	337/347 (97%)	311 (92%)	26 (8%)	16	40
1	F	330/347 (95%)	282 (86%)	48 (14%)	4	10
1	G	336/347 (97%)	315 (94%)	21 (6%)	22	52
1	H	333/347 (96%)	311 (93%)	22 (7%)	21	49
1	I	336/347 (97%)	312 (93%)	24 (7%)	18	45
1	J	333/347 (96%)	312 (94%)	21 (6%)	22	52
1	K	336/347 (97%)	312 (93%)	24 (7%)	18	45
1	L	321/347 (92%)	290 (90%)	31 (10%)	10	28
All	All	3991/4164 (96%)	3667 (92%)	324 (8%)	15	38

5 of 324 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	58	ASP
1	G	22	GLU
1	L	65	THR
1	F	81	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	155	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	367	GLN
1	F	149	GLN
1	L	29	ASN
1	F	18	GLN
1	F	29	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 12 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	EDO	C	502	-	3,3,3	0.31	0	2,2,2	0.84	0
3	EDO	I	502	-	3,3,3	0.32	0	2,2,2	0.67	0
3	EDO	I	503	-	3,3,3	0.60	0	2,2,2	0.51	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	J	502	-	3,3,3	0.34	0	2,2,2	0.72	0
3	EDO	L	502	-	3,3,3	0.28	0	2,2,2	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	C	502	-	-	0/1/1/1	0/0/0/0
3	EDO	I	502	-	-	0/1/1/1	0/0/0/0
3	EDO	I	503	-	-	0/1/1/1	0/0/0/0
3	EDO	J	502	-	-	0/1/1/1	0/0/0/0
3	EDO	L	502	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	502	EDO	2	0
3	J	502	EDO	1	0
3	L	502	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	379/406 (93%)	-0.21	7 (1%) 71 69	44, 69, 108, 132	0
1	B	381/406 (93%)	-0.22	10 (2%) 59 55	42, 64, 114, 149	0
1	C	387/406 (95%)	-0.27	10 (2%) 59 55	34, 56, 114, 146	0
1	D	379/406 (93%)	-0.14	11 (2%) 55 50	25, 66, 108, 129	0
1	E	388/406 (95%)	-0.22	11 (2%) 56 51	26, 59, 109, 131	0
1	F	380/406 (93%)	-0.03	21 (5%) 29 23	25, 66, 125, 151	0
1	G	387/406 (95%)	-0.06	26 (6%) 21 15	30, 65, 114, 150	0
1	H	383/406 (94%)	-0.19	13 (3%) 49 42	44, 66, 106, 128	0
1	I	387/406 (95%)	-0.23	11 (2%) 56 51	40, 65, 108, 125	0
1	J	382/406 (94%)	-0.37	6 (1%) 74 73	35, 59, 97, 117	0
1	K	387/406 (95%)	-0.28	6 (1%) 74 73	37, 57, 98, 124	0
1	L	369/406 (90%)	-0.13	8 (2%) 65 62	37, 67, 116, 132	0
All	All	4589/4872 (94%)	-0.20	140 (3%) 52 46	25, 64, 112, 151	0

The worst 5 of 140 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	89	ASP	7.6
1	G	21	PRO	5.8
1	G	395	VAL	5.6
1	C	159	VAL	5.1
1	L	395	VAL	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	I	503	4/4	0.89	0.37	6.73	61,75,76,77	0
2	FE	F	501	1/1	0.96	0.45	3.42	55,55,55,55	0
2	FE	B	501	1/1	0.95	0.44	3.40	60,60,60,60	0
2	FE	C	501	1/1	0.95	0.46	3.18	54,54,54,54	0
2	FE	H	501	1/1	0.97	0.48	-	60,60,60,60	0
2	FE	E	1400	1/1	0.96	0.44	-	57,57,57,57	0
3	EDO	C	502	4/4	0.95	0.18	-	40,43,48,59	0
2	FE	K	501	1/1	0.82	0.44	-	59,59,59,59	0
3	EDO	J	502	4/4	0.97	0.17	-	48,49,52,57	0
2	FE	I	501	1/1	0.97	0.46	-	58,58,58,58	0
2	FE	D	501	1/1	0.97	0.47	-	63,63,63,63	0
3	EDO	I	502	4/4	0.92	0.22	-	57,59,72,80	0
2	FE	A	501	1/1	0.98	0.43	-	59,59,59,59	0
3	EDO	L	502	4/4	0.97	0.15	-	51,52,59,62	0
2	FE	G	1400	1/1	0.96	0.47	-	57,57,57,57	0
2	FE	L	501	1/1	0.98	0.46	-	55,55,55,55	0
2	FE	J	501	1/1	0.96	0.48	-	56,56,56,56	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.