



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:17 PM GMT

PDB ID : 4RB4
Title : Crystal structure of dodecameric iron-containing heptosyltransferase TibC in complex with ADP-D-beta-D-heptose at 3.9 angstrom resolution
Authors : Yao, Q.; Lu, Q.; Shao, F.
Deposited on : 2014-09-12
Resolution : 3.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1005 (4.24-3.52)
Clashscore	102246	1026 (4.20-3.56)
Ramachandran outliers	100387	1003 (4.22-3.54)
Sidechain outliers	100360	1043 (4.24-3.52)
RSRZ outliers	91569	1009 (4.24-3.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	<div> <div></div> <div>72% 17% 6%</div> </div>
1	B	406	<div> <div></div> <div>74% 16% 5% 6%</div> </div>
1	C	406	<div> <div></div> <div>73% 19% . .</div> </div>
1	D	406	<div> <div></div> <div>69% 21% . 6%</div> </div>
1	E	406	<div> <div></div> <div>67% 24% 5% .</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	406	
1	G	406	
1	H	406	
1	I	406	
1	J	406	
1	K	406	
1	L	406	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FE	A	501	-	-	X	-
2	FE	D	501	-	-	X	-
2	FE	J	501	-	-	X	-
3	AQH	A	502	-	-	-	X
3	AQH	D	502	-	-	X	-
3	AQH	H	502	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 37611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyltransferase tibC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	K	390	Total	C	N	O	S	0	0	0
			3132	2016	546	557	13			
1	L	374	Total	C	N	O	S	0	0	0
			3008	1938	524	533	13			
1	C	389	Total	C	N	O	S	0	0	0
			3121	2007	545	556	13			
1	D	381	Total	C	N	O	S	0	0	0
			3061	1970	536	542	13			
1	E	390	Total	C	N	O	S	0	0	0
			3129	2013	546	557	13			
1	H	385	Total	C	N	O	S	0	0	0
			3093	1990	541	549	13			
1	B	383	Total	C	N	O	S	0	0	0
			3082	1987	538	544	13			
1	F	384	Total	C	N	O	S	0	0	0
			3084	1985	539	547	13			
1	G	389	Total	C	N	O	S	0	0	0
			3121	2007	545	556	13			
1	I	389	Total	C	N	O	S	0	0	0
			3121	2007	545	556	13			
1	J	385	Total	C	N	O	S	0	0	0
			3097	1996	541	547	13			
1	A	380	Total	C	N	O	S	0	0	0
			3058	1969	535	541	13			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
K	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
L	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
C	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
D	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
E	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
B	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
F	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
G	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
I	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
J	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13
A	110	ALA	ASP	ENGINEERED MUTATION	UNP H5MH13


- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Fe 1 1	0	0
2	J	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0
2	K	1	Total Fe 1 1	0	0
2	E	1	Total Fe 1 1	0	0
2	H	1	Total Fe 1 1	0	0
2	B	1	Total Fe 1 1	0	0
2	I	1	Total Fe 1 1	0	0
2	C	1	Total Fe 1 1	0	0
2	A	1	Total Fe 1 1	0	0
2	L	1	Total Fe 1 1	0	0
2	F	1	Total Fe 1 1	0	0

- Molecule 3 is [(2R,3S,4R,5R)-5-(6-AMINO-9H-PURIN-9-YL)-3,4-DIHYDROXYTETRAHYDROFURAN-2-YL]METHYL (2R,3R,4R,5R,6S)-6-[(1R)-1,2-DIHYDROXYETHYL]-3,4,5-TRIHYDROXYTETRAHYDRO-2H-PYRAN-2-YL DIHYDROGEN DIPHOSPHATE (three-letter code: AQH) (formula: C₁₇H₂₇N₅O₁₆P₂).



- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\text{C}_2\text{H}_6\text{O}_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			4	2	2		
4	I	1	Total	C	O	0	0
			4	2	2		
4	J	1	Total	C	O	0	0
			4	2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Glycosyltransferase tibC



• Molecule 1: Glycosyltransferase tibC




• Molecule 1: Glycosyltransferase tibC




- Molecule 1: Glycosyltransferase tibC

Chain B: 




- Molecule 1: Glycosyltransferase tibC

Chain F: 



- Molecule 1: Glycosyltransferase tibC


Chain G: 





- Molecule 1: Glycosyltransferase tibC

Chain I: 








• Molecule 1: Glycosyltransferase tibC



• Molecule 1: Glycosyltransferase tibC



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	88.29Å 313.20Å 164.69Å 90.00° 101.26° 90.00°	Depositor
Resolution (Å)	20.04 – 3.88 20.03 – 3.88	Depositor EDS
% Data completeness (in resolution range)	98.7 (20.04-3.88) 99.4 (20.03-3.88)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.74 (at 3.94Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.282 , 0.295 0.293 , 0.300	Depositor DCC
R_{free} test set	3995 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	98.2	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.22 , 59.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 79848 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	37611	wwPDB-VP
Average B, all atoms (Å ²)	149.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: AQH, EDO, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	3/3154 (0.1%)	0.34	0/4298
1	B	0.65	4/3181 (0.1%)	0.34	0/4338
1	C	0.58	0/3219	0.35	0/4390
1	D	0.59	3/3158 (0.1%)	0.34	0/4305
1	E	0.58	3/3227 (0.1%)	0.35	0/4401
1	F	0.65	3/3181 (0.1%)	0.34	0/4336
1	G	0.64	4/3219 (0.1%)	0.36	0/4390
1	H	0.60	3/3191 (0.1%)	0.37	1/4351 (0.0%)
1	I	0.61	6/3219 (0.2%)	0.37	1/4390 (0.0%)
1	J	0.62	3/3196 (0.1%)	0.37	1/4358 (0.0%)
1	K	0.58	2/3231 (0.1%)	0.36	0/4406
1	L	0.62	2/3102 (0.1%)	0.37	0/4225
All	All	0.61	36/38278 (0.1%)	0.35	3/52188 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	3
1	D	0	1
1	E	0	3
1	F	0	4
1	G	0	2
1	H	0	2
1	I	0	1
1	J	0	2
1	K	0	2

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	2
All	All	0	26

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	97	ARG	CZ-NH1	-8.61	1.21	1.33
1	H	265	PHE	CG-CD1	-8.42	1.26	1.38
1	L	291	ARG	CZ-NH1	-8.40	1.22	1.33
1	E	291	ARG	CZ-NH1	-7.90	1.22	1.33
1	A	82	GLY	CA-C	7.82	1.64	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	109	GLY	N-CA-C	6.28	128.81	113.10
1	I	103	PHE	C-N-CD	5.59	140.14	128.40
1	J	105	THR	N-CA-C	-5.05	97.36	111.00

There are no chirality outliers.

5 of 26 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	79	PHE	Peptide
1	K	251	ARG	Sidechain
1	K	357	TRP	Peptide
1	L	79	PHE	Peptide
1	L	80	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3058	0	2956	95	0
1	B	3082	0	2979	87	0
1	C	3121	0	3016	98	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3061	0	2952	137	0
1	E	3129	0	3027	134	0
1	F	3084	0	2979	96	0
1	G	3121	0	3016	128	0
1	H	3093	0	2988	116	0
1	I	3121	0	3015	120	0
1	J	3097	0	2993	100	0
1	K	3132	0	3024	118	0
1	L	3008	0	2882	268	0
2	A	1	0	0	2	0
2	B	1	0	0	1	0
2	C	1	0	0	1	0
2	D	1	0	0	3	0
2	E	1	0	0	1	0
2	F	1	0	0	1	0
2	G	1	0	0	1	0
2	H	1	0	0	1	0
2	I	1	0	0	1	0
2	J	1	0	0	3	0
2	K	1	0	0	1	0
2	L	1	0	0	1	0
3	A	40	0	25	13	0
3	B	40	0	25	8	0
3	C	40	0	25	12	0
3	D	40	0	25	23	0
3	E	40	0	25	10	0
3	F	40	0	25	19	0
3	G	40	0	25	12	0
3	H	40	0	25	24	0
3	I	40	0	25	11	0
3	J	40	0	25	8	0
3	K	40	0	25	16	0
3	L	40	0	25	17	0
4	C	4	0	6	1	0
4	I	4	0	6	1	0
4	J	4	0	6	0	0
All	All	37611	0	36145	1508	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 1508 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:296:PHE:CZ	1:L:298:GLY:HA3	1.21	1.68
1:L:107:THR:HG22	1:L:108:LEU:CD2	1.19	1.61
1:L:296:PHE:CE2	1:L:298:GLY:HA3	1.41	1.51
1:D:110:ALA:CB	1:D:137:ILE:HD12	1.07	1.50
1:D:110:ALA:HB1	1:D:137:ILE:CG2	1.43	1.49

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/406 (92%)	360 (96%)	10 (3%)	4 (1%)	17	63
1	B	381/406 (94%)	364 (96%)	12 (3%)	5 (1%)	15	60
1	C	387/406 (95%)	367 (95%)	15 (4%)	5 (1%)	15	60
1	D	377/406 (93%)	356 (94%)	15 (4%)	6 (2%)	12	56
1	E	388/406 (96%)	366 (94%)	18 (5%)	4 (1%)	19	64
1	F	380/406 (94%)	355 (93%)	20 (5%)	5 (1%)	15	60
1	G	387/406 (95%)	365 (94%)	16 (4%)	6 (2%)	12	56
1	H	383/406 (94%)	370 (97%)	7 (2%)	6 (2%)	12	56
1	I	387/406 (95%)	364 (94%)	21 (5%)	2 (0%)	34	76
1	J	383/406 (94%)	368 (96%)	11 (3%)	4 (1%)	19	64
1	K	388/406 (96%)	377 (97%)	9 (2%)	2 (0%)	34	76
1	L	364/406 (90%)	338 (93%)	14 (4%)	12 (3%)	5	43
All	All	4579/4872 (94%)	4350 (95%)	168 (4%)	61 (1%)	15	60

5 of 61 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	10	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	231	TYR
1	L	233	ASN
1	L	234	ASN
1	L	299	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/355 (93%)	311 (94%)	18 (6%)	27	66
1	B	331/355 (93%)	313 (95%)	18 (5%)	27	66
1	C	335/355 (94%)	320 (96%)	15 (4%)	34	70
1	D	328/355 (92%)	311 (95%)	17 (5%)	29	67
1	E	336/355 (95%)	315 (94%)	21 (6%)	22	61
1	F	331/355 (93%)	316 (96%)	15 (4%)	34	70
1	G	335/355 (94%)	308 (92%)	27 (8%)	15	52
1	H	332/355 (94%)	316 (95%)	16 (5%)	31	69
1	I	335/355 (94%)	312 (93%)	23 (7%)	19	58
1	J	332/355 (94%)	315 (95%)	17 (5%)	29	67
1	K	336/355 (95%)	313 (93%)	23 (7%)	20	59
1	L	322/355 (91%)	295 (92%)	27 (8%)	14	51
All	All	3982/4260 (94%)	3745 (94%)	237 (6%)	24	63

5 of 237 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	107	THR
1	B	205	ARG
1	A	10	ILE
1	H	170	TYR
1	B	40	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	284	GLN
1	G	367	GLN
1	I	367	GLN
1	A	337	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 12 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	AQH	A	502	-	35,43,43	1.83	10 (28%)	45,66,66	3.01	12 (26%)
3	AQH	B	502	-	35,43,43	1.84	10 (28%)	45,66,66	3.11	12 (26%)
4	EDO	C	502	-	3,3,3	0.32	0	2,2,2	0.40	0
3	AQH	C	503	-	35,43,43	1.83	10 (28%)	45,66,66	3.03	12 (26%)
3	AQH	D	502	-	35,43,43	1.81	10 (28%)	45,66,66	3.08	13 (28%)
3	AQH	E	502	-	35,43,43	1.82	10 (28%)	45,66,66	3.02	12 (26%)
3	AQH	F	502	-	35,43,43	1.89	10 (28%)	45,66,66	3.15	15 (33%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AQH	G	502	-	35,43,43	1.87	9 (25%)	45,66,66	3.15	14 (31%)
3	AQH	H	502	-	35,43,43	1.82	9 (25%)	45,66,66	3.02	12 (26%)
4	EDO	I	502	-	3,3,3	0.24	0	2,2,2	0.40	0
3	AQH	I	503	-	35,43,43	1.83	9 (25%)	45,66,66	3.02	12 (26%)
4	EDO	J	502	-	3,3,3	0.29	0	2,2,2	0.40	0
3	AQH	J	503	-	35,43,43	1.80	10 (28%)	45,66,66	3.11	12 (26%)
3	AQH	K	502	-	35,43,43	1.81	9 (25%)	45,66,66	3.11	12 (26%)
3	AQH	L	502	-	35,43,43	1.88	10 (28%)	45,66,66	3.14	15 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AQH	A	502	-	-	0/23/63/63	0/4/4/4
3	AQH	B	502	-	-	0/23/63/63	0/4/4/4
4	EDO	C	502	-	-	0/1/1/1	0/0/0/0
3	AQH	C	503	-	-	0/23/63/63	0/4/4/4
3	AQH	D	502	-	-	0/23/63/63	0/4/4/4
3	AQH	E	502	-	-	0/23/63/63	0/4/4/4
3	AQH	F	502	-	-	0/23/63/63	0/4/4/4
3	AQH	G	502	-	-	0/23/63/63	0/4/4/4
3	AQH	H	502	-	-	0/23/63/63	0/4/4/4
4	EDO	I	502	-	-	0/1/1/1	0/0/0/0
3	AQH	I	503	-	-	0/23/63/63	0/4/4/4
4	EDO	J	502	-	-	0/1/1/1	0/0/0/0
3	AQH	J	503	-	-	0/23/63/63	0/4/4/4
3	AQH	K	502	-	-	0/23/63/63	0/4/4/4
3	AQH	L	502	-	-	0/23/63/63	0/4/4/4

The worst 5 of 116 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	J	503	AQH	PA-O1A	-3.15	1.41	1.54
3	G	502	AQH	PA-O1A	-3.14	1.41	1.54
3	C	503	AQH	PA-O1A	-3.13	1.41	1.54
3	K	502	AQH	PA-O1A	-3.13	1.41	1.54
3	E	502	AQH	PA-O1A	-3.12	1.41	1.54

The worst 5 of 153 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	503	AQH	N3-C2-N1	-14.64	117.68	128.89
3	F	502	AQH	N3-C2-N1	-14.63	117.69	128.89
3	K	502	AQH	N3-C2-N1	-14.61	117.71	128.89
3	G	502	AQH	N3-C2-N1	-14.59	117.72	128.89
3	E	502	AQH	N3-C2-N1	-14.59	117.72	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

14 monomers are involved in 175 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	AQH	13	0
3	B	502	AQH	8	0
4	C	502	EDO	1	0
3	C	503	AQH	12	0
3	D	502	AQH	23	0
3	E	502	AQH	10	0
3	F	502	AQH	19	0
3	G	502	AQH	12	0
3	H	502	AQH	24	0
4	I	502	EDO	1	0
3	I	503	AQH	11	0
3	J	503	AQH	8	0
3	K	502	AQH	16	0
3	L	502	AQH	17	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	380/406 (93%)	-0.07	3 (0%) 87 81	150, 150, 150, 150	0
1	B	383/406 (94%)	-0.15	2 (0%) 91 88	150, 150, 150, 150	0
1	C	389/406 (95%)	-0.11	2 (0%) 91 88	150, 150, 150, 150	0
1	D	381/406 (93%)	-0.09	6 (1%) 74 64	30, 150, 150, 150	0
1	E	390/406 (96%)	-0.07	4 (1%) 84 77	124, 150, 150, 150	0
1	F	384/406 (94%)	-0.04	6 (1%) 74 64	150, 150, 150, 150	0
1	G	389/406 (95%)	-0.05	7 (1%) 71 61	150, 150, 150, 150	0
1	H	385/406 (94%)	-0.07	8 (2%) 67 56	25, 150, 150, 150	0
1	I	389/406 (95%)	-0.07	6 (1%) 76 66	85, 150, 150, 150	0
1	J	385/406 (94%)	-0.16	3 (0%) 87 81	150, 150, 150, 150	0
1	K	390/406 (96%)	-0.11	3 (0%) 87 81	150, 150, 150, 150	0
1	L	374/406 (92%)	0.02	8 (2%) 67 56	23, 150, 150, 150	0
All	All	4619/4872 (94%)	-0.08	58 (1%) 79 70	23, 150, 150, 150	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	57	CYS	6.4
1	D	110	ALA	5.7
1	A	85	THR	3.9
1	L	225	SER	3.7
1	F	160	ALA	3.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	AQH	A	502	40/40	0.76	0.36	2.08	150,150,150,150	40
3	AQH	I	503	40/40	0.81	0.34	0.87	150,150,150,150	0
3	AQH	L	502	40/40	0.75	0.40	0.73	150,150,150,150	0
3	AQH	F	502	40/40	0.84	0.28	0.73	150,150,150,150	0
3	AQH	G	502	40/40	0.87	0.27	0.69	150,150,150,150	40
3	AQH	H	502	40/40	0.80	0.33	0.59	150,150,150,150	0
3	AQH	J	503	40/40	0.88	0.26	0.31	150,150,150,150	0
3	AQH	B	502	40/40	0.88	0.27	0.29	150,150,150,150	0
3	AQH	D	502	40/40	0.83	0.28	0.12	150,150,150,150	40
3	AQH	C	503	40/40	0.87	0.25	0.01	150,150,150,150	0
3	AQH	K	502	40/40	0.89	0.23	-0.34	150,150,150,150	0
3	AQH	E	502	40/40	0.85	0.24	-0.37	150,150,150,150	0
2	FE	B	501	1/1	0.98	0.43	-	150,150,150,150	0
4	EDO	J	502	4/4	0.85	0.36	-	150,150,150,150	0
4	EDO	I	502	4/4	0.82	0.33	-	150,150,150,150	0
2	FE	J	501	1/1	0.98	0.48	-	150,150,150,150	0
2	FE	A	501	1/1	0.99	0.46	-	150,150,150,150	0
4	EDO	C	502	4/4	0.78	0.33	-	150,150,150,150	0
2	FE	I	501	1/1	0.94	0.52	-	150,150,150,150	0
2	FE	C	501	1/1	0.94	0.53	-	150,150,150,150	0
2	FE	D	501	1/1	0.98	0.47	-	150,150,150,150	0
2	FE	H	501	1/1	0.99	0.45	-	150,150,150,150	0
2	FE	E	501	1/1	0.91	0.52	-	150,150,150,150	0
2	FE	F	501	1/1	0.99	0.46	-	150,150,150,150	0
2	FE	L	501	1/1	0.99	0.46	-	150,150,150,150	0
2	FE	G	501	1/1	0.71	0.58	-	150,150,150,150	0
2	FE	K	501	1/1	0.91	0.66	-	150,150,150,150	0

6.5 Other polymers [i](#)

There are no such residues in this entry.