



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:36 PM GMT

PDB ID : 3RGN  
Title : Crystal structure of spin-labeled BtuB W371R1  
Authors : Freed, D.M.; Horanyi, P.S.; Wiener, M.C.; Cafiso, D.S.  
Deposited on : 2011-04-08  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	594	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	C8E	A	801	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	C8E	A	802	-	-	-	X
2	C8E	A	803	-	-	-	X
2	C8E	A	805	-	-	-	X
2	C8E	A	806	-	-	-	X
3	MG	A	596	-	-	-	X
4	MTN	A	598	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Vitamin B12 transporter BtuB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	549	Total	C	N	O	S	0	0	0
			4320	2720	743	854	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	371	CYS	TRP	ENGINEERED MUTATION	UNP P06129

- Molecule 2 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (three-letter code: C8E) (formula:  $C_{16}H_{34}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Mg	0	0
			3	3		

- Molecule 4 is S-[(1-OXYL-2,2,5,5-TETRAMETHYL-2,5-DIHYDRO-1H-PYRROL-3-YL)METHYL] METHANESULFONOTHIOATE (three-letter code: MTN) (formula: C<sub>10</sub>H<sub>18</sub>NO<sub>3</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	9	1	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	130	Total 130	O 130	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vitamin B12 transporter BtuB



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.68Å 81.68Å 227.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.38 – 2.30 33.38 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.2 (33.38-2.30) 95.1 (33.38-2.30)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.17 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.220 , 0.250 0.218 , 0.249	Depositor DCC
$R_{free}$ test set	1911 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	57.6	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 47.8	EDS
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 37884 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4612	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MTN, C8E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.12	6/4422 (0.1%)	1.01	15/6015 (0.2%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	372	GLN	CB-CG	-9.17	1.27	1.52
1	A	354	GLU	CD-OE2	7.57	1.33	1.25
1	A	354	GLU	CG-CD	5.95	1.60	1.51
1	A	320	GLN	CD-OE1	5.56	1.36	1.24
1	A	423	GLU	CB-CG	-5.35	1.42	1.52
1	A	381	GLU	CB-CG	5.16	1.61	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	384	ARG	NE-CZ-NH2	-9.90	115.35	120.30
1	A	121	ASP	CB-CG-OD2	-8.18	110.94	118.30
1	A	121	ASP	CB-CG-OD1	7.80	125.32	118.30
1	A	487	ARG	NE-CZ-NH2	-6.89	116.85	120.30
1	A	498	ARG	NE-CZ-NH1	6.89	123.74	120.30
1	A	384	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	A	460	ARG	NE-CZ-NH2	-6.78	116.91	120.30
1	A	47	ARG	NE-CZ-NH1	6.63	123.62	120.30
1	A	498	ARG	NE-CZ-NH2	-6.42	117.09	120.30
1	A	456	GLU	C-N-CA	-5.99	109.73	122.30
1	A	439	ASP	CB-CG-OD2	5.56	123.30	118.30
1	A	33	ASP	CB-CG-OD2	-5.38	113.46	118.30
1	A	6	ASP	CB-CG-OD1	-5.22	113.60	118.30
1	A	106	ARG	NE-CZ-NH2	-5.04	117.78	120.30
1	A	47	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4320	0	4073	41	0
2	A	147	0	238	11	0
3	A	3	0	0	0	0
4	A	12	0	15	1	0
5	A	130	0	0	1	0
All	All	4612	0	4326	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:802:C8E:H13	2:A:802:C8E:H82	1.51	0.93
2:A:802:C8E:H13	2:A:802:C8E:H52	1.50	0.92
1:A:166:VAL:HG22	1:A:206:LEU:CD1	2.01	0.91
1:A:544:VAL:HG23	1:A:571:LYS:HE2	1.57	0.85
1:A:529:LYS:NZ	1:A:538:THR:HG22	1.91	0.84
1:A:166:VAL:HG22	1:A:206:LEU:HD12	1.58	0.84
2:A:802:C8E:C1	2:A:802:C8E:H82	2.17	0.73
1:A:529:LYS:HZ3	1:A:538:THR:HG22	1.58	0.67
1:A:544:VAL:CG2	1:A:571:LYS:HE2	2.26	0.66
2:A:802:C8E:C1	2:A:802:C8E:H52	2.26	0.58
2:A:806:C8E:H61	4:A:598:MTN:H71	1.85	0.58
1:A:527:TYR:CE1	1:A:540:LYS:HG3	2.39	0.57
1:A:422:PHE:CG	2:A:801:C8E:H11	2.39	0.57
1:A:169:LEU:HD23	1:A:169:LEU:C	2.25	0.56
1:A:299:GLN:O	1:A:299:GLN:HG3	2.05	0.56
1:A:329:VAL:CG1	1:A:333:TYR:HE2	2.19	0.55
1:A:305:ILE:HD13	1:A:305:ILE:N	2.22	0.53
2:A:802:C8E:C8	2:A:802:C8E:H13	2.32	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:587:THR:HG22	1:A:588:LEU:N	2.24	0.52
1:A:114:ARG:HA	1:A:372:GLN:OE1	2.12	0.50
1:A:267:THR:HG23	1:A:298:VAL:HG22	1.92	0.49
2:A:802:C8E:C5	2:A:802:C8E:H13	2.33	0.49
1:A:529:LYS:HZ1	1:A:538:THR:HG22	1.77	0.48
1:A:116:ALA:O	1:A:358:ARG:HB2	2.14	0.48
1:A:329:VAL:HG11	1:A:333:TYR:CE2	2.49	0.47
1:A:19:ARG:HE	1:A:26:THR:CG2	2.27	0.47
1:A:323:THR:HG23	1:A:324:PRO:HD2	1.95	0.47
1:A:162:ASP:O	1:A:163:LYS:HB2	2.15	0.47
2:A:802:C8E:H171	2:A:802:C8E:H11	1.98	0.46
1:A:222:GLY:HA2	1:A:247:SER:O	2.16	0.46
1:A:98:GLN:HE22	1:A:227:THR:HG21	1.80	0.46
1:A:100:PRO:HG2	1:A:103:LEU:HD12	1.98	0.46
1:A:166:VAL:HG22	1:A:206:LEU:HD11	1.95	0.45
1:A:142:ALA:HA	1:A:151:ASN:O	2.17	0.45
1:A:329:VAL:HG11	1:A:333:TYR:HE2	1.82	0.45
2:A:802:C8E:C1	2:A:802:C8E:H171	2.47	0.44
1:A:529:LYS:CE	1:A:538:THR:HG22	2.47	0.44
1:A:227:THR:HG23	1:A:243:ARG:HB3	2.00	0.44
1:A:78:VAL:HG11	1:A:96:LEU:HD21	2.00	0.44
2:A:803:C8E:H22	2:A:805:C8E:H132	1.99	0.44
1:A:44:ASP:HA	1:A:47:ARG:HG2	1.99	0.44
1:A:529:LYS:NZ	1:A:538:THR:CG2	2.72	0.44
1:A:382:GLY:O	1:A:424:GLY:HA2	2.19	0.43
1:A:27:THR:HG21	1:A:49:LEU:HD22	2.00	0.43
1:A:19:ARG:HE	1:A:26:THR:HG23	1.84	0.42
1:A:169:LEU:HD23	1:A:170:GLY:N	2.34	0.42
1:A:436:TYR:CZ	1:A:463:GLY:HA3	2.54	0.42
1:A:74:SER:HB2	1:A:394:LYS:HE2	2.02	0.41
1:A:460:ARG:HG3	5:A:725:HOH:O	2.22	0.40
1:A:587:THR:CG2	1:A:588:LEU:N	2.83	0.40
1:A:19:ARG:NH2	1:A:26:THR:O	2.53	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	541/594 (91%)	522 (96%)	17 (3%)	2 (0%)	39	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	121	ASP
1	A	87	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	453/495 (92%)	438 (97%)	15 (3%)	45	61

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	LEU
1	A	47	ARG
1	A	93	SER
1	A	169	LEU
1	A	175	THR
1	A	202	LEU
1	A	227	THR
1	A	228	ASN
1	A	270	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	328	TYR
1	A	346	GLN
1	A	538	THR
1	A	540	LYS
1	A	557	SER
1	A	580	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	MTN	A	598	1	10,12,16	1.88	5 (50%)	13,20,27	4.30	5 (38%)
2	C8E	A	800	-	20,20,20	0.34	0	19,19,19	0.65	0
2	C8E	A	801	-	20,20,20	0.49	0	19,19,19	0.88	0
2	C8E	A	802	-	20,20,20	0.55	0	19,19,19	1.01	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	C8E	A	803	-	20,20,20	0.46	0	19,19,19	0.52	0
2	C8E	A	804	-	20,20,20	0.42	0	19,19,19	0.69	0
2	C8E	A	805	-	20,20,20	0.63	0	19,19,19	0.86	1 (5%)
2	C8E	A	806	-	20,20,20	0.54	0	19,19,19	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MTN	A	598	1	-	0/0/25/29	0/1/1/1
2	C8E	A	800	-	-	0/18/18/18	0/0/0/0
2	C8E	A	801	-	-	0/18/18/18	0/0/0/0
2	C8E	A	802	-	-	0/18/18/18	0/0/0/0
2	C8E	A	803	-	-	0/18/18/18	0/0/0/0
2	C8E	A	804	-	-	0/18/18/18	0/0/0/0
2	C8E	A	805	-	-	0/18/18/18	0/0/0/0
2	C8E	A	806	-	-	0/18/18/18	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	598	MTN	C5-N1	-3.11	1.46	1.50
4	A	598	MTN	C4-S1	2.00	1.86	1.81
4	A	598	MTN	C4-C3	2.16	1.55	1.50
4	A	598	MTN	C2-C3	2.54	1.35	1.32
4	A	598	MTN	C1-C2	2.82	1.54	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	598	MTN	C7-C5-C3	-4.98	106.97	113.23
2	A	802	C8E	C10-O9-C8	-2.23	103.70	113.31
2	A	805	C8E	O15-C16-C17	2.15	119.93	110.36
4	A	598	MTN	C3-C4-S1	2.30	117.76	110.44
4	A	598	MTN	O1-N1-C5	3.61	124.95	110.84
4	A	598	MTN	C2-C1-N1	4.50	101.97	99.36
4	A	598	MTN	C3-C5-N1	12.90	105.23	99.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	598	MTN	1	0
2	A	801	C8E	1	0
2	A	802	C8E	8	0
2	A	803	C8E	1	0
2	A	805	C8E	1	0
2	A	806	C8E	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	549/594 (92%)	0.39	48 (8%) <b>13</b> <b>18</b>	34, 53, 82, 105	3 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	594	PHE	8.1
1	A	288	ALA	7.9
1	A	287	SER	6.2
1	A	277	TYR	5.7
1	A	177	GLY	5.2
1	A	325	GLY	5.1
1	A	404	PHE	5.1
1	A	328	TYR	4.7
1	A	324	PRO	4.5
1	A	329	VAL	4.3
1	A	511	LEU	4.2
1	A	579	TYR	4.2
1	A	162	ASP	3.8
1	A	512	TYR	3.7
1	A	242	THR	3.7
1	A	289	THR	3.6
1	A	276	ASN	3.5
1	A	330	GLU	3.5
1	A	448	ASP	3.4
1	A	212	ASP	3.2
1	A	290	LEU	3.1
1	A	449	HIS	3.1
1	A	341	TYR	3.0
1	A	160	LEU	2.9
1	A	326	THR	2.8
1	A	176	HIS	2.7
1	A	313	ALA	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	118	TYR	2.6
1	A	312	GLY	2.6
1	A	331	ASP	2.6
1	A	275	TYR	2.6
1	A	510	GLN	2.6
1	A	555	VAL	2.5
1	A	314	GLY	2.5
1	A	76	VAL	2.5
1	A	228	ASN	2.5
1	A	6	ASP	2.4
1	A	257	ASN	2.3
1	A	592	TYR	2.3
1	A	343	THR	2.2
1	A	196	GLY	2.2
1	A	577	TYR	2.2
1	A	127	VAL	2.2
1	A	165	ARG	2.1
1	A	315	VAL	2.1
1	A	340	ILE	2.1
1	A	120	SER	2.0
1	A	115	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	MTN	A	598	12/16	0.75	0.32	12.38	51,57,60,61	12

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	596	1/1	0.87	0.33	10.29	69,69,69,69	0
2	C8E	A	805	21/21	0.65	0.39	6.36	76,97,100,101	0
2	C8E	A	803	21/21	0.56	0.48	5.72	85,97,107,107	0
2	C8E	A	806	21/21	0.42	0.42	5.23	58,94,121,122	0
2	C8E	A	801	21/21	0.76	0.28	2.92	68,76,91,92	0
2	C8E	A	802	21/21	0.93	0.33	2.58	45,56,70,74	0
2	C8E	A	800	21/21	0.77	0.24	1.24	44,56,77,81	0
2	C8E	A	804	21/21	0.90	0.13	-0.20	52,61,65,68	0
3	MG	A	595	1/1	0.96	0.06	-	76,76,76,76	0
3	MG	A	597	1/1	0.92	0.20	-	72,72,72,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.