



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:20 PM GMT

PDB ID : 4RIA
Title : FAN1 Nuclease bound to 5' phosphorylated nicked DNA
Authors : Pavletich, N.P.; Wang, R.
Deposited on : 2014-10-05
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	651	<div> <div>6%</div> <div> <div></div> <div>69%</div> <div>21%</div> <div>•</div> <div>6%</div> </div> </div>
1	B	651	<div> <div>7%</div> <div> <div></div> <div>68%</div> <div>23%</div> <div>•</div> <div>6%</div> </div> </div>
2	F	10	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>
2	L	10	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>
3	I	9	<div> <div></div> <div> <div>67%</div> <div>33%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	K	9	 67% 33%
4	G	14	 29% 21% 50%
4	J	14	 29% 21% 50%
5	E	12	 58% 42%
5	H	12	 58% 42%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fanconi-associated nuclease 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	613	Total	C	N	O	S	0	0	0
			4927	3136	877	887	27			
1	B	613	Total	C	N	O	S	0	0	0
			4927	3136	877	887	27			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	358	GLY	-	EXPRESSION TAG	UNP Q9Y2M0
A	359	ALA	-	EXPRESSION TAG	UNP Q9Y2M0
A	360	HIS	-	EXPRESSION TAG	UNP Q9Y2M0
A	361	MET	-	EXPRESSION TAG	UNP Q9Y2M0
A	362	THR	-	EXPRESSION TAG	UNP Q9Y2M0
A	363	ARG	-	EXPRESSION TAG	UNP Q9Y2M0
A	364	ASN	-	EXPRESSION TAG	UNP Q9Y2M0
A	365	GLY	-	EXPRESSION TAG	UNP Q9Y2M0
A	366	PRO	-	EXPRESSION TAG	UNP Q9Y2M0
A	367	GLY	-	EXPRESSION TAG	UNP Q9Y2M0
A	368	GLN	-	EXPRESSION TAG	UNP Q9Y2M0
A	369	THR	-	EXPRESSION TAG	UNP Q9Y2M0
A	487	ALA	VAL	ENGINEERED MUTATION	UNP Q9Y2M0
A	?	-	CYS	DELETION	UNP Q9Y2M0
A	?	-	THR	DELETION	UNP Q9Y2M0
A	?	-	TRP	DELETION	UNP Q9Y2M0
A	?	-	GLY	DELETION	UNP Q9Y2M0
A	?	-	LYS	DELETION	UNP Q9Y2M0
A	?	-	ASN	DELETION	UNP Q9Y2M0
A	?	-	LYS	DELETION	UNP Q9Y2M0
A	?	-	PRO	DELETION	UNP Q9Y2M0
A	?	-	GLY	DELETION	UNP Q9Y2M0
B	358	GLY	-	EXPRESSION TAG	UNP Q9Y2M0
B	359	ALA	-	EXPRESSION TAG	UNP Q9Y2M0
B	360	HIS	-	EXPRESSION TAG	UNP Q9Y2M0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	361	MET	-	EXPRESSION TAG	UNP Q9Y2M0
B	362	THR	-	EXPRESSION TAG	UNP Q9Y2M0
B	363	ARG	-	EXPRESSION TAG	UNP Q9Y2M0
B	364	ASN	-	EXPRESSION TAG	UNP Q9Y2M0
B	365	GLY	-	EXPRESSION TAG	UNP Q9Y2M0
B	366	PRO	-	EXPRESSION TAG	UNP Q9Y2M0
B	367	GLY	-	EXPRESSION TAG	UNP Q9Y2M0
B	368	GLN	-	EXPRESSION TAG	UNP Q9Y2M0
B	369	THR	-	EXPRESSION TAG	UNP Q9Y2M0
B	487	ALA	VAL	ENGINEERED MUTATION	UNP Q9Y2M0
B	?	-	CYS	DELETION	UNP Q9Y2M0
B	?	-	THR	DELETION	UNP Q9Y2M0
B	?	-	TRP	DELETION	UNP Q9Y2M0
B	?	-	GLY	DELETION	UNP Q9Y2M0
B	?	-	LYS	DELETION	UNP Q9Y2M0
B	?	-	ASN	DELETION	UNP Q9Y2M0
B	?	-	LYS	DELETION	UNP Q9Y2M0
B	?	-	PRO	DELETION	UNP Q9Y2M0
B	?	-	GLY	DELETION	UNP Q9Y2M0

- Molecule 2 is a DNA chain called DNA (5'-D(P*AP*GP*CP*CP*AP*CP*GP*CP*CP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	10	Total 202	C 95	N 37	O 60	P 10	0	0	0
2	L	10	Total 202	C 95	N 37	O 60	P 10	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(P*AP*GP*AP*CP*TP*CP*CP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	K	9	Total	C	N	O	P	0	0	0
			180	86	31	54	9			
3	I	9	Total	C	N	O	P	0	0	0
			180	86	31	54	9			

- Molecule 4 is a DNA chain called DNA (5'-D(*TP*TP*TP*TP*TP*TP*G*AP*GP*GP*CP*GP*TP*G)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	G	7	Total	C	N	O	P	0	0	0
			145	69	30	40	6			
4	J	7	Total	C	N	O	P	0	0	0
			145	69	30	40	6			

- Molecule 5 is a DNA chain called DNA (5'-D(P*GP*CP*TP*GP*AP*GP*GP*AP*GP*T P*CP*T)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	12	Total	C	N	O	P	0	0	0
			250	118	47	73	12			
5	H	12	Total	C	N	O	P	0	0	0
			250	118	47	73	12			

- Molecule 6 is BARIUM ION (three-letter code: BA) (formula: Ba).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Ba	0	0
			1	1		
6	A	1	Total	Ba	0	0
			1	1		



- Molecule 2: DNA (5'-D(P*AP*GP*CP*CP*AP*CP*GP*CP*CP*T)-3')



- Molecule 2: DNA (5'-D(P*AP*GP*CP*CP*AP*CP*GP*CP*CP*T)-3')





- Molecule 3: DNA (5'-D(P*AP*GP*AP*CP*TP*CP*CP*TP*C)-3')





- Molecule 3: DNA (5'-D(P*AP*GP*AP*CP*TP*CP*CP*TP*C)-3')



- Molecule 4: DNA (5'-D(*TP*TP*TP*TP*TP*TP*G*AP*GP*GP*CP*GP*TP*G)-3')




- Molecule 4: DNA (5'-D(*TP*TP*TP*TP*TP*TP*G*AP*GP*GP*CP*GP*TP*G)-3')




- Molecule 5: DNA (5'-D(P*GP*CP*TP*GP*AP*GP*GP*AP*GP*TP*CP*T)-3')

Chain E:  58% 42%



- Molecule 5: DNA (5'-D(P*GP*CP*TP*GP*AP*GP*GP*AP*GP*TP*CP*T)-3')

Chain H:  58% 42%



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	87.81Å 100.32Å 212.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 3.00 48.82 – 2.99	Depositor EDS
% Data completeness (in resolution range)	93.6 (40.00-3.00) 93.3 (48.82-2.99)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.224 , 0.267 0.224 , 0.271	Depositor DCC
R_{free} test set	1509 reflections (4.37%)	DCC
Wilson B-factor (Å ²)	55.4	Xtriage
Anisotropy	0.514	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	4 of 37624 reflections (0.011%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11410	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.72 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.3913e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/5029	0.65	0/6799
1	B	0.46	0/5029	0.65	0/6799
2	F	0.81	1/225 (0.4%)	0.89	0/342
2	L	0.78	1/225 (0.4%)	0.87	0/342
3	I	0.37	0/200	0.82	0/305
3	K	0.31	0/200	0.82	0/305
4	G	0.42	0/163	0.64	0/251
4	J	0.41	0/163	0.70	0/251
5	E	0.30	0/280	0.79	0/431
5	H	0.32	0/280	0.77	1/431 (0.2%)
All	All	0.46	2/11794 (0.0%)	0.68	1/16256 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	1	DA	OP3-P	-10.59	1.48	1.61
2	L	1	DA	OP3-P	-10.04	1.49	1.61

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	9	DG	C1'-O4'-C4'	-5.13	104.97	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4927	0	4960	108	0
1	B	4927	0	4960	114	0
2	F	202	0	112	2	0
2	L	202	0	112	4	0
3	I	180	0	102	2	0
3	K	180	0	102	2	0
4	G	145	0	80	7	0
4	J	145	0	80	6	0
5	E	250	0	136	8	0
5	H	250	0	136	5	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
All	All	11410	0	10780	241	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 241 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:682:MET:HB3	1:A:685:GLU:HG3	1.34	1.09
1:A:389:ASN:HB3	1:A:591:ARG:HH21	1.07	1.09
1:B:682:MET:HB3	1:B:685:GLU:HG3	1.39	1.05
1:B:836:SER:O	1:B:840:THR:HG23	1.66	0.95
1:A:836:SER:O	1:A:840:THR:HG23	1.67	0.93

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	607/651 (93%)	549 (90%)	44 (7%)	14 (2%)	8	36
1	B	607/651 (93%)	546 (90%)	52 (9%)	9 (2%)	13	50
All	All	1214/1302 (93%)	1095 (90%)	96 (8%)	23 (2%)	10	43

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	557	SER
1	A	558	MET
1	A	564	ALA
1	B	557	SER
1	A	760	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	536/563 (95%)	491 (92%)	45 (8%)	14	45
1	B	536/563 (95%)	492 (92%)	44 (8%)	14	46
All	All	1072/1126 (95%)	983 (92%)	89 (8%)	14	46

5 of 89 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	926	GLN
1	B	458	THR
1	B	840	THR
1	A	967	GLN
1	B	377	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	953	HIS
1	A	995	GLN
1	B	775	GLN
1	A	926	GLN
1	B	926	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	613/651 (94%)	0.05	40 (6%)	22	8	31, 54, 145, 173	0
1	B	613/651 (94%)	0.11	44 (7%)	18	7	33, 54, 144, 176	0
2	F	10/10 (100%)	-0.47	0	100	100	55, 67, 79, 86	0
2	L	10/10 (100%)	-0.48	0	100	100	59, 65, 73, 78	0
3	I	9/9 (100%)	-0.52	0	100	100	57, 63, 72, 73	0
3	K	9/9 (100%)	-0.57	0	100	100	53, 62, 70, 70	0
4	G	7/14 (50%)	-0.41	0	100	100	46, 51, 64, 68	0
4	J	7/14 (50%)	-0.57	0	100	100	45, 53, 72, 77	0
5	E	12/12 (100%)	-0.47	0	100	100	62, 79, 83, 84	0
5	H	12/12 (100%)	-0.52	0	100	100	55, 78, 84, 85	0
All	All	1302/1392 (93%)	0.05	84 (6%)	22	8	31, 55, 143, 176	0

The worst 5 of 84 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	568	GLN	9.3
1	A	568	GLN	7.2
1	B	484	PHE	6.6
1	B	796	VAL	6.4
1	B	480	LEU	5.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	BA	A	1101	1/1	1.00	0.19	-0.30	54,54,54,54	0
6	BA	B	1101	1/1	0.99	0.20	-0.36	61,61,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.