



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:37 PM GMT

PDB ID : 3RJ3  
Title : Complement components factor H CCP19-20 (S1191L mutant) and C3D in complex  
Authors : Morgan, H.P.; Hannan, J.P.  
Deposited on : 2011-04-15  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	<div> <div>2%</div> <div>76% 15% 8%</div> </div>
1	B	317	<div> <div>4%</div> <div>76% 17% 5%</div> </div>
1	C	317	<div> <div>6%</div> <div>69% 22% 8%</div> </div>
2	D	129	<div> <div>2%</div> <div>69% 25% ...</div> </div>
2	E	129	<div> <div>2%</div> <div>73% 19% 6%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	129	 A horizontal bar chart showing the quality of chain F. The bar is divided into five segments: red (29%), green (61%), yellow (15%), orange (6%), and grey (17%). The segments are labeled with their respective percentages: 29%, 61%, 15%, 6%, and 17%.

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement C3d fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	292	Total	C	N	O	S	0	4	0
			2328	1497	391	431	9			
1	B	300	Total	C	N	O	S	0	3	0
			2376	1526	395	446	9			
1	C	292	Total	C	N	O	S	0	1	0
			2306	1484	387	426	9			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLY	-	EXPRESSION TAG	UNP P01024
A	-5	PRO	-	EXPRESSION TAG	UNP P01024
A	-4	LEU	-	EXPRESSION TAG	UNP P01024
A	-3	GLY	-	EXPRESSION TAG	UNP P01024
A	-2	SER	-	EXPRESSION TAG	UNP P01024
A	-1	PRO	-	EXPRESSION TAG	UNP P01024
A	0	GLU	-	EXPRESSION TAG	UNP P01024
A	1	PHE	-	EXPRESSION TAG	UNP P01024
A	2	ARG	-	EXPRESSION TAG	UNP P01024
A	17	ALA	CYS	CONFLICT	UNP P01024
B	-6	GLY	-	EXPRESSION TAG	UNP P01024
B	-5	PRO	-	EXPRESSION TAG	UNP P01024
B	-4	LEU	-	EXPRESSION TAG	UNP P01024
B	-3	GLY	-	EXPRESSION TAG	UNP P01024
B	-2	SER	-	EXPRESSION TAG	UNP P01024
B	-1	PRO	-	EXPRESSION TAG	UNP P01024
B	0	GLU	-	EXPRESSION TAG	UNP P01024
B	1	PHE	-	EXPRESSION TAG	UNP P01024
B	2	ARG	-	EXPRESSION TAG	UNP P01024
B	17	ALA	CYS	CONFLICT	UNP P01024
C	-6	GLY	-	EXPRESSION TAG	UNP P01024
C	-5	PRO	-	EXPRESSION TAG	UNP P01024
C	-4	LEU	-	EXPRESSION TAG	UNP P01024

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	EXPRESSION TAG	UNP P01024
C	-2	SER	-	EXPRESSION TAG	UNP P01024
C	-1	PRO	-	EXPRESSION TAG	UNP P01024
C	0	GLU	-	EXPRESSION TAG	UNP P01024
C	1	PHE	-	EXPRESSION TAG	UNP P01024
C	2	ARG	-	EXPRESSION TAG	UNP P01024
C	17	ALA	CYS	CONFLICT	UNP P01024

- Molecule 2 is a protein called Complement Factor H-related protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	125	Total	C	N	O	S	0	1	0
			1005	632	181	183	9			
2	E	121	Total	C	N	O	S	0	2	0
			980	613	178	180	9			
2	F	107	Total	C	N	O	S	0	0	0
			845	535	145	156	9			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	1103	GLU	-	EXPRESSION TAG	UNP Q03591
D	1104	ALA	-	EXPRESSION TAG	UNP Q03591
D	1105	GLU	-	EXPRESSION TAG	UNP Q03591
D	1106	PHE	-	EXPRESSION TAG	UNP Q03591
D	1197	VAL	ALA	SEE REMARK 999	UNP Q03591
E	1103	GLU	-	EXPRESSION TAG	UNP Q03591
E	1104	ALA	-	EXPRESSION TAG	UNP Q03591
E	1105	GLU	-	EXPRESSION TAG	UNP Q03591
E	1106	PHE	-	EXPRESSION TAG	UNP Q03591
E	1197	VAL	ALA	SEE REMARK 999	UNP Q03591
F	1103	GLU	-	EXPRESSION TAG	UNP Q03591
F	1104	ALA	-	EXPRESSION TAG	UNP Q03591
F	1105	GLU	-	EXPRESSION TAG	UNP Q03591
F	1106	PHE	-	EXPRESSION TAG	UNP Q03591
F	1197	VAL	ALA	SEE REMARK 999	UNP Q03591

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	112	Total	O	0	0
			112	112		
4	B	97	Total	O	0	0
			97	97		
4	C	42	Total	O	0	0
			42	42		
4	D	59	Total	O	0	0
			59	59		
4	E	45	Total	O	0	0
			45	45		
4	F	17	Total	O	0	0
			17	17		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Complement C3d fragment




#### • Molecule 1: Complement C3d fragment




#### • Molecule 1: Complement C3d fragment




- Molecule 2: Complement Factor H-related protein 1



- Molecule 2: Complement Factor H-related protein 1



- Molecule 2: Complement Factor H-related protein 1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.75Å 82.62Å 85.40Å 112.61° 110.26° 99.84°	Depositor
Resolution (Å)	41.60 – 2.35 41.60 – 2.35	Depositor EDS
% Data completeness (in resolution range)	96.3 (41.60-2.35) 80.2 (41.60-2.35)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.27 (at 2.34Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.2_432)	Depositor
R, $R_{free}$	0.191 , 0.232 0.183 , 0.223	Depositor DCC
$R_{free}$ test set	3357 reflections (5.05%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.6	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 52.3	EDS
Estimated twinning fraction	0.014 for -h,-k,h+k+l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 66478 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10236	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/2383	0.40	0/3228
1	B	0.26	0/2432	0.42	0/3299
1	C	0.27	0/2358	0.40	0/3194
2	D	0.39	0/1034	0.51	1/1402 (0.1%)
2	E	0.23	0/1009	0.39	0/1364
2	F	0.35	0/866	0.51	1/1176 (0.1%)
All	All	0.29	0/10082	0.43	2/13663 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1184	THR	CB-CA-C	-5.41	96.99	111.60
2	D	1109	CYS	CA-CB-SG	5.35	123.64	114.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2328	0	2339	30	0
1	B	2376	0	2376	42	0
1	C	2306	0	2319	41	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1005	0	992	29	0
2	E	980	0	964	20	0
2	F	845	0	824	51	0
3	A	6	0	8	1	0
3	B	6	0	8	0	0
3	C	6	0	8	0	0
3	E	6	0	8	0	0
4	A	112	0	0	2	0
4	B	97	0	0	1	0
4	C	42	0	0	0	0
4	D	59	0	0	0	0
4	E	45	0	0	0	0
4	F	17	0	0	0	0
All	All	10236	0	9846	208	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 208 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:1184:THR:CG2	2:F:1185:ALA:HB2	1.14	1.59
2:F:1184:THR:HG23	2:F:1185:ALA:CB	1.26	1.57
2:F:1184:THR:HG23	2:F:1185:ALA:CA	1.52	1.36
2:F:1184:THR:CG2	2:F:1185:ALA:CB	1.90	1.32
2:F:1172:GLU:CB	2:F:1173:ILE:HG13	1.70	1.20

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/317 (92%)	285 (97%)	7 (2%)	1 (0%)	46	55
1	B	301/317 (95%)	286 (95%)	15 (5%)	0	100	100
1	C	291/317 (92%)	277 (95%)	13 (4%)	1 (0%)	46	55
2	D	124/129 (96%)	122 (98%)	2 (2%)	0	100	100
2	E	119/129 (92%)	119 (100%)	0	0	100	100
2	F	101/129 (78%)	96 (95%)	4 (4%)	1 (1%)	19	20
All	All	1229/1338 (92%)	1185 (96%)	41 (3%)	3 (0%)	52	63

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	1173	ILE
1	A	141	ARG
1	C	267	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/264 (93%)	237 (97%)	8 (3%)	45	58
1	B	253/264 (96%)	239 (94%)	14 (6%)	27	32
1	C	242/264 (92%)	233 (96%)	9 (4%)	41	53
2	D	113/115 (98%)	103 (91%)	10 (9%)	12	12
2	E	111/115 (96%)	108 (97%)	3 (3%)	52	67
2	F	96/115 (84%)	85 (88%)	11 (12%)	7	6
All	All	1060/1137 (93%)	1005 (95%)	55 (5%)	29	35

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	95	LEU
1	C	290	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	F	1197	VAL
1	C	98	ASN
1	C	143	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	133	HIS
1	C	225	ASN
2	E	1212	HIS
1	C	159	GLN
1	A	225	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	311	-	5,5,5	0.39	0	5,5,5	0.25	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	311	-	5,5,5	0.47	0	5,5,5	0.63	0
3	GOL	C	311	-	5,5,5	0.45	0	5,5,5	0.21	0
3	GOL	E	1	-	5,5,5	0.32	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	311	-	-	0/4/4/4	0/0/0/0
3	GOL	B	311	-	-	0/4/4/4	0/0/0/0
3	GOL	C	311	-	-	0/4/4/4	0/0/0/0
3	GOL	E	1	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	311	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	292/317 (92%)	-0.20	4 (1%) 78 87	21, 38, 74, 115	0
1	B	300/317 (94%)	0.25	14 (4%) 35 50	21, 45, 88, 177	0
1	C	292/317 (92%)	0.19	18 (6%) 24 36	28, 59, 113, 174	0
2	D	125/129 (96%)	-0.10	2 (1%) 74 84	28, 44, 76, 135	0
2	E	121/129 (93%)	-0.05	2 (1%) 73 83	32, 50, 82, 102	0
2	F	107/129 (82%)	1.27	38 (35%) 0 0	35, 77, 129, 184	0
All	All	1237/1338 (92%)	0.15	78 (6%) 23 35	21, 49, 103, 184	0

The worst 5 of 78 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	270	GLY	9.2
1	B	300	ASN	7.2
2	F	1183	TRP	7.0
1	B	301	LEU	7.0
1	B	269	TYR	6.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	311	6/6	0.85	0.18	1.72	61,67,68,68	0
3	GOL	A	311	6/6	0.98	0.13	0.02	29,39,43,45	0
3	GOL	E	1	6/6	0.90	0.19	-	51,58,67,68	0
3	GOL	C	311	6/6	0.84	0.17	-	62,64,67,79	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.