



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:43 PM GMT

PDB ID : 3RPN
Title : Crystal structure of human kappa class glutathione transferase in complex with S-hexylglutathione
Authors : Wang, B.; Peng, Y.; Zhang, T.; Ding, J.
Deposited on : 2011-04-27
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	<div> <div>3%</div> <div>84%</div> <div>9%</div> <div>6%</div> </div>
1	B	234	<div> <div>4%</div> <div>79%</div> <div>13%</div> <div>6%</div> </div>
1	C	234	<div> <div>3%</div> <div>85%</div> <div>9%</div> <div>6%</div> </div>
1	D	234	<div> <div>2%</div> <div>91%</div> <div>6%</div> </div>
1	E	234	<div> <div>3%</div> <div>83%</div> <div>10%</div> <div>6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	234	<div><div></div><div>7%</div><div>85%</div><div>9%</div><div>6%</div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11418 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutathione S-transferase kappa 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	0	0	0
			1743	1122	301	308	12			
1	B	220	Total	C	N	O	S	0	0	0
			1743	1122	301	308	12			
1	C	220	Total	C	N	O	S	0	0	0
			1743	1122	301	308	12			
1	D	220	Total	C	N	O	S	0	0	0
			1743	1122	301	308	12			
1	E	220	Total	C	N	O	S	0	0	0
			1743	1122	301	308	12			
1	F	220	Total	C	N	O	S	0	0	0
			1743	1122	301	308	12			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	227	LEU	-	EXPRESSION TAG	UNP Q9Y2Q3
A	228	GLU	-	EXPRESSION TAG	UNP Q9Y2Q3
A	229	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
A	230	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
A	231	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
A	232	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
A	233	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
A	234	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
B	227	LEU	-	EXPRESSION TAG	UNP Q9Y2Q3
B	228	GLU	-	EXPRESSION TAG	UNP Q9Y2Q3
B	229	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
B	230	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
B	231	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
B	232	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
B	233	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
B	234	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
C	227	LEU	-	EXPRESSION TAG	UNP Q9Y2Q3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	228	GLU	-	EXPRESSION TAG	UNP Q9Y2Q3
C	229	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
C	230	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
C	231	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
C	232	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
C	233	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
C	234	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
D	227	LEU	-	EXPRESSION TAG	UNP Q9Y2Q3
D	228	GLU	-	EXPRESSION TAG	UNP Q9Y2Q3
D	229	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
D	230	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
D	231	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
D	232	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
D	233	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
D	234	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
E	227	LEU	-	EXPRESSION TAG	UNP Q9Y2Q3
E	228	GLU	-	EXPRESSION TAG	UNP Q9Y2Q3
E	229	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
E	230	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
E	231	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
E	232	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
E	233	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
E	234	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
F	227	LEU	-	EXPRESSION TAG	UNP Q9Y2Q3
F	228	GLU	-	EXPRESSION TAG	UNP Q9Y2Q3
F	229	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
F	230	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
F	231	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
F	232	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
F	233	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3
F	234	HIS	-	EXPRESSION TAG	UNP Q9Y2Q3

- Molecule 2 is S-HEXYLGLUTATHIONE (three-letter code: GTX) (formula: C₁₆H₃₀N₃O₆S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 26	C 16	N 3	O 6	S 1	0	0
2	B	1	Total 26	C 16	N 3	O 6	S 1	0	0
2	C	1	Total 26	C 16	N 3	O 6	S 1	0	0
2	D	1	Total 26	C 16	N 3	O 6	S 1	0	0
2	E	1	Total 26	C 16	N 3	O 6	S 1	0	0
2	F	1	Total 26	C 16	N 3	O 6	S 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	124	Total O 124 124	0	0
3	B	128	Total O 128 128	0	0
3	C	117	Total O 117 117	0	0
3	D	160	Total O 160 160	0	0
3	E	154	Total O 154 154	0	0
3	F	121	Total O 121 121	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutathione S-transferase kappa 1




- Molecule 1: Glutathione S-transferase kappa 1




- Molecule 1: Glutathione S-transferase kappa 1




- Molecule 1: Glutathione S-transferase kappa 1



- Molecule 1: Glutathione S-transferase kappa 1



• Molecule 1: Glutathione S-transferase kappa 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.36Å 199.82Å 66.71Å 90.00° 116.12° 90.00°	Depositor
Resolution (Å)	29.95 – 1.90 29.95 – 1.90	Depositor EDS
% Data completeness (in resolution range)	95.5 (29.95-1.90) 95.6 (29.95-1.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.66 (at 1.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.183 , 0.218 0.180 , 0.214	Depositor DCC
R_{free} test set	5845 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.466	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 59.5	EDS
Estimated twinning fraction	0.025 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 116703 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11418	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/1787	0.47	0/2421
1	B	0.35	0/1787	0.49	0/2421
1	C	0.36	0/1787	0.49	0/2421
1	D	0.39	0/1787	0.51	0/2421
1	E	0.37	0/1787	0.51	0/2421
1	F	0.34	0/1787	0.47	0/2421
All	All	0.36	0/10722	0.49	0/14526

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1743	0	1775	19	0
1	B	1743	0	1775	25	0
1	C	1743	0	1775	13	0
1	D	1743	0	1775	6	0
1	E	1743	0	1775	20	0
1	F	1743	0	1775	18	0
2	A	26	0	28	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	26	0	28	3	0
2	C	26	0	28	2	0
2	D	26	0	28	2	0
2	E	26	0	28	2	0
2	F	26	0	28	5	0
3	A	124	0	0	0	0
3	B	128	0	0	1	0
3	C	117	0	0	2	0
3	D	160	0	0	2	0
3	E	154	0	0	5	0
3	F	121	0	0	3	0
All	All	11418	0	10818	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:74:ARG:HE	1:C:82:HIS:CE1	1.96	0.83
1:C:74:ARG:HE	1:C:82:HIS:HE1	1.35	0.75
1:F:17:PRO:HD2	2:F:302:GTX:HS21	1.70	0.73
1:A:169:LYS:O	1:A:173:GLU:HG2	1.89	0.72
1:A:88:LEU:HD22	2:A:302:GTX:HS52	1.73	0.69
1:D:88:LEU:HD22	2:D:302:GTX:HS32	1.74	0.69
1:F:221:ALA:HB2	3:F:787:HOH:O	1.92	0.68
1:E:47:ILE:HD13	1:E:176:CYS:SG	2.35	0.66
1:E:221:ALA:HB2	3:E:788:HOH:O	1.94	0.66
1:E:124:ARG:NH2	3:E:737:HOH:O	2.25	0.64
1:A:82:HIS:HD2	1:A:127:SER:O	1.80	0.63
1:F:43:LEU:HG	1:F:45:THR:HG22	1.80	0.63
1:B:15:LEU:HG	2:B:301:GTX:HS63	1.82	0.62
1:C:156:LEU:O	1:C:159:ILE:HG12	1.99	0.61
1:E:88:LEU:HD13	2:E:302:GTX:HS52	1.82	0.61
1:B:193:GLN:HB2	3:B:700:HOH:O	2.01	0.60
1:B:92:LEU:HG	2:B:301:GTX:HS62	1.84	0.59
1:D:128:ARG:NH2	3:D:762:HOH:O	2.37	0.58
1:A:102:PHE:HA	1:A:159:ILE:HD11	1.87	0.57
1:F:88:LEU:HD11	3:F:658:HOH:O	2.05	0.57
1:B:169:LYS:O	1:B:173:GLU:HG3	2.05	0.57
1:B:156:LEU:O	1:B:159:ILE:HG12	2.04	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:PHE:HA	1:B:159:ILE:HD11	1.87	0.57
1:C:124:ARG:HD2	1:C:141:ALA:HB2	1.87	0.57
1:F:197:LEU:HD12	1:F:197:LEU:N	2.20	0.56
1:B:61:ARG:NH2	1:E:176:CYS:HB3	2.20	0.56
1:C:66:MET:HE2	1:C:83:PHE:CE1	2.41	0.56
1:F:121:LEU:O	1:F:125:VAL:HG12	2.06	0.54
1:C:108:LEU:HD13	1:C:163:LYS:HE2	1.89	0.54
1:C:72:LEU:CD1	3:C:723:HOH:O	2.54	0.54
1:A:67:ALA:O	1:A:71:LYS:HG3	2.07	0.53
1:A:15:LEU:HG	2:A:302:GTX:HS63	1.89	0.53
1:B:66:MET:HE2	1:B:88:LEU:HD11	1.91	0.53
1:E:196:MET:C	1:E:197:LEU:HD12	2.28	0.53
1:A:177:ARG:HG2	1:A:177:ARG:HH11	1.74	0.53
1:A:82:HIS:CD2	1:A:127:SER:O	2.61	0.52
1:F:17:PRO:CD	2:F:302:GTX:HS21	2.39	0.52
1:B:124:ARG:HD2	1:B:141:ALA:HB2	1.91	0.51
1:C:74:ARG:NE	1:C:82:HIS:HE1	2.04	0.51
1:E:4:LEU:HD22	1:E:4:LEU:N	2.26	0.51
1:F:57:GLY:HA3	1:F:88:LEU:HD11	1.93	0.51
1:A:170:GLU:HA	1:A:173:GLU:CG	2.42	0.50
1:C:89:SER:O	1:C:93:GLU:HG3	2.12	0.50
1:B:156:LEU:HD12	1:B:159:ILE:HG12	1.94	0.49
1:B:82:HIS:HD2	1:B:127:SER:O	1.96	0.49
1:F:59:LEU:HD23	1:F:62:LYS:HG3	1.94	0.49
1:E:156:LEU:O	1:E:159:ILE:HG12	2.12	0.49
1:B:67:ALA:O	1:B:71:LYS:HG3	2.13	0.49
1:F:74:ARG:HB2	1:F:81:ILE:HG22	1.95	0.49
1:B:61:ARG:NH2	1:E:50:ASP:OD2	2.45	0.49
1:B:66:MET:CE	1:B:83:PHE:HE1	2.26	0.49
1:A:84:PRO:HB3	1:A:129:ASN:HD21	1.78	0.48
1:B:132:ILE:HD13	1:B:132:ILE:H	1.78	0.47
1:A:72:LEU:HG	1:D:209:LEU:HD21	1.96	0.47
1:E:72:LEU:HD13	3:E:511:HOH:O	2.15	0.47
1:F:109:GLU:HG3	3:F:626:HOH:O	2.14	0.47
1:B:61:ARG:HH22	1:E:177:ARG:NH2	2.13	0.46
1:B:154:GLY:O	1:B:158:LYS:HG3	2.15	0.46
1:F:66:MET:HE2	2:F:302:GTX:HS52	1.98	0.46
1:A:45:THR:O	1:A:49:LYS:HD3	2.15	0.45
1:B:183:LEU:HB3	1:B:184:PRO:HA	1.98	0.45
1:B:66:MET:HE2	1:B:83:PHE:HE1	1.82	0.44
1:A:84:PRO:HB3	1:A:129:ASN:ND2	2.31	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:180:ALA:HB2	1:F:185:ILE:HD13	1.99	0.44
1:B:83:PHE:CD2	1:B:84:PRO:HD2	2.53	0.44
1:E:4:LEU:HD22	1:E:4:LEU:H	1.83	0.44
1:F:17:PRO:HG3	2:F:302:GTX:HS42	2.00	0.44
1:C:17:PRO:HD2	2:C:301:GTX:HS22	1.99	0.44
1:A:163:LYS:HB2	1:A:163:LYS:HE3	1.80	0.43
1:A:83:PHE:CG	1:A:84:PRO:HD2	2.54	0.43
1:F:57:GLY:HA3	1:F:88:LEU:CD1	2.49	0.43
1:D:17:PRO:HD2	2:D:302:GTX:HS21	2.00	0.43
1:C:183:LEU:HB3	1:C:184:PRO:HA	2.01	0.43
1:B:102:PHE:HA	1:B:159:ILE:CD1	2.48	0.42
1:B:59:LEU:HD23	1:B:62:LYS:HG3	2.01	0.42
1:B:131:ASP:O	1:B:137:SER:HB3	2.19	0.42
1:E:43:LEU:HD11	1:E:97:LEU:HD13	2.01	0.42
1:F:132:ILE:O	1:F:138:ILE:HD11	2.19	0.41
1:B:17:PRO:HD2	2:B:301:GTX:SG2	2.60	0.41
1:E:109:GLU:OE2	1:E:158:LYS:HE3	2.21	0.41
1:E:66:MET:HB3	2:E:302:GTX:HS42	2.03	0.41
1:E:197:LEU:HD12	1:E:197:LEU:N	2.35	0.41
1:C:58:LEU:O	1:C:60:PRO:HD3	2.21	0.41
1:A:66:MET:CE	1:A:88:LEU:HD11	2.50	0.41
1:A:170:GLU:HA	1:A:173:GLU:HG2	2.02	0.41
1:A:123:MET:O	1:A:127:SER:HB3	2.20	0.41
2:C:301:GTX:O31	1:F:62:LYS:HE2	2.21	0.41
1:E:136:GLN:HG2	3:E:746:HOH:O	2.21	0.41
1:E:47:ILE:HD11	1:E:175:ALA:HB3	2.03	0.41
1:E:109:GLU:HG3	3:E:306:HOH:O	2.21	0.41
1:E:30:GLN:HA	1:E:35:ILE:HD11	2.02	0.41
1:B:83:PHE:CG	1:B:84:PRO:HD2	2.56	0.41
1:A:17:PRO:HD2	2:A:302:GTX:SG2	2.62	0.40
1:C:124:ARG:NH2	3:C:345:HOH:O	2.54	0.40
1:F:88:LEU:HD12	2:F:302:GTX:HS52	2.03	0.40
1:D:177:ARG:NH1	3:D:536:HOH:O	2.51	0.40
1:D:49:LYS:HE2	1:D:49:LYS:HB3	1.82	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/234 (93%)	214 (98%)	4 (2%)	0	100	100
1	B	218/234 (93%)	216 (99%)	2 (1%)	0	100	100
1	C	218/234 (93%)	215 (99%)	3 (1%)	0	100	100
1	D	218/234 (93%)	216 (99%)	2 (1%)	0	100	100
1	E	218/234 (93%)	217 (100%)	1 (0%)	0	100	100
1	F	218/234 (93%)	216 (99%)	2 (1%)	0	100	100
All	All	1308/1404 (93%)	1294 (99%)	14 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/201 (94%)	183 (97%)	5 (3%)	52	43
1	B	188/201 (94%)	175 (93%)	13 (7%)	19	8
1	C	188/201 (94%)	185 (98%)	3 (2%)	70	66
1	D	188/201 (94%)	185 (98%)	3 (2%)	70	66
1	E	188/201 (94%)	184 (98%)	4 (2%)	61	55
1	F	188/201 (94%)	185 (98%)	3 (2%)	70	66
All	All	1128/1206 (94%)	1097 (97%)	31 (3%)	52	43

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	LEU
1	A	122	TRP
1	A	159	ILE
1	A	177	ARG
1	A	197	LEU
1	B	44	ILE
1	B	61	ARG
1	B	72	LEU
1	B	88	LEU
1	B	92	LEU
1	B	122	TRP
1	B	124	ARG
1	B	129	ASN
1	B	132	ILE
1	B	143	GLU
1	B	165	LYS
1	B	193	GLN
1	B	197	LEU
1	C	61	ARG
1	C	122	TRP
1	C	144	LYS
1	D	72	LEU
1	D	122	TRP
1	D	177	ARG
1	E	72	LEU
1	E	90	VAL
1	E	122	TRP
1	E	190	VAL
1	F	72	LEU
1	F	122	TRP
1	F	190	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	A	75	HIS
1	A	78	GLN
1	A	82	HIS
1	A	129	ASN
1	B	68	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	78	GLN
1	B	82	HIS
1	B	151	GLN
1	B	153	GLN
1	B	208	HIS
1	C	31	ASN
1	C	38	GLN
1	C	82	HIS
1	C	110	HIS
1	C	208	HIS
1	D	38	GLN
1	D	78	GLN
1	D	110	HIS
1	D	166	ASN
1	E	36	ASN
1	E	38	GLN
1	E	78	GLN
1	F	31	ASN
1	F	78	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GTX	A	302	-	19,25,25	2.82	2 (10%)	23,30,30	1.30	1 (4%)
2	GTX	B	301	-	19,25,25	2.82	2 (10%)	23,30,30	1.21	1 (4%)
2	GTX	C	301	-	19,25,25	2.73	2 (10%)	23,30,30	1.30	2 (8%)
2	GTX	D	302	-	19,25,25	2.83	2 (10%)	23,30,30	1.17	2 (8%)
2	GTX	E	302	-	19,25,25	2.76	2 (10%)	23,30,30	1.23	1 (4%)
2	GTX	F	302	-	19,25,25	2.76	2 (10%)	23,30,30	1.51	5 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTX	A	302	-	-	0/24/30/30	0/0/0/0
2	GTX	B	301	-	-	0/24/30/30	0/0/0/0
2	GTX	C	301	-	-	0/24/30/30	0/0/0/0
2	GTX	D	302	-	-	0/24/30/30	0/0/0/0
2	GTX	E	302	-	-	0/24/30/30	0/0/0/0
2	GTX	F	302	-	-	0/24/30/30	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	302	GTX	OE1-CD1	8.00	1.40	1.23
2	D	302	GTX	OE1-CD1	8.17	1.40	1.23
2	C	301	GTX	O2-C2	8.19	1.39	1.23
2	E	302	GTX	O2-C2	8.28	1.39	1.23
2	E	302	GTX	OE1-CD1	8.37	1.40	1.23
2	C	301	GTX	OE1-CD1	8.37	1.40	1.23
2	A	302	GTX	OE1-CD1	8.40	1.40	1.23
2	B	301	GTX	OE1-CD1	8.41	1.40	1.23
2	A	302	GTX	O2-C2	8.64	1.40	1.23
2	F	302	GTX	O2-C2	8.69	1.40	1.23
2	B	301	GTX	O2-C2	8.72	1.40	1.23
2	D	302	GTX	O2-C2	8.95	1.40	1.23

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	GTX	CB2-SG2-C1S	-4.53	88.81	102.41
2	A	302	GTX	CB2-SG2-C1S	-4.30	89.49	102.41
2	C	301	GTX	CB2-SG2-C1S	-3.99	90.43	102.41
2	F	302	GTX	CB2-SG2-C1S	-3.36	92.32	102.41
2	E	302	GTX	CB2-SG2-C1S	-3.32	92.43	102.41
2	F	302	GTX	CA3-N3-C2	-3.23	117.89	122.34
2	D	302	GTX	CB2-SG2-C1S	-2.69	94.33	102.41
2	C	301	GTX	CB1-CG1-CD1	-2.33	107.74	113.27
2	F	302	GTX	CB1-CG1-CD1	-2.13	108.21	113.27
2	D	302	GTX	CB1-CG1-CD1	-2.11	108.25	113.27
2	F	302	GTX	CG1-CB1-CA1	-2.05	110.48	114.03
2	F	302	GTX	CG1-CD1-N2	2.36	119.68	115.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	302	GTX	3	0
2	B	301	GTX	3	0
2	C	301	GTX	2	0
2	D	302	GTX	2	0
2	E	302	GTX	2	0
2	F	302	GTX	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	220/234 (94%)	0.04	8 (3%)	46	50	23, 36, 61, 104	0
1	B	220/234 (94%)	0.06	9 (4%)	41	45	25, 38, 69, 94	0
1	C	220/234 (94%)	-0.06	7 (3%)	51	54	23, 36, 60, 85	0
1	D	220/234 (94%)	-0.26	5 (2%)	64	67	21, 31, 57, 80	0
1	E	220/234 (94%)	-0.12	7 (3%)	51	54	21, 32, 63, 98	0
1	F	220/234 (94%)	0.10	16 (7%)	18	20	25, 38, 68, 100	0
All	All	1320/1404 (94%)	-0.04	52 (3%)	43	47	21, 35, 63, 104	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	GLY	9.0
1	C	221	ALA	7.9
1	A	221	ALA	6.2
1	F	2	GLY	6.1
1	B	221	ALA	5.8
1	E	2	GLY	4.7
1	D	2	GLY	4.5
1	F	88	LEU	4.3
1	B	2	GLY	4.2
1	A	4	LEU	3.9
1	E	58	LEU	3.8
1	F	92	LEU	3.7
1	A	3	PRO	3.7
1	D	88	LEU	3.5
1	F	86	ASP	3.5
1	F	3	PRO	3.4
1	B	150	GLU	3.3
1	F	150	GLU	3.3
1	F	221	ALA	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	3	PRO	3.2
1	B	3	PRO	3.1
1	B	10	LEU	3.1
1	E	88	LEU	3.0
1	F	58	LEU	2.9
1	F	4	LEU	2.8
1	A	220	PRO	2.7
1	C	2	GLY	2.7
1	F	85	LYS	2.6
1	F	143	GLU	2.6
1	F	191	ASP	2.5
1	C	49	LYS	2.5
1	F	149	ALA	2.5
1	E	92	LEU	2.5
1	C	149	ALA	2.4
1	E	86	ASP	2.4
1	C	150	GLU	2.3
1	C	12	TYR	2.3
1	F	18	TYR	2.3
1	B	146	GLY	2.2
1	A	49	LYS	2.2
1	D	221	ALA	2.2
1	B	52	GLY	2.2
1	D	85	LYS	2.2
1	B	93	GLU	2.1
1	F	21	LEU	2.1
1	F	89	SER	2.1
1	A	10	LEU	2.1
1	A	12	TYR	2.1
1	E	59	LEU	2.1
1	D	3	PRO	2.0
1	C	54	LYS	2.0
1	B	136	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GTX	A	302	26/26	0.89	0.15	0.65	35,48,60,62	0
2	GTX	B	301	26/26	0.92	0.15	0.54	35,44,90,93	0
2	GTX	E	302	26/26	0.93	0.14	0.41	25,32,73,76	0
2	GTX	F	302	26/26	0.94	0.15	0.27	30,35,94,95	0
2	GTX	D	302	26/26	0.94	0.13	0.07	23,30,58,64	0
2	GTX	C	301	26/26	0.94	0.12	-0.03	30,38,60,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.