



# Full wwPDB X-ray Structure Validation Report ⓘ

May 6, 2016 – 02:19 AM EDT

PDB ID : 4RQ9  
Title : Crystal structure of the chromophore-binding domain of Stigmatella auranti-  
aca bacteriophytochrome (Thr289His mutant) in the Pr state  
Authors : Woitowich, N.C.; Halavaty, A.S.; Gallagher, K.D.; Nugent, A.C.; Patel, H.;  
Duong, P.; Kovaleva, S.E.; St. Peter, S.; Ozarowski, W.B.; Hernandez, C.N.;  
Stojkovic, E.A.  
Deposited on : 2014-10-31  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027457  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	<div> <div>3%</div> <div>81%</div> <div>5%</div> <div>13%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SRT	A	402	-	-	-	X
5	GOL	A	407	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	410	-	-	-	X
5	GOL	A	418	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2802 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Photoreceptor-histidine kinase BphP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	0	10	0
			2471	1564	451	448	8			

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP Q097N3
A	-18	GLY	-	EXPRESSION TAG	UNP Q097N3
A	-17	SER	-	EXPRESSION TAG	UNP Q097N3
A	-16	SER	-	EXPRESSION TAG	UNP Q097N3
A	-15	HIS	-	EXPRESSION TAG	UNP Q097N3
A	-14	HIS	-	EXPRESSION TAG	UNP Q097N3
A	-13	HIS	-	EXPRESSION TAG	UNP Q097N3
A	-12	HIS	-	EXPRESSION TAG	UNP Q097N3
A	-11	HIS	-	EXPRESSION TAG	UNP Q097N3
A	-10	HIS	-	EXPRESSION TAG	UNP Q097N3
A	-9	SER	-	EXPRESSION TAG	UNP Q097N3
A	-8	SER	-	EXPRESSION TAG	UNP Q097N3
A	-7	GLY	-	EXPRESSION TAG	UNP Q097N3
A	-6	LEU	-	EXPRESSION TAG	UNP Q097N3
A	-5	VAL	-	EXPRESSION TAG	UNP Q097N3
A	-4	PRO	-	EXPRESSION TAG	UNP Q097N3
A	-3	ARG	-	EXPRESSION TAG	UNP Q097N3
A	-2	GLY	-	EXPRESSION TAG	UNP Q097N3
A	-1	SER	-	EXPRESSION TAG	UNP Q097N3
A	0	HIS	-	EXPRESSION TAG	UNP Q097N3
A	289	HIS	THR	ENGINEERED MUTATION	UNP Q097N3

- Molecule 2 is BILIVERDINE IX ALPHA (three-letter code: BLA) (formula:  $C_{33}H_{34}N_4O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			43	33	4	6		

- Molecule 3 is S,R MESO-TARTARIC ACID (three-letter code: SRT) (formula:  $C_4H_6O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	178	Total	O	0	8
			184	184		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Photoreceptor-histidine kinase BphP





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	131.81Å 131.81Å 97.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.96 – 2.50 29.96 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.96-2.50) 99.8 (29.96-2.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.65 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.155 , 0.178 0.158 , 0.179	Depositor DCC
$R_{free}$ test set	1738 reflections (5.40%)	DCC
Wilson B-factor (Å <sup>2</sup> )	55.0	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 55.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2802	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BLA, EDO, SRT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.57	0/2530	0.82	5/3439 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	ASP	CB-CG-OD1	8.41	125.87	118.30
1	A	173	ARG	NE-CZ-NH1	-6.75	116.93	120.30
1	A	195	ARG	NE-CZ-NH1	5.95	123.27	120.30
1	A	214	ARG	NE-CZ-NH2	5.78	123.19	120.30
1	A	173	ARG	NE-CZ-NH2	5.33	122.97	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2471	0	2451	13	0
2	A	43	0	30	3	0
3	A	10	0	4	0	0
4	A	4	0	6	0	0
5	A	90	0	120	2	0
6	A	184	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2802	0	2611	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:ARG:HH11	1:A:195:ARG:HG2	1.59	0.67
1:A:269:GLN:HG3	6:A:595:HOH:O	1.95	0.67
2:A:401:BLA:NB	2:A:401:BLA:HMA1	2.13	0.64
1:A:78:GLU:HB2	1:A:79:PRO:HD3	1.82	0.60
1:A:195:ARG:HH11	1:A:195:ARG:CG	2.19	0.55
1:A:177:TYR:CZ	1:A:185:GLY:HA3	2.43	0.53
1:A:317:ALA:N	6:A:543:HOH:O	2.41	0.51
2:A:401:BLA:CB	2:A:401:BLA:HMC1	2.44	0.48
1:A:173:ARG:NH2	1:A:197:ASP:O	2.49	0.46
2:A:401:BLA:HB	2:A:401:BLA:HMA1	1.80	0.45
1:A:195:ARG:NH1	1:A:195:ARG:HG2	2.30	0.45
1:A:249:PHE:CE2	5:A:409:GOL:H12	2.53	0.43
1:A:236[A]:ARG:HH11	1:A:236[A]:ARG:HB3	1.83	0.42
1:A:166:ARG:O	5:A:414:GOL:H31	2.20	0.42
1:A:102:ASP:O	1:A:104:VAL:HG23	2.20	0.42
1:A:193:ASP:OD1	1:A:195:ARG:NH1	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	312/348 (90%)	306 (98%)	5 (2%)	1 (0%)	46 68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	128	HIS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	266 / 294 (90%)	260 (98%)	6 (2%)	58 83

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	GLN
1	A	195	ARG
1	A	223	ARG
1	A	236[A]	ARG
1	A	236[B]	ARG
1	A	295	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	155	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	BLA	A	401	-	36,46,46	4.57	14 (38%)	45,67,67	2.37	15 (33%)
3	SRT	A	402	-	3,9,9	0.84	0	6,12,12	0.80	0
4	EDO	A	403	-	3,3,3	0.51	0	2,2,2	0.31	0
5	GOL	A	404	-	5,5,5	0.22	0	5,5,5	0.60	0
5	GOL	A	405	-	5,5,5	0.32	0	5,5,5	0.36	0
5	GOL	A	406	-	5,5,5	0.23	0	5,5,5	0.24	0
5	GOL	A	407	-	5,5,5	0.76	0	5,5,5	0.72	0
5	GOL	A	408	-	5,5,5	0.21	0	5,5,5	0.91	0
5	GOL	A	409	-	5,5,5	0.60	0	5,5,5	0.68	0
5	GOL	A	410	-	5,5,5	0.57	0	5,5,5	0.59	0
5	GOL	A	411	-	5,5,5	0.42	0	5,5,5	0.40	0
5	GOL	A	412	-	5,5,5	0.32	0	5,5,5	0.18	0
5	GOL	A	413	-	5,5,5	0.52	0	5,5,5	0.51	0
5	GOL	A	414	-	5,5,5	0.72	0	5,5,5	0.93	0
5	GOL	A	415	-	5,5,5	0.51	0	5,5,5	0.73	0
5	GOL	A	416	-	5,5,5	0.77	0	5,5,5	1.12	0
5	GOL	A	417	-	5,5,5	0.47	0	5,5,5	0.28	0
5	GOL	A	418	-	5,5,5	0.28	0	5,5,5	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BLA	A	401	-	-	0/22/74/74	0/4/4/4
3	SRT	A	402	-	-	0/4/12/12	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	403	-	-	0/1/1/1	0/0/0/0
5	GOL	A	404	-	-	0/4/4/4	0/0/0/0
5	GOL	A	405	-	-	0/4/4/4	0/0/0/0
5	GOL	A	406	-	-	0/4/4/4	0/0/0/0
5	GOL	A	407	-	-	0/4/4/4	0/0/0/0
5	GOL	A	408	-	-	0/4/4/4	0/0/0/0
5	GOL	A	409	-	-	0/4/4/4	0/0/0/0
5	GOL	A	410	-	-	0/4/4/4	0/0/0/0
5	GOL	A	411	-	-	0/4/4/4	0/0/0/0
5	GOL	A	412	-	-	0/4/4/4	0/0/0/0
5	GOL	A	413	-	-	0/4/4/4	0/0/0/0
5	GOL	A	414	-	-	0/4/4/4	0/0/0/0
5	GOL	A	415	-	-	0/4/4/4	0/0/0/0
5	GOL	A	416	-	-	0/4/4/4	0/0/0/0
5	GOL	A	417	-	-	0/4/4/4	0/0/0/0
5	GOL	A	418	-	-	0/4/4/4	0/0/0/0

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	BLA	C4D-C3D	-3.97	1.38	1.45
2	A	401	BLA	C3B-C4B	-3.22	1.37	1.47
2	A	401	BLA	C1D-C2D	-3.15	1.38	1.45
2	A	401	BLA	C1B-C2B	-3.08	1.39	1.45
2	A	401	BLA	C1C-C2C	-2.53	1.40	1.47
2	A	401	BLA	C1A-CHA	2.15	1.48	1.40
2	A	401	BLA	C4A-CHB	2.47	1.49	1.40
2	A	401	BLA	CHD-C1D	4.63	1.51	1.40
2	A	401	BLA	CBC-CAC	4.78	1.53	1.30
2	A	401	BLA	CHD-C4C	5.14	1.49	1.37
2	A	401	BLA	CHB-C1B	7.13	1.48	1.34
2	A	401	BLA	OB-C4B	11.36	1.45	1.23
2	A	401	BLA	OC-C1C	11.86	1.46	1.23
2	A	401	BLA	CHA-C4D	16.76	1.50	1.35

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	BLA	C4C-NC-C1C	-7.10	100.44	110.73
2	A	401	BLA	C1B-NB-C4B	-4.70	103.93	110.73
2	A	401	BLA	CHB-C1B-C2B	-4.58	118.07	126.96
2	A	401	BLA	CHA-C4D-C3D	-3.52	117.47	125.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	BLA	CHD-C4C-C3C	-3.31	119.31	128.02
2	A	401	BLA	CHD-C1D-C2D	-3.05	117.42	124.71
2	A	401	BLA	OB-C4B-C3B	-2.33	124.26	130.06
2	A	401	BLA	CHA-C4D-ND	2.24	132.69	128.71
2	A	401	BLA	CMC-C2C-C1C	2.32	127.37	121.40
2	A	401	BLA	C2B-C1B-NB	2.41	110.34	106.94
2	A	401	BLA	CMB-C2B-C1B	2.47	127.43	124.20
2	A	401	BLA	CAB-C3B-C4B	2.86	131.81	123.58
2	A	401	BLA	C3C-C4C-NC	3.75	112.60	106.85
2	A	401	BLA	C2C-C1C-NC	4.08	118.08	106.40
2	A	401	BLA	CHD-C1D-ND	5.08	134.38	124.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	BLA	3	0
5	A	409	GOL	1	0
5	A	414	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	304/348 (87%)	-0.24	12 (3%) 43 48	38, 50, 89, 140	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	16	THR	6.5
1	A	128	HIS	5.3
1	A	130	GLU	4.4
1	A	318	ALA	2.8
1	A	286	ILE	2.4
1	A	20	ARG	2.3
1	A	222	LEU	2.3
1	A	274	ILE	2.3
1	A	131	ALA	2.3
1	A	273	SER	2.2
1	A	292	SER	2.1
1	A	251[A]	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron



density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	A	410	6/6	0.81	0.35	11.66	77,94,102,103	0
3	SRT	A	402	10/10	0.86	0.45	10.23	67,87,98,99	0
5	GOL	A	418	6/6	0.86	0.36	6.16	103,110,113,119	0
5	GOL	A	407	6/6	0.62	0.32	5.30	77,92,96,101	0
4	EDO	A	403	4/4	0.86	0.20	1.78	69,75,85,88	0
5	GOL	A	404	6/6	0.92	0.23	1.65	60,75,77,81	0
5	GOL	A	411	6/6	0.91	0.18	1.54	65,84,91,101	0
2	BLA	A	401	43/43	0.96	0.18	0.52	38,42,55,76	1
5	GOL	A	415	6/6	0.89	0.17	0.39	80,82,90,99	0
5	GOL	A	408	6/6	0.90	0.16	-0.05	67,89,93,94	0
5	GOL	A	405	6/6	0.91	0.32	-	77,91,97,103	0
5	GOL	A	406	6/6	0.86	0.36	-	98,106,112,112	0
5	GOL	A	409	6/6	0.75	0.28	-	75,85,93,96	0
5	GOL	A	417	6/6	0.69	0.31	-	103,118,122,124	0
5	GOL	A	412	6/6	0.74	0.24	-	93,100,106,108	0
5	GOL	A	416	6/6	0.70	0.50	-	69,92,98,98	0
5	GOL	A	414	6/6	0.71	0.53	-	83,88,92,97	0
5	GOL	A	413	6/6	0.59	0.39	-	92,108,120,120	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.