



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:43 PM GMT

PDB ID : 3RUS  
Title : Crystal structure of Cpn-rls in complex with ADP from Methanococcus maripaludis  
Authors : Pereira, J.H.; Ralston, C.Y.; Douglas, N.R.; Kumar, R.; McAndrew, R.P.; Knee, K.M.; King, J.A.; Frydman, J.; Adams, P.D.  
Deposited on : 2011-05-05  
Resolution : 2.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1406 (2.36-2.32)
Clashscore	102246	1509 (2.36-2.32)
Ramachandran outliers	100387	1490 (2.36-2.32)
Sidechain outliers	100360	1491 (2.36-2.32)
RSRZ outliers	91569	1412 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	<div> <div></div> <div>72% 21% • 5%</div> </div>
1	B	543	<div> <div></div> <div>73% 21% • 5%</div> </div>
1	C	543	<div> <div></div> <div>76% 17% • 5%</div> </div>
1	D	543	<div> <div></div> <div>73% 21% • 5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ADP	A	545	-	-	-	X
3	ADP	B	545	-	-	-	X
3	ADP	C	545	-	-	-	X
3	ADP	D	545	-	-	-	X
4	SO4	A	547	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 16140 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chaperonin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	516	Total	C	N	O	S	0	0	0
			3859	2397	666	771	25			
1	B	516	Total	C	N	O	S	0	0	0
			3859	2397	666	771	25			
1	C	516	Total	C	N	O	S	0	0	0
			3859	2397	666	771	25			
1	D	516	Total	C	N	O	S	0	0	0
			3859	2397	666	771	25			


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	327	ALA	THR	ENGINEERED MUTATION	UNP Q877G8
A	328	ALA	ASN	ENGINEERED MUTATION	UNP Q877G8
A	330	ALA	LYS	ENGINEERED MUTATION	UNP Q877G8
A	331	ALA	ASP	ENGINEERED MUTATION	UNP Q877G8
B	327	ALA	THR	ENGINEERED MUTATION	UNP Q877G8
B	328	ALA	ASN	ENGINEERED MUTATION	UNP Q877G8
B	330	ALA	LYS	ENGINEERED MUTATION	UNP Q877G8
B	331	ALA	ASP	ENGINEERED MUTATION	UNP Q877G8
C	327	ALA	THR	ENGINEERED MUTATION	UNP Q877G8
C	328	ALA	ASN	ENGINEERED MUTATION	UNP Q877G8
C	330	ALA	LYS	ENGINEERED MUTATION	UNP Q877G8
C	331	ALA	ASP	ENGINEERED MUTATION	UNP Q877G8
D	327	ALA	THR	ENGINEERED MUTATION	UNP Q877G8
D	328	ALA	ASN	ENGINEERED MUTATION	UNP Q877G8
D	330	ALA	LYS	ENGINEERED MUTATION	UNP Q877G8
D	331	ALA	ASP	ENGINEERED MUTATION	UNP Q877G8

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $\text{C}_{10}\text{H}_{15}\text{N}_5\text{O}_{10}\text{P}_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	B	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	C	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	D	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	163	Total	O	0	0
			163	163		
5	B	128	Total	O	0	0
			128	128		
5	C	136	Total	O	0	0
			136	136		
5	D	125	Total	O	0	0
			125	125		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Chaperonin



#### • Molecule 1: Chaperonin



[illegible]

Chain D: 

Label	Category	Value
G404	Met	1252
V409	Met	1258
S412	Met	1264
R416	Met	1268
R425	Met	1272
A429	Met	1273
A434	Met	1284
R442	Met	1287
A452	Met	1288
I456	Met	1290
V459	Met	1291
A464	Met	1293
S465	Met	1294
M466	Met	1295
G467	Met	1296
C484	Met	1297
K495	Met	1298
Q500	Met	1299
V515	Met	1300
I516	Met	1301
E519	Met	1302
LYS	Met	1303
LEU	Met	1304
ARG	Met	1305
GLY	Met	1306
ALA	Met	1307
PRO	Met	1308
ASP	Met	1309
MET	Met	1310
GLY	Met	1311
ASP	Met	1312
MET	Met	1313
GLY	Met	1314
PRO	Met	1315
GLY	Met	1316
MET	Met	1317
GLY	Met	1318
MET	Met	1319
GLY	Met	1320
PRO	Met	1321
GLY	Met	1322
MET	Met	1323
GLY	Met	1324
MET	Met	1325
Q125	Met	1326
E141	Met	1327
D146	Met	1328
I149	Met	1329
K161	Met	1330
I173	Met	1331
I174	Met	1332
A177	Met	1333
V178	Met	1334
D184	Met	1335
E185	Met	1336
V188	Met	1337
I193	Met	1338
K194	Met	1339
I195	Met	1340
E196	Met	1341
K197	Met	1342
K198	Met	1343
S199	Met	1344
G200	Met	1345
A201	Met	1346
S202	Met	1347
I203	Met	1348
D204	Met	1349
D205	Met	1350
L208	Met	1351
I209	Met	1352
K210	Met	1353
G211	Met	1354
V212	Met	1355
L213	Met	1356
V227	Met	1357
A230	Met	1358
L234	Met	1359
L235	Met	1360
M236	Met	1361
C237	Met	1362
A238	Met	1363
I239	Met	1364
E240	Met	1365
I241	Met	1366
T246	Met	1367
E249	Met	1368
T250	Met	1369
R251	Met	1370
L252	Met	1371
L258	Met	1372
Q264	Met	1373
M288	Met	1374
L269	Met	1375
M272	Met	1376
V273	Met	1377
L284	Met	1378
Q287	Met	1379
K288	Met	1380
G289	Met	1381
I290	Met	1382
D291	Met	1383
D292	Met	1384
L293	Met	1385
A294	Met	1386
Q295	Met	1387
E296	Met	1388
Y297	Met	1389
A305	Met	1390
K311	Met	1391
S312	Met	1392
D313	Met	1393
M314	Met	1394
E315	Met	1395
K316	Met	1396
K319	Met	1397
A327	Met	1398
A328	Met	1399
D339	Met	1400
S352	Met	1401
V356	Met	1402
M367	Met	1403
L368	Met	1404
I369	Met	1405
R370	Met	1406
MET	Met	1407
PRO	Met	1408
GLY	Met	1409
MET	Met	1410
GLY	Met	1411
MET	Met	1412
GLY	Met	1413
MET	Met	1414
GLY	Met	1415
MET	Met	1416
GLY	Met	1417
MET	Met	1418
GLY	Met	1419
MET	Met	1420
GLY	Met	1421
MET	Met	1422
GLY	Met	1423
MET	Met	1424
GLY	Met	1425
MET	Met	1426
GLY	Met	1427
MET	Met	1428
GLY	Met	1429
MET	Met	1430
GLY	Met	1431
MET	Met	1432

## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	161.56Å 185.59Å 185.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.04 – 2.34 49.04 – 2.34	Depositor EDS
% Data completeness (in resolution range)	93.5 (49.04-2.34) 99.9 (49.04-2.34)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.87 (at 2.34Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, $R_{free}$	0.173 , 0.221 0.178 , 0.220	Depositor DCC
$R_{free}$ test set	5866 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.5	Xtriage
Anisotropy	0.158	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 49.1	EDS
Estimated twinning fraction	0.006 for -h,-l,-k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 117127 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	16140	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	0/3883	0.58	1/5228 (0.0%)
1	B	0.39	0/3883	0.54	0/5228
1	C	0.40	0/3883	0.54	0/5228
1	D	0.39	0/3883	0.52	0/5228
All	All	0.40	0/15532	0.55	1/20912 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	78	LEU	CA-CB-CG	5.29	127.46	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3859	0	4001	96	0
1	B	3859	0	4001	102	0
1	C	3859	0	4001	85	0
1	D	3859	0	4001	93	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	27	0	12	4	0
3	B	27	0	12	0	0
3	C	27	0	12	2	0
3	D	27	0	12	1	0
4	A	10	0	0	0	0
4	B	10	0	0	0	0
4	C	10	0	0	0	0
4	D	10	0	0	0	0
5	A	163	0	0	3	0
5	B	128	0	0	1	0
5	C	136	0	0	4	0
5	D	125	0	0	2	0
All	All	16140	0	16052	331	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 331 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:161:LYS:O	5:C:553:HOH:O	1.91	0.87
1:B:477:THR:HG22	1:B:479:ALA:H	1.40	0.87
1:D:141:GLU:HG2	1:D:399:ARG:HH11	1.39	0.87
1:A:61:GLY:HA3	1:A:94:THR:HG22	1.58	0.85
1:B:442:ARG:HG3	1:B:452:ALA:HB1	1.59	0.84

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	514/543 (95%)	505 (98%)	8 (2%)	1 (0%)	52	61
1	B	514/543 (95%)	496 (96%)	17 (3%)	1 (0%)	52	61
1	C	514/543 (95%)	502 (98%)	10 (2%)	2 (0%)	39	45
1	D	514/543 (95%)	501 (98%)	13 (2%)	0	100	100
All	All	2056/2172 (95%)	2004 (98%)	48 (2%)	4 (0%)	52	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	163	ALA
1	C	465	SER
1	C	5	PRO
1	A	465	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	412/430 (96%)	384 (93%)	28 (7%)	20	22
1	B	412/430 (96%)	392 (95%)	20 (5%)	31	38
1	C	412/430 (96%)	397 (96%)	15 (4%)	42	54
1	D	412/430 (96%)	391 (95%)	21 (5%)	29	36
All	All	1648/1720 (96%)	1564 (95%)	84 (5%)	29	36

5 of 84 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	244	THR
1	B	456	LEU
1	D	319	LYS
1	B	250	ILE
1	B	337	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such

sidechains are listed below:

Mol	Chain	Res	Type
1	C	264	GLN
1	C	500	GLN
1	D	287	GLN
1	C	11	ASN
1	D	264	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ADP	A	545	2	22,29,29	0.97	1 (4%)	27,45,45	1.97	5 (18%)
4	SO4	A	546	-	4,4,4	0.14	0	6,6,6	0.15	0
4	SO4	A	547	-	4,4,4	0.24	0	6,6,6	0.13	0
3	ADP	B	545	2	22,29,29	0.97	1 (4%)	27,45,45	1.77	3 (11%)
4	SO4	B	546	-	4,4,4	0.20	0	6,6,6	0.13	0
4	SO4	B	547	-	4,4,4	0.23	0	6,6,6	0.21	0
3	ADP	C	545	2	22,29,29	1.06	2 (9%)	27,45,45	1.91	5 (18%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	C	546	-	4,4,4	0.25	0	6,6,6	0.12	0
4	SO4	C	547	-	4,4,4	0.22	0	6,6,6	0.13	0
3	ADP	D	545	2	22,29,29	0.99	1 (4%)	27,45,45	1.80	4 (14%)
4	SO4	D	546	-	4,4,4	0.26	0	6,6,6	0.15	0
4	SO4	D	547	-	4,4,4	0.22	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	545	2	-	0/12/32/32	0/3/3/3
4	SO4	A	546	-	-	0/0/0/0	0/0/0/0
4	SO4	A	547	-	-	0/0/0/0	0/0/0/0
3	ADP	B	545	2	-	0/12/32/32	0/3/3/3
4	SO4	B	546	-	-	0/0/0/0	0/0/0/0
4	SO4	B	547	-	-	0/0/0/0	0/0/0/0
3	ADP	C	545	2	-	0/12/32/32	0/3/3/3
4	SO4	C	546	-	-	0/0/0/0	0/0/0/0
4	SO4	C	547	-	-	0/0/0/0	0/0/0/0
3	ADP	D	545	2	-	0/12/32/32	0/3/3/3
4	SO4	D	546	-	-	0/0/0/0	0/0/0/0
4	SO4	D	547	-	-	0/0/0/0	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	545	ADP	O4'-C1'	2.06	1.43	1.41
3	A	545	ADP	C5-C4	2.75	1.46	1.40
3	D	545	ADP	C5-C4	2.85	1.46	1.40
3	B	545	ADP	C5-C4	2.97	1.47	1.40
3	C	545	ADP	C5-C4	3.02	1.47	1.40

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	545	ADP	N3-C2-N1	-7.36	123.25	128.89
3	C	545	ADP	N3-C2-N1	-6.70	123.77	128.89
3	D	545	ADP	N3-C2-N1	-6.61	123.83	128.89
3	B	545	ADP	N3-C2-N1	-6.60	123.84	128.89
3	B	545	ADP	C4-C5-N7	-3.38	106.37	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	545	ADP	4	0
3	C	545	ADP	2	0
3	D	545	ADP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	516/543 (95%)	-0.29	1 (0%) 95 98	21, 48, 81, 119	0
1	B	516/543 (95%)	-0.21	6 (1%) 81 88	25, 50, 86, 127	0
1	C	516/543 (95%)	-0.15	7 (1%) 78 85	22, 51, 89, 125	0
1	D	516/543 (95%)	-0.09	7 (1%) 78 85	27, 55, 86, 137	0
All	All	2064/2172 (95%)	-0.19	21 (1%) 84 91	21, 51, 86, 137	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	11	ASN	7.6
1	C	4	GLN	3.8
1	A	11	ASN	3.6
1	B	11	ASN	3.5
1	D	52	LEU	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ADP	B	545	27/27	0.97	0.19	4.57	35,52,71,74	0
3	ADP	A	545	27/27	0.96	0.17	3.92	30,43,52,57	0
3	ADP	D	545	27/27	0.93	0.19	3.14	30,46,69,194	0
3	ADP	C	545	27/27	0.96	0.17	2.89	32,42,55,73	0
4	SO4	A	547	5/5	0.86	0.24	2.08	132,133,134,135	0
4	SO4	D	547	5/5	0.89	0.25	-	130,132,132,132	0
2	MG	A	544	1/1	0.97	0.10	-	30,30,30,30	0
4	SO4	B	547	5/5	0.88	0.23	-	116,116,118,121	0
4	SO4	C	546	5/5	0.95	0.09	-	108,108,109,111	0
4	SO4	A	546	5/5	0.93	0.14	-	89,94,96,99	0
4	SO4	B	546	5/5	0.92	0.10	-	94,98,98,101	0
2	MG	C	544	1/1	0.97	0.13	-	27,27,27,27	0
2	MG	D	544	1/1	0.85	0.15	-	39,39,39,39	0
4	SO4	D	546	5/5	0.89	0.16	-	104,106,110,110	0
2	MG	B	544	1/1	0.93	0.10	-	31,31,31,31	0
4	SO4	C	547	5/5	0.90	0.13	-	121,123,124,125	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.