Full wwPDB NMR Structure Validation Report (i)

Apr 27, 2016 – 05:08 AM BST

PDB ID	:	2RVJ
Title	:	NMR structure of Epithelial splicing regulatory protein 1
Authors	:	Yang, Y.; Allemand, F.; Guichou, J.; Labesse, G.
Deposited on	:	2015-10-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 46%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	104	

2 Ensemble composition and analysis (i)

This entry contains 15 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:443-A:532 (90)	0.42	9

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
2	2, 15

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1638 atoms, of which 816 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Epithelial splicing regulatory protein 1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	104	1638	514	816	150	148	10	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	438	MET	VAL	ENGINEERED MUTATION	UNP Q6NXG1
A	540	LEU	-	EXPRESSION TAG	UNP Q6NXG1
A	541	GLU	-	EXPRESSION TAG	UNP Q6NXG1

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:



4.2.2 Score per residue for model 2

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:

4.2.3 Score per residue for model 3

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A: 79% 8% 13%

M438 P439 P440 T441 N442 A455 R473 R500 K511 K514 D515 R516 M528 L532 M533 G534 G535 T536 L537 N538 R539 L540 E541

4.2.4 Score per residue for model 4

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A: 82% 5% 13%

M438 P439 P440 T441 N442 A455 A456 T457 I458 E459 Q506 K514 M533 G534 G535 T536 L537 N538 R539 L540 E541

4.2.5 Score per residue for model 5

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A: 77% 9% • 13%

M438 P439 P440 T441 N442 R448 L449 R450 A455 I458 E459 D460 I461 L462 D463 R500 K514 Q522 M533 G534 G535 T536 L537 N538 R539 L540 E541

4.2.6 Score per residue for model 6

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A: 81% 5% • 13%

M438 P439 P440 T441 N442 R448 E459 D460 D463 F492 K514 M533 G534 G535 T536 L537 N538 R539 L540 E541

4.2.7 Score per residue for model 7

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A: 78% 9% 13%

M438 P439 P440 T441 N442 C446 I447 A455 L481 Q494 M495 R500 K514 D515 R516 M533 G534 G535 T536 L537 N538 R539 L540 E541

4.2.8 Score per residue for model 8

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:



4.2.9 Score per residue for model 9 (medoid)

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:



4.2.10 Score per residue for model 10

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:



4.2.11 Score per residue for model 11

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:



4.2.12 Score per residue for model 12

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:



4.2.13 Score per residue for model 13

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:

4.2.14 Score per residue for model 14

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:



4.2.15 Score per residue for model 15

- Molecule 1: Epithelial splicing regulatory protein 1

Chain A:

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 15 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	geometry optimization	2.1
CNS	refinement	
CNS, RECOORDscripts	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2rvj_cs.str
Number of chemical shift lists	1
Total number of shifts	591
Number of shifts mapped to atoms	591
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	46%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.3±0.6
All	All	0	5

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	500	ARG	Sidechain	3
1	A	473	ARG	Sidechain	1
1	A	448	ARG	Sidechain	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	719	709	709	2±1
All	All	10785	10635	10635	29

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:447:ILE:HD11	1:A:495:MET:SD	0.61	2.35	12	3
1:A:444:ARG:HB2	1:A:522:GLN:NE2	0.55	2.16	15	1
1:A:458:ILE:O	1:A:462:LEU:HG	0.54	2.02	1	5
1:A:446:CYS:SG	1:A:494:GLN:HG3	0.48	2.48	7	1
1:A:460:ASP:O	1:A:463:ASP:HB3	0.48	2.09	6	2
1:A:446:CYS:SG	1:A:525:ALA:HA	0.47	2.49	9	1
1:A:528:MET:O	1:A:532:LEU:HG	0.47	2.10	10	4
1:A:457:THR:HB	1:A:459:GLU:OE1	0.46	2.10	4	1
1:A:507:LYS:HA	1:A:507:LYS:HE3	0.45	1.88	12	1
1:A:508:CYS:HA	1:A:511:LYS:HB2	0.44	1.90	11	1
1:A:448:ARG:HD3	1:A:492:PHE:CZ	0.43	2.48	6	1
1:A:505:ALA:O	1:A:509:HIS:HB2	0.43	2.14	1	1
1:A:448:ARG:HD3	1:A:450:ARG:NH2	0.43	2.29	5	1
1:A:479:MET:SD	1:A:487:PRO:HB3	0.43	2.53	2	2
1:A:444:ARG:HG3	1:A:524:SER:OG	0.42	2.14	14	1
1:A:480:VAL:O	1:A:487:PRO:HA	0.42	2.14	10	1
1:A:514:LYS:HD2	1:A:514:LYS:N	0.41	2.29	6	1
1:A:449:LEU:O	1:A:490:ASP:HA	0.41	2.16	12	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	90/104 (87%)	85±1 (95±1%)	3±1 (3±2%)	2±1 (2±1%)	15	58
All	All	1350/1560 (87%)	1279 (95%)	47 (3%)	24 (2%)	15	58

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	514	LYS	13
1	A	455	ALA	11

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	76/88 (86%)	74±1 (97±1%)	2±1 (3±1%)	52 90
All	All	1140/1320 (86%)	1106 (97%)	34 (3%)	52 90

All 13 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	522	GLN	6
1	A	500	ARG	5
1	A	495	MET	4
1	A	514	LYS	3
1	A	511	LYS	3
1	A	506	GLN	3
1	A	516	ARG	2
1	A	459	GLU	2
1	A	512	ASN	2
1	A	450	ARG	1
1	A	481	LEU	1
1	A	486	ARG	1
1	A	457	THR	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 46% for the well-defined parts and 45% for the entire structure.

7.1 Chemical shift list 1

File name: 2rvj_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	591
Number of shifts mapped to atoms	591
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	101	-0.36 ± 0.15	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	91	0.22 ± 0.18	None needed (< 0.5 ppm)
$^{13}C'$	97	-0.20 ± 0.20	None needed (< 0.5 ppm)
^{15}N	97	0.12 ± 0.20	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 46%, i.e. 523 atoms were assigned a chemical shift out of a possible 1136. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	441/446 (99%)	177/178 (99%)	177/180 (98%)	87/88 (99%)
Sidechain	82/592 (14%)	0/348 (0%)	82/211 (39%)	0/33 (0%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	0/98 (0%)	0/54 (0%)	0/40 (0%)	0/4 (0%)
Overall	523/1136 (46%)	177/580 (31%)	259/431 (60%)	87/125 (70%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 45%, i.e. 584 atoms were assigned a chemical shift out of a possible 1301. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	493/512 (96%)	198/204 (97%)	198/208 (95%)	97/100 (97%)
Sidechain	91/691 (13%)	0/407 (0%)	91/246 (37%)	0/38 (0%)
Aromatic	0/98 (0%)	0/54 (0%)	0/40 (0%)	0/4 (0%)
Overall	584/1301 (45%)	198/665 (30%)	289/494 (59%)	97/142 (68%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: