



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:05 PM GMT

PDB ID : 1RXC  
Title : E. COLI uridine phosphorylase: 5-fluorouracil ribose-1-phosphate complex  
Authors : Caradoc-Davies, T.T.; Cutfield, S.M.; Lamont, I.L.; Cutfield, J.F.  
Deposited on : 2003-12-18  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	<div> <div>4%</div> <div>85%</div> <div>12%</div> <div>.</div> </div>
1	B	253	<div> <div>%</div> <div>85%</div> <div>13%</div> <div>..</div> </div>
1	C	253	<div> <div>4%</div> <div>84%</div> <div>12%</div> <div>..</div> </div>
1	D	253	<div> <div>4%</div> <div>83%</div> <div>15%</div> <div>..</div> </div>
1	E	253	<div> <div>3%</div> <div>88%</div> <div>10%</div> <div>..</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	253	
1	G	253	
1	H	253	
1	I	253	
1	J	253	
1	K	253	
1	L	253	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	R1P	B	2012	-	-	-	X
2	R1P	C	2082	-	-	-	X
2	R1P	D	2022	-	-	-	X
2	R1P	E	2032	-	-	-	X
2	R1P	F	2002	-	-	-	X
2	R1P	I	2042	-	-	-	X
2	R1P	K	2062	-	-	-	X
2	R1P	L	2072	-	-	-	X

## 2 Entry composition


There are 7 unique types of molecules in this entry. The entry contains 23681 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	246	Total	C	N	O	S	0	0	0
			1847	1158	323	356	10			
1	B	250	Total	C	N	O	S	0	0	0
			1880	1178	328	363	11			
1	C	246	Total	C	N	O	S	0	0	0
			1851	1162	323	355	11			
1	D	250	Total	C	N	O	S	0	0	0
			1880	1178	328	363	11			
1	E	250	Total	C	N	O	S	0	0	0
			1880	1178	328	363	11			
1	F	250	Total	C	N	O	S	0	0	0
			1880	1178	328	363	11			
1	G	245	Total	C	N	O	S	0	0	0
			1839	1153	321	354	11			
1	H	240	Total	C	N	O	S	0	0	0
			1801	1130	315	346	10			
1	I	250	Total	C	N	O	S	0	0	0
			1880	1178	328	363	11			
1	J	250	Total	C	N	O	S	0	0	0
			1880	1178	328	363	11			
1	K	243	Total	C	N	O	S	0	0	0
			1822	1144	319	349	10			
1	L	250	Total	C	N	O	S	0	0	0
			1880	1178	328	363	11			

- Molecule 2 is SUGAR (ALPHA-D-RIBOSE-1-PHOSPHATE) (three-letter code: R1P) (formula: C<sub>5</sub>H<sub>11</sub>O<sub>8</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	F	1	Total	C	O	P	0	0
			14	5	8	1		
2	B	1	Total	C	O	P	0	0
			14	5	8	1		
2	D	1	Total	C	O	P	0	0
			14	5	8	1		
2	E	1	Total	C	O	P	0	0
			14	5	8	1		
2	I	1	Total	C	O	P	0	0
			14	5	8	1		
2	K	1	Total	C	O	P	0	0
			14	5	8	1		
2	L	1	Total	C	O	P	0	0
			14	5	8	1		
2	C	1	Total	C	O	P	0	0
			14	5	8	1		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	J	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	K	1	Total	K	0	0
			1	1		
4	I	1	Total	K	0	0
			1	1		
4	A	1	Total	K	0	0
			1	1		
4	C	1	Total	K	0	0
			1	1		
4	E	1	Total	K	0	0
			1	1		

- Molecule 5 is 5-FLUOROURACIL (three-letter code: URF) (formula: C<sub>4</sub>H<sub>3</sub>FN<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	F	1	Total	C	F	N	O	0	0
			9	4	1	2	2		
5	B	1	Total	C	F	N	O	0	0
			9	4	1	2	2		
5	D	1	Total	C	F	N	O	0	0
			9	4	1	2	2		
5	E	1	Total	C	F	N	O	0	0
			9	4	1	2	2		
5	I	1	Total	C	F	N	O	0	0
			9	4	1	2	2		
5	K	1	Total	C	F	N	O	0	0
			9	4	1	2	2		
5	L	1	Total	C	F	N	O	0	0
			9	4	1	2	2		
5	C	1	Total	C	F	N	O	0	0
			9	4	1	2	2		

- Molecule 6 is 5-FLUOROURIDINE (three-letter code: 5UD) (formula: C<sub>9</sub>H<sub>11</sub>FN<sub>2</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
6	J	1	18	9	1	2	6	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	90	Total	O	0	0
			90	90		
7	B	101	Total	O	0	0
			101	101		
7	C	85	Total	O	0	0
			85	85		
7	D	88	Total	O	0	0
			88	88		
7	E	98	Total	O	0	0
			98	98		
7	F	96	Total	O	0	0
			96	96		
7	G	89	Total	O	0	0
			89	89		
7	H	113	Total	O	0	0
			113	113		
7	I	95	Total	O	0	0
			95	95		
7	J	120	Total	O	0	0
			120	120		
7	K	90	Total	O	0	0
			90	90		

Continued on next page...




*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	L	84	Total	O	0	0
			84	84		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase





- Molecule 1: Uridine phosphorylase









- Molecule 1: Uridine phosphorylase





- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase





- Molecule 1: Uridine phosphorylase









- Molecule 1: Uridine phosphorylase



- Molecule 1: Uridine phosphorylase



- Molecule 1: Uridine phosphorylase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.19Å 191.70Å 91.91Å 90.00° 118.50° 90.00°	Depositor
Resolution (Å)	16.93 – 2.35 16.93 – 2.35	Depositor EDS
% Data completeness (in resolution range)	92.7 (16.93-2.35) 92.7 (16.93-2.35)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.68 (at 2.35Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.150 , 0.203 0.161 , 0.208	Depositor DCC
$R_{free}$ test set	5294 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.7	Xtriage
Anisotropy	0.128	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 39.8	EDS
Estimated twinning fraction	0.002 for -h-l,k,h 0.002 for l,k,-h-l 0.015 for h,-k,-h-l 0.018 for -h-l,-k,l 0.019 for l,-k,h	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 105171 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	23681	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, 5UD, URF, PO4, R1P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/1878	0.74	6/2549 (0.2%)
1	B	0.43	0/1912	0.74	8/2595 (0.3%)
1	C	0.42	0/1882	0.73	4/2552 (0.2%)
1	D	0.42	0/1912	0.75	8/2595 (0.3%)
1	E	0.44	0/1912	0.74	7/2595 (0.3%)
1	F	0.42	0/1912	0.72	4/2595 (0.2%)
1	G	0.41	0/1869	0.73	7/2534 (0.3%)
1	H	0.43	0/1831	0.72	4/2484 (0.2%)
1	I	0.41	0/1912	0.74	5/2595 (0.2%)
1	J	0.44	0/1912	0.76	5/2595 (0.2%)
1	K	0.43	0/1853	0.74	4/2515 (0.2%)
1	L	0.40	0/1912	0.74	6/2595 (0.2%)
All	All	0.42	0/22697	0.74	68/30799 (0.2%)

There are no bond length outliers.

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	27	ASP	CB-CG-OD2	7.44	125.00	118.30
1	A	106	ASP	CB-CG-OD2	7.04	124.64	118.30
1	D	170	ASP	CB-CG-OD2	7.01	124.61	118.30
1	B	117	ASP	CB-CG-OD2	6.80	124.42	118.30
1	D	27	ASP	CB-CG-OD2	6.72	124.35	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1847	0	1845	8	0
1	B	1880	0	1884	8	0
1	C	1851	0	1859	10	0
1	D	1880	0	1884	11	0
1	E	1880	0	1884	6	0
1	F	1880	0	1884	9	0
1	G	1839	0	1846	9	0
1	H	1801	0	1806	12	0
1	I	1880	0	1884	11	0
1	J	1880	0	1884	11	0
1	K	1822	0	1827	13	0
1	L	1880	0	1884	9	0
2	B	14	0	8	0	0
2	C	14	0	8	0	0
2	D	14	0	8	0	0
2	E	14	0	8	0	0
2	F	14	0	8	0	0
2	I	14	0	8	0	0
2	K	14	0	8	0	0
2	L	14	0	8	0	0
3	J	5	0	0	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
4	I	1	0	0	0	0
4	K	1	0	0	0	0
5	B	9	0	3	0	0
5	C	9	0	3	0	0
5	D	9	0	3	0	0
5	E	9	0	3	0	0
5	F	9	0	3	0	0
5	I	9	0	3	0	0
5	K	9	0	3	0	0
5	L	9	0	3	0	0
6	J	18	0	10	0	0
7	A	90	0	0	0	0
7	B	101	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	85	0	0	0	0
7	D	88	0	0	0	0
7	E	98	0	0	0	0
7	F	96	0	0	1	0
7	G	89	0	0	2	0
7	H	113	0	0	2	0
7	I	95	0	0	2	0
7	J	120	0	0	3	0
7	K	90	0	0	1	0
7	L	84	0	0	0	0
All	All	23681	0	22469	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 111 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:185:GLU:HA	1:G:188:GLN:HE21	1.50	0.76
1:C:58:ASP:OD2	1:C:250:ARG:HD3	1.87	0.73
7:G:310:HOH:O	1:I:186:GLU:HG3	1.92	0.68
1:G:152:HIS:HD2	7:G:335:HOH:O	1.80	0.65
1:K:170:ASP:HB2	7:K:1147:HOH:O	1.99	0.63

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/253 (96%)	239 (99%)	2 (1%)	1 (0%)	39	46
1	B	248/253 (98%)	242 (98%)	5 (2%)	1 (0%)	39	46

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	242/253 (96%)	232 (96%)	9 (4%)	1 (0%)	39	46
1	D	248/253 (98%)	241 (97%)	6 (2%)	1 (0%)	39	46
1	E	248/253 (98%)	242 (98%)	5 (2%)	1 (0%)	39	46
1	F	248/253 (98%)	244 (98%)	3 (1%)	1 (0%)	39	46
1	G	241/253 (95%)	235 (98%)	5 (2%)	1 (0%)	39	46
1	H	236/253 (93%)	234 (99%)	1 (0%)	1 (0%)	39	46
1	I	248/253 (98%)	245 (99%)	2 (1%)	1 (0%)	39	46
1	J	248/253 (98%)	245 (99%)	2 (1%)	1 (0%)	39	46
1	K	239/253 (94%)	233 (98%)	5 (2%)	1 (0%)	39	46
1	L	248/253 (98%)	245 (99%)	2 (1%)	1 (0%)	39	46
All	All	2936/3036 (97%)	2877 (98%)	47 (2%)	12 (0%)	39	46

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	163	TYR
1	C	163	TYR
1	F	163	TYR
1	G	163	TYR
1	I	163	TYR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/204 (97%)	188 (95%)	9 (5%)	33	42
1	B	201/204 (98%)	186 (92%)	15 (8%)	17	18
1	C	198/204 (97%)	180 (91%)	18 (9%)	12	11
1	D	201/204 (98%)	188 (94%)	13 (6%)	21	24
1	E	201/204 (98%)	190 (94%)	11 (6%)	27	32
1	F	201/204 (98%)	186 (92%)	15 (8%)	17	18

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	196/204 (96%)	182 (93%)	14 (7%)	18	20
1	H	192/204 (94%)	179 (93%)	13 (7%)	20	22
1	I	201/204 (98%)	192 (96%)	9 (4%)	34	43
1	J	201/204 (98%)	186 (92%)	15 (8%)	17	18
1	K	194/204 (95%)	186 (96%)	8 (4%)	37	48
1	L	201/204 (98%)	185 (92%)	16 (8%)	15	16
All	All	2384/2448 (97%)	2228 (94%)	156 (6%)	21	24

5 of 156 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	92	ILE
1	G	42	VAL
1	L	145	LYS
1	F	150	THR
1	F	240	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	103	ASN
1	G	225	GLN
1	K	188	GLN
1	G	188	GLN
1	H	14	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 23 ligands modelled in this entry, 5 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	URF	B	2011	-	5,9,9	3.54	1 (20%)	5,12,12	6.24	4 (80%)
2	R1P	B	2012	-	13,14,14	0.61	0	18,21,21	4.75	8 (44%)
5	URF	C	2081	-	5,9,9	3.80	1 (20%)	5,12,12	6.41	4 (80%)
2	R1P	C	2082	-	13,14,14	0.66	0	18,21,21	4.95	9 (50%)
5	URF	D	2021	-	5,9,9	3.75	1 (20%)	5,12,12	6.08	4 (80%)
2	R1P	D	2022	-	13,14,14	0.66	0	18,21,21	4.60	9 (50%)
5	URF	E	2031	-	5,9,9	3.39	1 (20%)	5,12,12	6.92	4 (80%)
2	R1P	E	2032	-	13,14,14	0.55	0	18,21,21	4.92	10 (55%)
5	URF	F	2001	-	5,9,9	3.23	1 (20%)	5,12,12	6.52	4 (80%)
2	R1P	F	2002	-	13,14,14	0.68	0	18,21,21	4.91	8 (44%)
5	URF	I	2041	-	5,9,9	3.83	1 (20%)	5,12,12	6.55	5 (100%)
2	R1P	I	2042	-	13,14,14	0.78	0	18,21,21	4.78	8 (44%)
6	5UD	J	2051	-	14,19,19	2.40	2 (14%)	13,28,28	4.17	11 (84%)
3	PO4	J	2052	-	4,4,4	0.45	0	6,6,6	0.28	0
5	URF	K	2061	-	5,9,9	3.16	1 (20%)	5,12,12	6.77	4 (80%)
2	R1P	K	2062	-	13,14,14	0.65	0	18,21,21	4.71	9 (50%)
5	URF	L	2071	-	5,9,9	3.41	1 (20%)	5,12,12	6.30	4 (80%)
2	R1P	L	2072	-	13,14,14	0.65	0	18,21,21	4.53	9 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	URF	B	2011	-	-	0/0/0/0	0/1/1/1
2	R1P	B	2012	-	-	0/6/23/23	0/1/1/1
5	URF	C	2081	-	-	0/0/0/0	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R1P	C	2082	-	-	0/6/23/23	0/1/1/1
5	URF	D	2021	-	-	0/0/0/0	0/1/1/1
2	R1P	D	2022	-	-	0/6/23/23	0/1/1/1
5	URF	E	2031	-	-	0/0/0/0	0/1/1/1
2	R1P	E	2032	-	-	0/6/23/23	0/1/1/1
5	URF	F	2001	-	-	0/0/0/0	0/1/1/1
2	R1P	F	2002	-	-	0/6/23/23	0/1/1/1
5	URF	I	2041	-	-	0/0/0/0	0/1/1/1
2	R1P	I	2042	-	-	0/6/23/23	0/1/1/1
6	5UD	J	2051	-	-	0/2/22/22	0/2/2/2
3	PO4	J	2052	-	-	0/0/0/0	0/0/0/0
5	URF	K	2061	-	-	0/0/0/0	0/1/1/1
2	R1P	K	2062	-	-	0/6/23/23	0/1/1/1
5	URF	L	2071	-	-	0/0/0/0	0/1/1/1
2	R1P	L	2072	-	-	0/6/23/23	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	J	2051	5UD	C6-C5	2.39	1.42	1.38
5	K	2061	URF	C4-C5	6.99	1.47	1.38
5	F	2001	URF	C4-C5	7.07	1.47	1.38
5	L	2071	URF	C4-C5	7.57	1.47	1.38
5	E	2031	URF	C4-C5	7.58	1.47	1.38

The worst 5 of 114 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2032	R1P	O4'-C1'-C2'	-12.75	86.87	104.78
2	F	2002	R1P	O4'-C1'-C2'	-12.65	87.01	104.78
2	C	2082	R1P	O4'-C1'-C2'	-12.26	87.55	104.78
5	E	2031	URF	N1-C2-N3	-12.15	120.58	128.33
5	K	2061	URF	N1-C2-N3	-12.05	120.65	128.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	246/253 (97%)	-0.01	9 (3%) 45 59	11, 19, 36, 63	0
1	B	250/253 (98%)	-0.25	3 (1%) 81 89	10, 18, 29, 35	0
1	C	246/253 (97%)	-0.06	10 (4%) 41 55	11, 19, 38, 66	0
1	D	250/253 (98%)	-0.11	9 (3%) 46 60	11, 20, 44, 63	0
1	E	250/253 (98%)	-0.21	7 (2%) 56 69	11, 17, 32, 50	0
1	F	250/253 (98%)	-0.23	11 (4%) 38 52	10, 15, 31, 58	0
1	G	245/253 (96%)	-0.03	9 (3%) 45 59	15, 23, 35, 62	0
1	H	240/253 (94%)	-0.22	5 (2%) 67 79	12, 19, 29, 42	0
1	I	250/253 (98%)	-0.25	5 (2%) 68 79	10, 17, 31, 51	0
1	J	250/253 (98%)	-0.37	3 (1%) 81 89	9, 15, 25, 41	0
1	K	243/253 (96%)	-0.24	5 (2%) 67 79	9, 16, 28, 39	0
1	L	250/253 (98%)	-0.07	11 (4%) 38 52	15, 21, 35, 55	0
All	All	2970/3036 (97%)	-0.17	87 (2%) 55 67	9, 19, 33, 66	0

The worst 5 of 87 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	228	ILE	7.1
1	G	232	GLU	6.0
1	F	232	GLU	5.9
1	A	230	ASN	5.6
1	L	233	THR	5.3

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	R1P	L	2072	14/14	0.93	0.19	4.64	38,38,41,42	2
2	R1P	D	2022	14/14	0.88	0.20	3.63	38,39,43,43	3
2	R1P	K	2062	14/14	0.85	0.21	3.41	31,32,39,40	2
2	R1P	C	2082	14/14	0.80	0.24	3.11	42,44,50,50	2
2	R1P	I	2042	14/14	0.95	0.15	2.82	27,29,33,33	2
2	R1P	E	2032	14/14	0.91	0.17	2.73	31,34,39,39	2
2	R1P	F	2002	14/14	0.94	0.15	2.66	30,32,38,39	2
2	R1P	B	2012	14/14	0.94	0.15	2.58	26,29,30,30	2
6	5UD	J	2051	18/18	0.91	0.16	1.76	29,31,33,35	2
3	PO4	J	2052	5/5	0.98	0.10	0.48	18,18,19,19	0
5	URF	K	2061	9/9	0.95	0.10	0.20	19,21,22,27	0
5	URF	C	2081	9/9	0.92	0.16	0.17	41,44,44,45	1
5	URF	B	2011	9/9	0.96	0.10	-0.38	11,13,15,17	0
5	URF	D	2021	9/9	0.95	0.10	-0.47	30,32,33,33	0
5	URF	F	2001	9/9	0.96	0.10	-0.63	18,22,23,25	0
5	URF	E	2031	9/9	0.97	0.09	-0.72	25,27,28,29	0
5	URF	L	2071	9/9	0.97	0.10	-0.76	27,28,29,32	0
5	URF	I	2041	9/9	0.97	0.07	-1.27	19,20,21,22	0
4	K	C	2102	1/1	0.99	0.05	-2.41	21,21,21,21	0
4	K	E	2103	1/1	0.99	0.03	-2.65	16,16,16,16	0
4	K	I	2104	1/1	1.00	0.03	-3.06	16,16,16,16	0
4	K	K	2105	1/1	0.99	0.03	-3.14	17,17,17,17	0
4	K	A	2101	1/1	0.99	0.03	-3.21	20,20,20,20	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.