



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:09 PM GMT

PDB ID : 1S0V  
Title : Structural basis for substrate selection by T7 RNA polymerase  
Authors : Temiakov, D.; Patlan, V.; Anikin, M.; McAllister, W.T.; Yokoyama, S.; Vassilyev, D.G.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2004-01-05  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)
RNA backbone	2183	1079 (3.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	18	<div> <div>11%</div> <div>39%</div> <div>11%</div> <div>33%</div> <div>6%</div> </div>
1	H	18	<div> <div>6%</div> <div>17%</div> <div>28%</div> <div>22%</div> <div>28%</div> <div>6%</div> </div>
1	K	18	<div> <div>56%</div> <div>11%</div> <div>28%</div> <div>6%</div> </div>
1	N	18	<div> <div>6%</div> <div>56%</div> <div>22%</div> <div>11%</div> <div>6%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	12	
2	I	12	
2	L	12	
2	O	12	
3	G	10	
3	J	10	
3	M	10	
3	P	10	
4	A	883	
4	B	883	
4	C	883	
4	D	883	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	APC	A	2000	-	-	X	-
6	APC	B	2001	-	-	X	-
6	APC	C	2002	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 30899 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*G\*GP\*GP\*AP\*AP\*TP\*CP\*GP\*AP\*TP\*AP\*TP\*CP\*GP\*CP\*CP\*GP\*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	17	Total	C	N	O	P	0	0	0
			346	165	66	99	16			
1	H	17	Total	C	N	O	P	0	0	0
			346	165	66	99	16			
1	K	17	Total	C	N	O	P	0	0	0
			346	165	66	99	16			
1	N	17	Total	C	N	O	P	0	0	0
			346	165	66	99	16			

- Molecule 2 is a RNA chain called 5'-R(\*AP\*AP\*CP\*U\*GP\*CP\*GP\*GP\*CP\*GP\*AP\*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	8	Total	C	N	O	P	0	0	0
			171	77	33	54	7			
2	I	8	Total	C	N	O	P	0	0	0
			171	77	33	54	7			
2	L	8	Total	C	N	O	P	0	0	0
			171	77	33	54	7			
2	O	8	Total	C	N	O	P	0	0	0
			171	77	33	54	7			

- Molecule 3 is a DNA chain called 5'-D(\*GP\*TP\*CP\*GP\*AP\*TP\*TP\*CP\*CP\*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	9	Total	C	N	O	P	0	0	0
			179	87	30	54	8			
3	J	9	Total	C	N	O	P	0	0	0
			179	87	30	54	8			
3	M	9	Total	C	N	O	P	0	0	0
			179	87	30	54	8			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	9	Total	C	N	O	P	0	0	0
			179	87	30	54	8			

- Molecule 4 is a protein called DNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	857	Total	C	N	O	S	0	0	0
			6746	4296	1173	1242	35			
4	B	857	Total	C	N	O	S	0	0	0
			6746	4296	1173	1242	35			
4	C	857	Total	C	N	O	S	0	0	0
			6746	4296	1173	1242	35			
4	D	857	Total	C	N	O	S	0	0	0
			6746	4296	1173	1242	35			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	497	LEU	-	INSERTION	UNP P00573
B	497	LEU	-	INSERTION	UNP P00573
C	497	LEU	-	INSERTION	UNP P00573
D	497	LEU	-	INSERTION	UNP P00573

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Mg	0	0
			2	2		
5	A	1	Total	Mg	0	0
			1	1		
5	D	2	Total	Mg	0	0
			2	2		
5	C	2	Total	Mg	0	0
			2	2		
5	F	1	Total	Mg	0	0
			1	1		

- Molecule 6 is DIPHOSPHOMETHYLPHOSPHONIC ACID ADENOSYL ESTER (three-letter code: APC) (formula: C<sub>11</sub>H<sub>18</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
6	B	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
6	C	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
6	D	1	Total	C	N	O	P	0	0
			31	11	5	12	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	237	Total	O	0	0
			237	237		
7	B	212	Total	O	0	0
			212	212		
7	C	201	Total	O	0	0
			201	201		
7	D	173	Total	O	0	0
			173	173		
7	E	39	Total	O	0	0
			39	39		
7	F	9	Total	O	0	0
			9	9		
7	G	9	Total	O	0	0
			9	9		
7	H	19	Total	O	0	0
			19	19		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	I	14	Total 14	O 14	0	0
7	J	13	Total 13	O 13	0	0
7	K	20	Total 20	O 20	0	0
7	L	8	Total 8	O 8	0	0
7	M	10	Total 10	O 10	0	0
7	N	14	Total 14	O 14	0	0
7	O	15	Total 15	O 15	0	0
7	P	6	Total 6	O 6	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(\*G\*GP\*GP\*AP\*AP\*TP\*CP\*GP\*AP\*TP\*AP\*TP\*CP\*GP\*CP\*CP\*GP\*C)-3'

Chain E: 




- Molecule 1: 5'-D(\*G\*GP\*GP\*AP\*AP\*TP\*CP\*GP\*AP\*TP\*AP\*TP\*CP\*GP\*CP\*CP\*GP\*C)-3'

Chain H: 




- Molecule 1: 5'-D(\*G\*GP\*GP\*AP\*AP\*TP\*CP\*GP\*AP\*TP\*AP\*TP\*CP\*GP\*CP\*CP\*GP\*C)-3'

Chain K: 



- Molecule 1: 5'-D(\*G\*GP\*GP\*AP\*AP\*TP\*CP\*GP\*AP\*TP\*AP\*TP\*CP\*GP\*CP\*CP\*GP\*C)-3'

Chain N: 



- Molecule 2: 5'-R(\*AP\*AP\*CP\*U\*GP\*CP\*GP\*GP\*CP\*GP\*AP\*U)-3'


Chain F: 



- Molecule 2: 5'-R(\*AP\*AP\*CP\*U\*GP\*CP\*GP\*GP\*CP\*GP\*AP\*U)-3'




Chain I: 



- Molecule 2: 5'-R(\*AP\*AP\*CP\*U\*GP\*CP\*GP\*GP\*CP\*GP\*AP\*U)-3'

Chain L: 





- Molecule 2: 5'-R(\*AP\*AP\*CP\*U\*GP\*CP\*GP\*GP\*CP\*GP\*AP\*U)-3'

Chain O: 





- Molecule 3: 5'-D(\*GP\*TP\*CP\*GP\*AP\*TP\*TP\*CP\*CP\*C)-3'

Chain G: 




- Molecule 3: 5'-D(\*GP\*TP\*CP\*GP\*AP\*TP\*TP\*CP\*CP\*C)-3'

Chain J: 





- Molecule 3: 5'-D(\*GP\*TP\*CP\*GP\*AP\*TP\*TP\*CP\*CP\*C)-3'

Chain M: 




- Molecule 3: 5'-D(\*GP\*TP\*CP\*GP\*AP\*TP\*TP\*CP\*CP\*C)-3'


Chain P: 





- Molecule 4: DNA-directed RNA polymerase

Chain A: 







### • Molecule 4: DNA-directed RNA polymerase







## • Molecule 4: DNA-directed RNA polymerase



A843	L778	G716	Q649	A575	A513	L452	A390	N325	L261	S201
Y847	A779	E717	V650	K576	F514	G453	A391	T326	A261	S202
D846	P780	I718	L651	K577	C515	K454	R391	G263	G263	S203
Q848	N781	L719	E652	V578	F516	E455	R394	W328	W204	W204
F849	F782	K720	D653	N579	E517	G456	R395	K329	S265	H205
A850	V783	K721	T654	E580	Y518	Y457	R396	I330	H206	H206
D851	H784	R722	L655	T581	A519	Y458	I396	I331	E207	E207
Q852	S785	G723	Q656	L582	G520	W459	S397	K332	D208	D208
D853	Q786	A724	Q657	Q583	V521	L460	I398	K333	S209	S209
H854	D787	H725	A584	D585	Q522	E399	F400	V334	C271	H211
S855	G788	H726	D585	D585	H523	F400	I335	I210	H211	H211
S856	S789	W727	D660	N588	H524	W401	I402	V272	V212	V212
Q857	H730	W728	S661	G589	G525	L402	V337	V273	G213	G213
L858	L791	T729	K663	G590	L526	E403	A338	P274	V214	V214
D859	R792	P730	G664	T590	S527	Q404	N339	P275	R215	R215
K860	K793	D731	L665	Y528	Y528	A405	V340	K276	C216	C216
M861	T794	G732	M666	N592	N529	I406	I341	P277	I217	I217
P862	W795	F733	F667	E593	C530	W470	T342	W278	E218	E218
A863	W796	P734	T668	V594	S531	D471	K343	I281	W219	W219
L864	W797	W735	Q669	M601	L532	K472	I344	I282	L220	L220
P865	K801	W736	Q672	T602	P533	V473	K412	G283	I221	I221
G868	Y802	E738	M677	G603	A535	F475	I414	E283	E222	E222
N869	G803	Y739	Y676	S606	F536	P476	W415	W286	T224	T224
L870	I804	K741	A678	W609	D537	E477	F416	W287	G225	G225
N871	E885	P742	K679	L611	G538	R478	I417	A288	M226	M226
R873	S806	I743	L680	L611	S539	I479	Y418	M289	V227	V227
D874	L809	Q744	I681	G612	S541	F481	K480	G290	S228	S228
I875	T810	T745	G683	T513	G542	I482	I419	R291	L229	L229
L876	D812	R746	S684	K614	I543	E483	W420	R292	H230	H230
S877	S813	N748	V685	A615	Q544	E484	Q422	P293	R231	R231
S878	F814	L749	S686	L616	Q546	N485	R423	L294	Q232	Q232
D879	G815	W750	V687	Q619	N549	N486	Q424	A295	N233	N233
F880	T816	F751	T688	W620	L550	E487	Y426	L296	A234	A234
A881	L817	L752	V689	L621	R551	I489	Y427	R298	W236	W236
F882	P818	G753	A691	A622	D552	M490	A428	H300	V237	V237
A883	A819	Q754	A692	W623	E553	A491	V429	E301	G238	G238
	D820	F755	V693	V625	V554	C492		Q239	Q239	Q239
	N823	R756	M693	W625	G555	K494	F432	D240	D240	D240
	L824	GLN	M696	V629	G556	S495	I433	K302	S241	S241
	F825	PRO	M697	T630	R557	P496	P434	K303	E242	E242
	K826	THR	W698	R631	A558	E498	Q435	A304	T243	T243
	A827	ILE	L699	K632	V559	E498	I437	I305	I244	I244
	V828	ASN	K700	R632	N560	N499	D438	N306	E245	E245
	R829	THR	S701	S633	L561	T500	I439	Y308	L246	L246
	E830	ASN	A702	V634	L562	W501	N439	E309	A247	A247
	T831	LYS	A703	W635	P563	W502	T440	D310	P248	P248
	M832	D766	K704	W635	S564	A503	I441	V311	E249	E249
	V833		L705	W639		E504	G442	Y312	Y250	Y250
	D834	I769	L706	Q640	V567	E504	L443	K313	A251	A251
	T835		S641	S641	Q505	Q505	L444	E252	E252	E252
	Y836	H772	E709	K642	Q568	D506	K378	E315	A253	A253
	E837	K773	K711	E643	D569	S507	I446	I254	I254	I254
	D840	Q774	W710	P646	I570	P508	A447	Y317	A255	A255
	V841	E775	D712	F647	W571	F509	K448	T256	T256	T256
	L842	W776	T715	Q648	G572	C510	Q449	R257	R257	R257
		G777			I573	F511	K450	A258	A258	A258
					V574	L512	P451	G259	G259	G259
								Q324	Q324	Q324

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.13Å 87.70Å 206.53Å 91.93° 91.02° 110.66°	Depositor
Resolution (Å)	40.00 – 3.20 39.88 – 3.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-3.20) 82.3 (39.88-3.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 3.01Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.255 , 0.307 0.255 , 0.305	Depositor DCC
$R_{free}$ test set	3685 reflections (5.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	59.0	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 107.2	EDS
Estimated twinning fraction	0.014 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.37$	Xtriage
Outliers	0 of 88725 reflections	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	30899	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: APC, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	E	1.74	9/388 (2.3%)	1.60	9/597 (1.5%)
1	H	1.62	7/388 (1.8%)	1.47	6/597 (1.0%)
1	K	1.59	7/388 (1.8%)	1.31	1/597 (0.2%)
1	N	1.26	3/388 (0.8%)	1.27	4/597 (0.7%)
2	F	2.39	9/191 (4.7%)	2.00	11/297 (3.7%)
2	I	2.25	7/191 (3.7%)	1.68	2/297 (0.7%)
2	L	1.65	1/191 (0.5%)	1.36	0/297
2	O	1.47	0/191	1.38	0/297
3	G	0.82	0/199	0.92	0/305
3	J	0.78	0/199	0.93	0/305
3	M	0.88	0/199	0.86	0/305
3	P	0.94	0/199	1.07	0/305
4	A	1.20	22/6897 (0.3%)	1.14	24/9329 (0.3%)
4	B	1.21	24/6897 (0.3%)	1.14	24/9329 (0.3%)
4	C	0.97	5/6897 (0.1%)	0.97	7/9329 (0.1%)
4	D	0.92	3/6897 (0.0%)	0.91	4/9329 (0.0%)
All	All	1.14	97/30700 (0.3%)	1.09	92/42112 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	6
1	H	0	9
1	K	0	6
1	N	0	4
2	F	0	4
2	I	0	2
2	L	0	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	#Chirality outliers	#Planarity outliers
2	O	0	3
3	P	0	1
4	A	0	1
4	B	0	1
4	D	0	1
All	All	0	39

The worst 5 of 97 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	502	TRP	CB-CG	-13.18	1.26	1.50
1	K	14	DG	C5-C6	-11.28	1.31	1.42
4	A	467	CYS	CB-SG	-10.63	1.64	1.82
2	I	8	U	N1-C6	-9.75	1.29	1.38
2	I	7	A	C5-C6	-8.93	1.33	1.41

The worst 5 of 92 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	460	LEU	CB-CG-CD1	-10.01	93.98	111.00
1	E	14	DG	O5'-P-OP1	-9.17	97.45	105.70
4	A	460	LEU	CB-CG-CD2	9.03	126.35	111.00
4	D	791	LEU	CA-CB-CG	8.79	135.52	115.30
4	A	425	ARG	NE-CZ-NH1	-8.45	116.07	120.30

There are no chirality outliers.

5 of 39 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	11	DA	Sidechain
1	E	12	DT	Sidechain
1	E	13	DC	Sidechain
1	E	14	DG	Sidechain
1	E	16	DC	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	346	0	190	28	0
1	H	346	0	192	66	0
1	K	346	0	190	57	0
1	N	346	0	192	38	0
2	F	171	0	89	10	0
2	I	171	0	89	11	0
2	L	171	0	89	14	0
2	O	171	0	89	5	0
3	G	179	0	104	13	0
3	J	179	0	104	17	0
3	M	179	0	104	13	0
3	P	179	0	104	8	0
4	A	6746	0	6708	1151	0
4	B	6746	0	6708	1190	0
4	C	6746	0	6708	1007	0
4	D	6746	0	6708	899	0
5	A	1	0	0	0	0
5	B	2	0	0	0	0
5	C	2	0	0	0	0
5	D	2	0	0	0	0
5	F	1	0	0	0	0
6	A	31	0	13	16	0
6	B	31	0	13	9	0
6	C	31	0	14	16	0
6	D	31	0	13	7	0
7	A	237	0	0	82	0
7	B	212	0	0	68	0
7	C	201	0	0	70	0
7	D	173	0	0	53	0
7	E	39	0	0	1	0
7	F	9	0	0	1	0
7	G	9	0	0	1	0
7	H	19	0	0	9	0
7	I	14	0	0	4	0
7	J	13	0	0	0	0
7	K	20	0	0	7	0
7	L	8	0	0	0	0
7	M	10	0	0	2	0
7	N	14	0	0	3	0
7	O	15	0	0	2	0
7	P	6	0	0	2	0
All	All	30899	0	28421	4458	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 76.

The worst 5 of 4458 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:313:MET:SD	4:A:313:MET:CE	2.05	1.44
4:C:631:LYS:NZ	6:C:2002:APC:H3A2	1.52	1.22
2:F:1:G:H5"	7:F:3008:HOH:O	1.40	1.19
4:A:631:LYS:NZ	6:A:2000:APC:H3A2	1.58	1.18
4:A:546:PHE:CE1	4:A:783:VAL:HG22	1.78	1.16

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	A	851/883 (96%)	570 (67%)	198 (23%)	83 (10%)	1	4
4	B	851/883 (96%)	578 (68%)	188 (22%)	85 (10%)	1	4
4	C	851/883 (96%)	608 (71%)	178 (21%)	65 (8%)	1	9
4	D	851/883 (96%)	623 (73%)	174 (20%)	54 (6%)	2	13
All	All	3404/3532 (96%)	2379 (70%)	738 (22%)	287 (8%)	1	6

5 of 287 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	A	7	ALA
4	A	194	GLY
4	A	199	GLU
4	A	281	ILE
4	A	288	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	A	703/729 (96%)	573 (82%)	130 (18%)	2	10
4	B	703/729 (96%)	573 (82%)	130 (18%)	2	10
4	C	703/729 (96%)	594 (84%)	109 (16%)	3	15
4	D	703/729 (96%)	598 (85%)	105 (15%)	4	17
All	All	2812/2916 (96%)	2338 (83%)	474 (17%)	2	13

5 of 474 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	B	569	ASP
4	C	32	LEU
4	D	571	TYR
4	B	633	SER
4	B	776	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 125 such sidechains are listed below:

Mol	Chain	Res	Type
4	B	545	HIS
4	B	848	GLN
4	D	726	HIS
4	B	588	ASN
4	B	748	ASN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	F	7/12 (58%)	0	0
2	I	7/12 (58%)	1 (14%)	0
2	L	7/12 (58%)	0	0
2	O	7/12 (58%)	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
All	All	28/48 (58%)	1 (3%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	I	8	U

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	APC	A	2000	5	25,33,33	1.95	5 (20%)	30,52,52	1.85	9 (30%)
6	APC	B	2001	5	25,33,33	2.13	7 (28%)	30,52,52	2.14	6 (20%)
6	APC	C	2002	5	25,33,33	2.23	7 (28%)	30,52,52	1.35	5 (16%)
6	APC	D	2003	5	25,33,33	2.00	4 (16%)	30,52,52	1.85	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	APC	A	2000	5	-	0/15/38/38	0/3/3/3
6	APC	B	2001	5	-	0/15/38/38	0/3/3/3
6	APC	C	2002	5	-	0/15/38/38	0/3/3/3
6	APC	D	2003	5	-	0/15/38/38	0/3/3/3

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	2001	APC	PB-O2B	-2.76	1.49	1.56
6	B	2001	APC	PA-O2A	-2.50	1.50	1.56
6	C	2002	APC	PA-O2A	-2.47	1.50	1.56
6	A	2000	APC	C8-N7	-2.24	1.30	1.34
6	A	2000	APC	PB-O2B	-2.24	1.50	1.56

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	2001	APC	O1A-PA-C3A	-4.68	97.24	109.02
6	A	2000	APC	O5'-PA-O1A	-4.59	101.79	113.98
6	B	2001	APC	PG-O3B-PB	-3.67	120.37	132.67
6	A	2000	APC	PG-O3B-PB	-2.88	123.02	132.67
6	C	2002	APC	PG-O3B-PB	-2.43	124.51	132.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 48 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	2000	APC	16	0
6	B	2001	APC	9	0
6	C	2002	APC	16	0
6	D	2003	APC	7	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	E	17/18 (94%)	-0.20	0 100 100	19, 58, 179, 181	0
1	H	17/18 (94%)	0.20	1 (5%) 26 14	21, 75, 158, 171	0
1	K	17/18 (94%)	0.14	0 100 100	48, 94, 157, 159	0
1	N	17/18 (94%)	0.15	0 100 100	51, 102, 172, 173	0
2	F	8/12 (66%)	-0.46	0 100 100	17, 23, 83, 93	0
2	I	8/12 (66%)	-0.01	0 100 100	17, 38, 82, 93	0
2	L	8/12 (66%)	0.21	0 100 100	47, 53, 113, 118	0
2	O	8/12 (66%)	0.16	0 100 100	51, 80, 132, 144	0
3	G	9/10 (90%)	1.06	2 (22%) 1 1	146, 153, 176, 183	0
3	J	9/10 (90%)	0.07	1 (11%) 7 4	130, 138, 162, 163	0
3	M	9/10 (90%)	-0.13	0 100 100	134, 146, 155, 158	0
3	P	9/10 (90%)	1.39	1 (11%) 7 4	157, 167, 172, 172	0
4	A	857/883 (97%)	-0.28	37 (4%) 39 25	15, 76, 141, 153	0
4	B	857/883 (97%)	-0.41	15 (1%) 71 58	10, 69, 131, 150	0
4	C	857/883 (97%)	-0.30	19 (2%) 65 50	47, 93, 134, 160	0
4	D	857/883 (97%)	-0.04	29 (3%) 49 34	41, 102, 144, 156	0
All	All	3564/3692 (96%)	-0.24	105 (2%) 55 41	10, 89, 142, 183	0

The worst 5 of 105 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	P	1	DG	8.5
4	D	235	GLY	7.1
3	G	1	DG	6.9
4	A	235	GLY	6.2
4	A	601	ASN	5.5

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
6	APC	B	2001	31/31	0.94	0.22	1.24	28,47,63,68	0
6	APC	A	2000	31/31	0.95	0.20	1.19	34,58,80,82	0
6	APC	C	2002	31/31	0.93	0.24	0.47	65,70,77,78	0
6	APC	D	2003	31/31	0.94	0.17	-0.82	48,57,61,63	0
5	MG	D	3004	1/1	0.70	0.31	-	79,79,79,79	0
5	MG	B	3002	1/1	0.95	0.55	-	65,65,65,65	0
5	MG	C	3007	1/1	0.99	0.18	-	46,46,46,46	0
5	MG	F	3005	1/1	0.96	0.16	-	18,18,18,18	0
5	MG	C	3003	1/1	0.99	0.27	-	41,41,41,41	0
5	MG	D	3008	1/1	0.97	0.18	-	16,16,16,16	0
5	MG	B	3006	1/1	0.98	0.17	-	34,34,34,34	0
5	MG	A	3001	1/1	0.94	0.13	-	22,22,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.