



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:09 PM GMT

PDB ID : 1SA1
Title : Tubulin-podophyllotoxin: stathmin-like domain complex
Authors : Ravelli, R.B.; Gigant, B.; Curmi, P.A.; Jourdain, I.; Lachkar, S.; Sobel, A.; Knossow, M.
Deposited on : 2004-02-06
Resolution : 4.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1140 (4.80-3.60)
Ramachandran outliers	100387	1083 (4.80-3.60)
Sidechain outliers	100360	1067 (4.80-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	451	
1	C	451	
2	B	445	
2	D	445	
3	E	142	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	POD	B	700	X	-	-	-
7	POD	D	701	X	-	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14180 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	433	Total	C	N	O	S	0	0	0
			3299	2089	559	631	20			
1	C	430	Total	C	N	O	S	0	0	0
			3275	2072	555	628	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	265	ILE	ALA	SEE REMARK 999	UNP P02550
C	265	ILE	ALA	SEE REMARK 999	UNP P02550

- Molecule 2 is a protein called Tubulin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	422	Total	C	N	O	S	0	0	0
			3241	2040	545	631	25			
2	D	426	Total	C	N	O	S	0	0	0
			3278	2059	556	638	25			

- Molecule 3 is a protein called Stathmin 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	134	Total	C	N	O	S	0	0	0
			905	555	169	176	5			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	4	ALA	-	SEE REMARK 999	UNP P02554

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	C	1	Total Mg 1 1	0	0

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



- [illegible]

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			30	22	8		



Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	D	1	30	22	8	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: Tubulin alpha chain




- Molecule 1: Tubulin alpha chain






• Molecule 2: Tubulin beta chain


Chain B: 24% 44% 21% 6% 5%




• Molecule 2: Tubulin beta chain

Chain D: 26% 44% 20% 6%





● Molecule 3: Stathmin 4



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	328.06 Å 328.06 Å 54.30 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 4.20	Depositor
% Data completeness (in resolution range)	98.3 (20.00-4.20)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.204 , 0.259	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	14180	wwPDB-VP
Average B, all atoms (Å ²)	116.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, GTP, MG, POD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.89	1/3374 (0.0%)	1.09	19/4593 (0.4%)
1	C	0.74	0/3349	1.00	11/4561 (0.2%)
2	B	0.86	3/3314 (0.1%)	1.08	25/4506 (0.6%)
2	D	0.76	2/3352 (0.1%)	1.04	20/4556 (0.4%)
3	E	0.87	0/914	0.95	2/1238 (0.2%)
All	All	0.82	6/14303 (0.0%)	1.05	77/19454 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	C	0	3
2	B	0	3
2	D	0	2
3	E	0	3
All	All	0	15

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	44	LEU	C-N	11.14	1.59	1.34
2	B	44	LEU	C-N	9.19	1.55	1.34
2	B	360	PRO	C-N	8.02	1.52	1.34
2	D	360	PRO	C-N	6.55	1.49	1.34
2	B	2	ARG	NE-CZ	5.14	1.39	1.33

The worst 5 of 77 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	427	ASP	CB-CG-OD2	8.82	126.24	118.30
2	D	205	ASP	CB-CG-OD2	8.78	126.20	118.30
2	D	163	ASP	CB-CG-OD2	8.20	125.68	118.30
2	B	205	ASP	CB-CG-OD2	7.81	125.33	118.30
1	C	211	ASP	CB-CG-OD2	7.70	125.22	118.30

There are no chirality outliers.

5 of 15 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	146	GLY	Peptide
1	A	220	GLU	Peptide
1	A	221	ARG	Peptide
1	A	339	ARG	Peptide
2	B	49	ILE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3299	0	3129	275	0
1	C	3275	0	3104	271	0
2	B	3241	0	3033	321	0
2	D	3278	0	3074	297	0
3	E	905	0	730	59	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	32	0	12	3	0
5	C	32	0	12	2	0
6	B	28	0	12	1	0
6	D	28	0	12	2	0
7	B	30	0	19	5	0
7	D	30	0	19	4	0
All	All	14180	0	13156	1177	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 1177 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:33:PRO:HB3	3:E:35:PHE:CZ	1.64	1.33
1:C:240:ALA:HB2	1:C:243:ARG:NH1	1.39	1.33
2:D:273:ALA:CB	2:D:274:PRO:HD3	1.74	1.18
2:B:191:VAL:HG11	2:B:425:MET:HE3	1.22	1.15
2:D:191:VAL:HG11	2:D:425:MET:HE3	1.22	1.14

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	429/451 (95%)	288 (67%)	88 (20%)	53 (12%)	0	8
1	C	426/451 (94%)	289 (68%)	90 (21%)	47 (11%)	0	11
2	B	418/445 (94%)	274 (66%)	87 (21%)	57 (14%)	0	6
2	D	424/445 (95%)	276 (65%)	87 (20%)	61 (14%)	0	6
3	E	130/142 (92%)	81 (62%)	29 (22%)	20 (15%)	0	5
All	All	1827/1934 (94%)	1208 (66%)	381 (21%)	238 (13%)	0	7

5 of 238 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	33	ASP
1	A	47	ASP
1	A	62	VAL
1	A	73	THR
1	A	96	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/378 (90%)	217 (64%)	125 (36%)	0	1
1	C	340/378 (90%)	219 (64%)	121 (36%)	0	1
2	B	343/381 (90%)	216 (63%)	127 (37%)	0	1
2	D	348/381 (91%)	214 (62%)	134 (38%)	0	0
3	E	65/126 (52%)	45 (69%)	20 (31%)	0	3
All	All	1438/1644 (88%)	911 (63%)	527 (37%)	0	1

5 of 527 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	382	THR
1	C	178	SER
2	D	373	MET
2	B	406	HIS
1	C	92	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	406	HIS
1	C	133	GLN
2	D	309	HIS
1	C	8	HIS
1	C	88	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GTP	A	600	4	25,34,34	1.08	1 (4%)	34,54,54	1.95	8 (23%)
6	GDP	B	602	-	23,30,30	1.14	2 (8%)	30,47,47	2.30	10 (33%)
7	POD	B	700	-	34,34,34	3.20	11 (32%)	51,51,51	4.21	21 (41%)
5	GTP	C	601	4	25,34,34	1.13	2 (8%)	34,54,54	1.97	9 (26%)
6	GDP	D	603	-	23,30,30	0.96	1 (4%)	30,47,47	1.95	8 (26%)
7	POD	D	701	-	34,34,34	3.19	11 (32%)	51,51,51	3.98	22 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	A	600	4	-	0/18/38/38	0/3/3/3
6	GDP	B	602	-	-	0/12/32/32	0/3/3/3
7	POD	B	700	-	1/1/5/5	0/10/45/45	0/5/5/5
5	GTP	C	601	4	-	0/18/38/38	0/3/3/3
6	GDP	D	603	-	-	0/12/32/32	0/3/3/3
7	POD	D	701	-	1/1/5/5	0/10/45/45	0/5/5/5

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	D	701	POD	C7-C6	-10.33	1.21	1.39
7	D	701	POD	C7-C2	-9.32	1.21	1.38
7	B	700	POD	C7-C6	-9.19	1.23	1.39
7	B	700	POD	C7-C2	-8.85	1.22	1.38
7	B	700	POD	C10-C13	-6.33	1.42	1.51

The worst 5 of 78 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	701	POD	C7-C6-C5	-12.43	103.81	120.11
7	B	700	POD	C7-C6-C5	-12.11	104.22	120.11
7	B	700	POD	C7-C2-C3	-10.53	107.77	121.96
7	D	701	POD	C7-C2-C3	-9.27	109.47	121.96
6	B	602	GDP	N3-C2-N1	-5.96	118.36	127.44

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
7	D	701	POD	C10
7	B	700	POD	C10

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	600	GTP	3	0
6	B	602	GDP	1	0
7	B	700	POD	5	0
5	C	601	GTP	2	0
6	D	603	GDP	2	0
7	D	701	POD	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.