



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:59 PM GMT

PDB ID : 3SBU
Title : Crystal structure of a ntf2-like protein (BF2862) from Bacteroides fragilis
NCTC 9343 at 2.15 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2011-06-06
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	<div> <div>11%</div> <div>72%</div> <div>16%</div> <div>12%</div> </div>
1	B	261	<div> <div>13%</div> <div>77%</div> <div>11%</div> <div>12%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	285	-	-	-	X
2	GOL	A	287	-	-	-	X
2	GOL	A	288	-	-	-	X
2	GOL	A	289	-	-	X	-
2	GOL	A	290	-	-	-	X
2	GOL	A	291	-	-	-	X
2	GOL	B	286	-	-	-	X
3	PEG	A	293	-	-	-	X
3	PEG	A	294	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4167 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical ntf2-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	Se	0	4	0
			1947	1265	309	370	3			
1	B	230	Total	C	N	O	Se	0	12	0
			1986	1291	314	378	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	leader sequence	UNP Q5LBG4
B	0	GLY	-	leader sequence	UNP Q5LBG4

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	79	Total 79	O 79	0	0
4	B	79	Total 79	O 79	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hypothetical ntf2-like protein



• Molecule 1: Hypothetical ntf2-like protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	80.48 Å 80.48 Å 196.10 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.59 – 2.15 29.59 – 2.15	Depositor EDS
% Data completeness (in resolution range)	(Not available) (29.59-2.15) 99.9 (29.59-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.16 Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.176 , 0.192 0.182 , 0.203	Depositor DCC
R_{free} test set	1951 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	46.0	Xtriage
Anisotropy	0.128	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 60.9	EDS
Estimated twinning fraction	0.064 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 38909 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4167	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG, MLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	1/1870 (0.1%)	0.61	0/2541
1	B	0.49	0/1936	0.63	0/2629
All	All	0.51	1/3806 (0.0%)	0.62	0/5170

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	124	MSE	SE-CE	-6.27	1.58	1.95

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1947	0	1824	30	0
1	B	1986	0	1864	16	0
2	A	42	0	56	9	0
2	B	6	0	8	0	0
3	A	21	0	30	2	0
3	B	7	0	10	0	0
4	A	79	0	0	1	0
4	B	79	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4167	0	3792	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:192:SER:HB2	1:B:219[B]:LEU:HD21	1.65	0.76
1:A:64:PHE:H	2:A:289:GOL:H12	1.55	0.71
1:A:85:PRO:HB2	1:A:96:MLY:HG3	1.73	0.69
1:A:115:THR:HG22	1:A:254:LEU:HD23	1.74	0.67
1:A:64:PHE:N	2:A:289:GOL:H12	2.14	0.62
1:B:115:THR:HG22	1:B:254:LEU:HD23	1.80	0.62
1:A:86:LEU:HB3	1:A:97:ILE:HG23	1.84	0.59
1:A:139:TRP:CE2	1:A:233:MLY:HH23	2.38	0.58
1:B:196:ARG:HH21	1:B:270:MLY:HH23	1.70	0.56
1:B:203:THR:HG22	1:B:281:ASP:HB3	1.86	0.55
1:B:280[B]:GLU:HG2	1:B:282:THR:HG23	1.88	0.54
1:B:77:LEU:O	1:B:81:ARG:HG2	2.09	0.53
1:A:167:ARG:HE	2:A:291:GOL:H12	1.74	0.53
1:B:143:MLY:HH23	1:B:186:THR:HG21	1.92	0.52
1:A:91:GLY:O	2:A:291:GOL:H2	2.11	0.51
1:B:87:PRO:HG3	1:B:96:MLY:HH11	1.92	0.51
1:A:198:PRO:HB2	1:A:216:THR:HB	1.93	0.50
1:B:198:PRO:HB2	1:B:216:THR:HB	1.92	0.50
1:A:147:VAL:HG21	2:A:285:GOL:H12	1.94	0.49
1:A:230:PRO:HB3	3:A:293:PEG:H22	1.96	0.48
1:A:249:HIS:HE1	4:A:338:HOH:O	1.97	0.47
1:A:127:VAL:HG11	1:A:255:MLY:HH11	1.96	0.46
1:A:204:THR:CG2	1:A:209:ASP:HA	2.45	0.46
1:A:202:VAL:HB	1:A:280[A]:GLU:HG2	1.96	0.45
1:A:78:GLN:O	1:A:82:VAL:HG23	2.17	0.45
1:A:129:ASP:O	2:A:289:GOL:H32	2.18	0.43
1:A:77:LEU:HD11	1:A:81:ARG:HH11	1.82	0.43
1:A:266:TYR:HB2	1:A:278:MLY:HB2	2.01	0.43
1:A:230:PRO:HG2	1:A:234:LEU:HD23	2.02	0.42
1:A:215:THR:HG21	1:B:168:PRO:HD3	2.01	0.42
1:A:255:MLY:HH22	2:A:290:GOL:H2	2.02	0.42
1:A:148:MLY:HH13	1:A:150:TYR:OH	2.20	0.42
1:A:77:LEU:O	1:A:81:ARG:HG2	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:TRP:CE2	1:B:233:MLY:HH23	2.55	0.42
1:B:127:VAL:HG11	1:B:255:MLY:HH11	2.01	0.42
1:A:191:GLN:OE1	1:A:226:MLY:HH12	2.20	0.41
1:A:63:SER:HB2	2:A:289:GOL:H11	2.02	0.41
1:A:111:GLN:HG2	3:A:293:PEG:H42	2.03	0.41
1:A:139:TRP:CE3	1:A:233:MLY:HD3	2.55	0.41
1:A:207:ASP:OD2	1:B:148:MLY:HH22	2.21	0.41
1:B:202:VAL:O	1:B:280[B]:GLU:HA	2.21	0.41
1:B:246[B]:ASN:ND2	4:B:335:HOH:O	2.54	0.41
1:A:167:ARG:NE	2:A:291:GOL:H12	2.35	0.40
1:B:96:MLY:HG3	1:B:96:MLY:HH12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/261 (83%)	215 (99%)	2 (1%)	0	100	100
1	B	225/261 (86%)	222 (99%)	3 (1%)	0	100	100
All	All	442/522 (85%)	437 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/214 (91%)	193 (99%)	2 (1%)	82	87
1	B	201/214 (94%)	199 (99%)	2 (1%)	82	87
All	All	396/428 (92%)	392 (99%)	4 (1%)	80	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	PHE
1	A	243	ASN
1	B	180	PHE
1	B	243	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	243	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

30 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	MLY	A	100	1	3,4,11	0.59	0	0,4,13	0.00	-
1	MLY	A	103	1	3,4,11	0.65	0	0,4,13	0.00	-
1	MLY	A	110	1	4,5,11	0.61	0	3,5,13	1.53	1 (33%)
1	MLY	A	143	1	8,9,11	1.42	1 (12%)	7,9,13	1.22	1 (14%)
1	MLY	A	144	1	9,10,11	1.11	1 (11%)	9,11,13	1.60	3 (33%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MLY	A	148	1	9,10,11	1.17	1 (11%)	9,11,13	1.51	3 (33%)
1	MLY	A	149	1	9,10,11	1.05	1 (11%)	9,11,13	1.88	4 (44%)
1	MLY	A	156	1	7,8,11	0.54	0	6,8,13	1.14	0
1	MLY	A	226	1	9,10,11	1.08	1 (11%)	9,11,13	1.56	2 (22%)
1	MLY	A	233	1	9,10,11	1.34	1 (11%)	9,11,13	1.02	0
1	MLY	A	255	1	9,10,11	0.78	0	9,11,13	1.46	2 (22%)
1	MLY	A	270	1	9,10,11	0.71	0	9,11,13	1.67	3 (33%)
1	MLY	A	278	1	9,10,11	1.00	0	9,11,13	1.44	3 (33%)
1	MLY	A	56	1	3,4,11	0.61	0	0,4,13	0.00	-
1	MLY	A	96	1	9,10,11	0.86	0	9,11,13	1.68	2 (22%)
1	MLY	B	100	1	3,4,11	0.70	0	0,4,13	0.00	-
1	MLY	B	103	1	3,4,11	0.73	0	0,4,13	0.00	-
1	MLY	B	110	1	9,10,11	0.59	0	9,11,13	1.16	1 (11%)
1	MLY	B	143	1	9,10,11	1.07	0	9,11,13	1.36	2 (22%)
1	MLY	B	144	1	3,4,11	0.63	0	0,4,13	0.00	-
1	MLY	B	148	1	9,10,11	0.87	0	9,11,13	1.77	2 (22%)
1	MLY	B	149	1	9,10,11	0.82	0	9,11,13	2.30	5 (55%)
1	MLY	B	156	1	3,4,11	0.74	0	0,4,13	0.00	-
1	MLY	B	226	1	9,10,11	1.13	0	9,11,13	1.34	1 (11%)
1	MLY	B	233	1	9,10,11	0.96	1 (11%)	9,11,13	2.08	2 (22%)
1	MLY	B	255	1	9,10,11	0.95	0	9,11,13	1.48	2 (22%)
1	MLY	B	270	1	9,10,11	0.85	0	9,11,13	1.35	1 (11%)
1	MLY	B	278	1	9,10,11	0.97	1 (11%)	9,11,13	1.27	1 (11%)
1	MLY	B	56	1	3,4,11	0.64	0	0,4,13	0.00	-
1	MLY	B	96	1	9,10,11	0.85	0	9,11,13	1.09	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	A	100	1	-	0/0/2/11	0/0/0/0
1	MLY	A	103	1	-	0/0/2/11	0/0/0/0
1	MLY	A	110	1	-	0/2/4/11	0/0/0/0
1	MLY	A	143	1	-	0/6/8/11	0/0/0/0
1	MLY	A	144	1	-	0/7/9/11	0/0/0/0
1	MLY	A	148	1	-	0/7/9/11	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	A	149	1	-	0/7/9/11	0/0/0/0
1	MLY	A	156	1	-	0/5/7/11	0/0/0/0
1	MLY	A	226	1	-	0/7/9/11	0/0/0/0
1	MLY	A	233	1	-	0/7/9/11	0/0/0/0
1	MLY	A	255	1	-	0/7/9/11	0/0/0/0
1	MLY	A	270	1	-	0/7/9/11	0/0/0/0
1	MLY	A	278	1	-	0/7/9/11	0/0/0/0
1	MLY	A	56	1	-	0/0/2/11	0/0/0/0
1	MLY	A	96	1	-	0/7/9/11	0/0/0/0
1	MLY	B	100	1	-	0/0/2/11	0/0/0/0
1	MLY	B	103	1	-	0/0/2/11	0/0/0/0
1	MLY	B	110	1	-	0/7/9/11	0/0/0/0
1	MLY	B	143	1	-	0/7/9/11	0/0/0/0
1	MLY	B	144	1	-	0/0/2/11	0/0/0/0
1	MLY	B	148	1	-	0/7/9/11	0/0/0/0
1	MLY	B	149	1	-	0/7/9/11	0/0/0/0
1	MLY	B	156	1	-	0/0/2/11	0/0/0/0
1	MLY	B	226	1	-	0/7/9/11	0/0/0/0
1	MLY	B	233	1	-	0/7/9/11	0/0/0/0
1	MLY	B	255	1	-	0/7/9/11	0/0/0/0
1	MLY	B	270	1	-	0/7/9/11	0/0/0/0
1	MLY	B	278	1	-	0/7/9/11	0/0/0/0
1	MLY	B	56	1	-	0/0/2/11	0/0/0/0
1	MLY	B	96	1	-	0/7/9/11	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	148	MLY	CB-CA	-3.07	1.50	1.53
1	A	149	MLY	CB-CA	-2.61	1.51	1.53
1	B	278	MLY	CH2-NZ	2.12	1.53	1.46
1	B	233	MLY	CB-CA	2.45	1.56	1.53
1	A	226	MLY	CB-CA	2.58	1.56	1.53
1	A	144	MLY	CH2-NZ	2.69	1.54	1.46
1	A	143	MLY	CB-CA	3.51	1.57	1.53
1	A	233	MLY	CB-CA	3.63	1.57	1.53

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	233	MLY	CB-CA-N	-4.71	97.13	110.52
1	B	149	MLY	CH1-NZ-CE	-4.29	93.72	110.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	148	MLY	CH2-NZ-CH1	-3.67	99.91	109.72
1	A	226	MLY	CH2-NZ-CH1	-3.30	100.91	109.72
1	A	255	MLY	CD-CE-NZ	-3.16	105.79	113.92
1	A	149	MLY	CD-CG-CB	-3.00	102.99	113.66
1	A	149	MLY	CB-CA-N	-2.99	102.03	110.52
1	B	255	MLY	CD-CE-NZ	-2.95	106.33	113.92
1	B	149	MLY	CD-CG-CB	-2.86	103.51	113.66
1	A	270	MLY	CH2-NZ-CE	-2.72	99.98	110.79
1	A	96	MLY	O-C-CA	-2.60	118.72	125.49
1	B	148	MLY	O-C-CA	-2.57	118.79	125.49
1	A	144	MLY	O-C-CA	-2.55	118.85	125.49
1	B	270	MLY	O-C-CA	-2.51	118.95	125.49
1	B	143	MLY	O-C-CA	-2.51	118.95	125.49
1	B	233	MLY	CD-CG-CB	-2.50	104.78	113.66
1	A	278	MLY	CD-CG-CB	-2.49	104.81	113.66
1	A	96	MLY	CB-CA-N	-2.45	103.57	110.52
1	B	255	MLY	O-C-CA	-2.43	119.16	125.49
1	A	270	MLY	O-C-CA	-2.41	119.21	125.49
1	A	148	MLY	CD-CG-CB	-2.40	105.14	113.66
1	A	144	MLY	CH2-NZ-CH1	-2.39	103.32	109.72
1	A	110	MLY	O-C-CA	-2.39	119.27	125.49
1	A	148	MLY	CH1-NZ-CE	-2.38	101.33	110.79
1	A	270	MLY	CH2-NZ-CH1	-2.38	103.37	109.72
1	B	110	MLY	O-C-CA	-2.35	119.36	125.49
1	A	143	MLY	O-C-CA	-2.34	119.39	125.49
1	B	143	MLY	CH2-NZ-CH1	-2.32	103.50	109.72
1	B	149	MLY	O-C-CA	-2.25	119.63	125.49
1	B	278	MLY	CD-CG-CB	-2.21	105.80	113.66
1	B	96	MLY	O-C-CA	-2.21	119.74	125.49
1	B	149	MLY	CH2-NZ-CH1	-2.20	103.84	109.72
1	A	255	MLY	CG-CB-CA	-2.14	105.18	114.16
1	A	144	MLY	CD-CG-CB	-2.12	106.14	113.66
1	A	149	MLY	O-C-CA	-2.08	120.06	125.49
1	A	148	MLY	O-C-CA	-2.08	120.08	125.49
1	A	278	MLY	O-C-CA	-2.02	120.22	125.49
1	A	278	MLY	CD-CE-NZ	-2.01	108.74	113.92
1	A	149	MLY	CH2-NZ-CH1	2.02	115.12	109.72
1	B	149	MLY	CD-CE-NZ	2.09	119.30	113.92
1	A	226	MLY	CB-CA-N	2.15	116.63	110.52
1	B	226	MLY	CH2-NZ-CE	2.62	121.22	110.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

12 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	148	MLY	1	0
1	A	226	MLY	1	0
1	A	233	MLY	2	0
1	A	255	MLY	2	0
1	A	278	MLY	1	0
1	A	96	MLY	1	0
1	B	143	MLY	1	0
1	B	148	MLY	1	0
1	B	233	MLY	1	0
1	B	255	MLY	1	0
1	B	270	MLY	1	0
1	B	96	MLY	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	285	-	5,5,5	0.96	0	5,5,5	0.97	0
2	GOL	A	287	-	5,5,5	0.72	0	5,5,5	0.69	0
2	GOL	A	288	-	5,5,5	0.73	0	5,5,5	0.73	0
2	GOL	A	289	-	5,5,5	0.60	0	5,5,5	1.55	1 (20%)
2	GOL	A	290	-	5,5,5	0.64	0	5,5,5	0.80	0
2	GOL	A	291	-	5,5,5	0.56	0	5,5,5	1.18	0
2	GOL	A	292	-	5,5,5	0.50	0	5,5,5	0.79	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PEG	A	293	-	6,6,6	0.35	0	5,5,5	0.33	0
3	PEG	A	294	-	6,6,6	0.58	0	5,5,5	0.41	0
3	PEG	A	295	-	6,6,6	0.16	0	5,5,5	0.14	0
2	GOL	B	286	-	5,5,5	0.55	0	5,5,5	0.83	0
3	PEG	B	296	-	6,6,6	0.67	0	5,5,5	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	285	-	-	0/4/4/4	0/0/0/0
2	GOL	A	287	-	-	0/4/4/4	0/0/0/0
2	GOL	A	288	-	-	0/4/4/4	0/0/0/0
2	GOL	A	289	-	-	0/4/4/4	0/0/0/0
2	GOL	A	290	-	-	0/4/4/4	0/0/0/0
2	GOL	A	291	-	-	0/4/4/4	0/0/0/0
2	GOL	A	292	-	-	0/4/4/4	0/0/0/0
3	PEG	A	293	-	-	0/4/4/4	0/0/0/0
3	PEG	A	294	-	-	0/4/4/4	0/0/0/0
3	PEG	A	295	-	-	0/4/4/4	0/0/0/0
2	GOL	B	286	-	-	0/4/4/4	0/0/0/0
3	PEG	B	296	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	289	GOL	O2-C2-C3	2.45	119.89	108.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	285	GOL	1	0
2	A	289	GOL	4	0
2	A	290	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	291	GOL	3	0
3	A	293	PEG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/261 (81%)	0.65	28 (13%) 4 7	33, 60, 101, 130	0
1	B	212/261 (81%)	0.79	33 (15%) 3 4	32, 61, 126, 155	0
All	All	424/522 (81%)	0.72	61 (14%) 3 5	32, 61, 115, 155	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	101	TYR	11.9
1	B	101	TYR	11.3
1	A	210	PHE	7.0
1	B	55	PRO	6.1
1	B	59	GLU	5.9
1	B	158	ALA	5.9
1	A	55	PRO	5.7
1	B	60	ALA	5.6
1	B	57	PRO	5.5
1	A	57	PRO	5.3
1	B	85	PRO	5.1
1	B	97	ILE	5.0
1	A	284	ILE	4.7
1	A	155	ILE	4.6
1	A	60	ALA	4.5
1	A	69	TYR	4.4
1	B	157	GLY	4.4
1	A	102	TRP	3.9
1	A	62[A]	GLU	3.9
1	B	98	ASP	3.7
1	B	155	ILE	3.7
1	B	137	VAL	3.7
1	A	58	ILE	3.6
1	B	271	ASP	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	252	LEU	3.4
1	B	284	ILE	3.4
1	B	69	TYR	3.2
1	B	84	PHE	3.2
1	B	252	LEU	3.0
1	A	234	LEU	3.0
1	B	258	GLY	3.0
1	A	140	ILE	3.0
1	B	62	GLU	2.9
1	A	147	VAL	2.9
1	A	237	ILE	2.8
1	A	116	LEU	2.8
1	B	210	PHE	2.8
1	B	99	ARG	2.7
1	A	117	LEU	2.7
1	B	58	ILE	2.6
1	A	85	PRO	2.6
1	A	137	VAL	2.6
1	A	251	ILE	2.5
1	B	66[A]	ASP	2.5
1	B	237	ILE	2.4
1	B	117[A]	LEU	2.3
1	B	102	TRP	2.3
1	A	158	ALA	2.3
1	A	154	ARG	2.2
1	A	157	GLY	2.2
1	A	173	GLU	2.2
1	B	223	PHE	2.2
1	B	150	TYR	2.2
1	A	59	GLU	2.1
1	B	76	ALA	2.1
1	B	265	LEU	2.1
1	B	80	GLN	2.1
1	B	88[A]	TYR	2.1
1	B	147	VAL	2.1
1	A	189	ILE	2.0
1	A	68	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	MLY	B	233	11/12	0.98	0.12	-	54,59,66,67	0
1	MLY	A	96	11/12	0.87	0.26	-	95,105,108,109	0
1	MLY	B	143	11/12	0.88	0.28	-	65,74,86,89	0
1	MLY	B	56	5/12	0.86	0.33	-	153,156,158,162	0
1	MLY	B	103	5/12	0.95	0.12	-	100,104,105,106	0
1	MLY	B	255	11/12	0.96	0.11	-	45,51,65,65	0
1	MLY	B	110	11/12	0.91	0.35	-	81,87,104,107	0
1	MLY	B	270	11/12	0.85	0.41	-	58,70,79,81	0
1	MLY	A	100	5/12	0.86	0.38	-	119,122,125,126	0
1	MLY	A	156	9/12	0.92	0.36	-	93,100,112,115	0
1	MLY	B	149	11/12	0.96	0.18	-	56,59,63,63	0
1	MLY	A	233	11/12	0.95	0.12	-	51,54,62,62	0
1	MLY	A	148	11/12	0.97	0.21	-	48,55,69,71	0
1	MLY	A	103	5/12	0.94	0.13	-	91,94,96,96	0
1	MLY	B	278	11/12	0.98	0.16	-	35,36,36,38	0
1	MLY	B	96	11/12	0.90	0.28	-	101,108,111,113	0
1	MLY	A	255	11/12	0.96	0.10	-	47,52,63,64	0
1	MLY	A	144	11/12	0.95	0.09	-	46,47,52,55	0
1	MLY	A	270	11/12	0.94	0.15	-	45,54,69,70	0
1	MLY	A	149	11/12	0.92	0.21	-	55,60,66,67	0
1	MLY	A	110	6/12	0.93	0.27	-	81,83,85,89	0
1	MLY	B	156	5/12	0.91	0.30	-	105,109,114,115	0
1	MLY	A	226	11/12	0.94	0.13	-	52,53,67,69	0
1	MLY	B	148	11/12	0.94	0.22	-	52,57,69,71	0
1	MLY	A	56	5/12	0.86	0.35	-	128,130,131,135	0
1	MLY	B	100	5/12	0.89	0.43	-	127,129,130,133	0
1	MLY	A	143	10/12	0.92	0.14	-	49,56,72,72	0
1	MLY	B	226	11/12	0.94	0.12	-	49,51,65,68	0
1	MLY	A	278	11/12	0.97	0.12	-	34,35,38,39	0
1	MLY	B	144	5/12	0.97	0.09	-	66,66,67,69	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	A	288	6/6	0.79	0.20	4.45	96,96,97,97	0
2	GOL	A	291	6/6	0.74	0.32	4.38	94,95,95,96	0
2	GOL	A	285	6/6	0.76	0.44	4.34	81,83,84,84	0
3	PEG	A	293	7/7	0.52	0.39	3.70	95,97,98,99	0
2	GOL	A	287	6/6	0.76	0.19	2.61	73,76,77,77	0
2	GOL	A	290	6/6	0.62	0.28	2.41	83,85,86,87	0
2	GOL	B	286	6/6	0.55	0.24	2.07	90,91,91,92	0
3	PEG	A	294	7/7	0.45	0.28	2.02	96,97,98,99	0
2	GOL	A	289	6/6	0.91	0.23	0.54	76,76,77,78	0
2	GOL	A	292	6/6	0.85	0.37	-	90,92,93,93	0
3	PEG	A	295	7/7	0.79	0.34	-	101,102,105,105	0
3	PEG	B	296	7/7	0.81	0.19	-	83,85,86,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.