



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:00 PM GMT

PDB ID : 3SG6
Title : Crystal Structure of Dimeric GCaMP2-LIA(linker 1)
Authors : Schreiter, E.R.; Akerboom, J.; Looger, L.L.
Deposited on : 2011-06-14
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called myosin light chain kinase, green fluorescent protein, calmodulin chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	S	0	5	0
			3082	1942	517	608	15			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP Q6LDG3
A	2	ARG	-	EXPRESSION TAG	UNP Q6LDG3
A	3	GLY	-	EXPRESSION TAG	UNP Q6LDG3
A	4	SER	-	EXPRESSION TAG	UNP Q6LDG3
A	5	HIS	-	EXPRESSION TAG	UNP Q6LDG3
A	6	HIS	-	EXPRESSION TAG	UNP Q6LDG3
A	7	HIS	-	EXPRESSION TAG	UNP Q6LDG3
A	8	HIS	-	EXPRESSION TAG	UNP Q6LDG3
A	9	HIS	-	EXPRESSION TAG	UNP Q6LDG3
A	10	HIS	-	EXPRESSION TAG	UNP Q6LDG3
A	11	GLY	-	EXPRESSION TAG	UNP Q6LDG3
A	12	MET	-	EXPRESSION TAG	UNP Q6LDG3
A	13	ALA	-	EXPRESSION TAG	UNP Q6LDG3
A	14	SER	-	EXPRESSION TAG	UNP Q6LDG3
A	15	MET	-	EXPRESSION TAG	UNP Q6LDG3
A	16	THR	-	EXPRESSION TAG	UNP Q6LDG3
A	17	GLY	-	EXPRESSION TAG	UNP Q6LDG3
A	18	GLY	-	EXPRESSION TAG	UNP Q6LDG3
A	19	GLN	-	EXPRESSION TAG	UNP Q6LDG3
A	20	GLN	-	EXPRESSION TAG	UNP Q6LDG3
A	21	MET	-	EXPRESSION TAG	UNP Q6LDG3
A	22	GLY	-	EXPRESSION TAG	UNP Q6LDG3
A	23	ARG	-	EXPRESSION TAG	UNP Q6LDG3
A	24	ASP	-	EXPRESSION TAG	UNP Q6LDG3
A	25	LEU	-	EXPRESSION TAG	UNP Q6LDG3
A	26	TYR	-	EXPRESSION TAG	UNP Q6LDG3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	ASP	-	EXPRESSION TAG	UNP Q6LDG3
A	28	ASP	-	EXPRESSION TAG	UNP Q6LDG3
A	29	ASP	-	EXPRESSION TAG	UNP Q6LDG3
A	30	ASP	-	EXPRESSION TAG	UNP Q6LDG3
A	31	LYS	-	EXPRESSION TAG	UNP Q6LDG3
A	32	ASP	-	EXPRESSION TAG	UNP Q6LDG3
A	33	LEU	-	EXPRESSION TAG	UNP Q6LDG3
A	34	ALA	-	EXPRESSION TAG	UNP Q6LDG3
A	35	THR	-	EXPRESSION TAG	UNP Q6LDG3
A	36	MET	-	EXPRESSION TAG	UNP Q6LDG3
A	37	VAL	-	EXPRESSION TAG	UNP Q6LDG3
A	38	ASP	-	EXPRESSION TAG	UNP Q6LDG3
A	39	SER	-	EXPRESSION TAG	UNP Q6LDG3
A	40	SER	-	EXPRESSION TAG	UNP Q6LDG3
A	45	ASN	GLN	ENGINEERED MUTATION	UNP Q6LDG3
A	60	LEU	-	LINKER	UNP P42212
A	61	ILE	-	LINKER	UNP P42212
A	62	ALA	-	LINKER	UNP P42212
A	77	ALA	VAL	ENGINEERED MUTATION	UNP P42212
A	89	GLY	SER	ENGINEERED MUTATION	UNP P42212
A	94	TYR	ASP	ENGINEERED MUTATION	UNP P42212
A	120	LYS	ALA	ENGINEERED MUTATION	UNP P42212
A	145	LEU	HIS	ENGINEERED MUTATION	UNP P42212
A	153	GLY	-	LINKER	UNP P42212
A	154	GLY	-	LINKER	UNP P42212
A	155	THR	-	LINKER	UNP P42212
A	156	GLY	-	LINKER	UNP P42212
A	157	GLY	-	LINKER	UNP P42212
A	158	SER	-	LINKER	UNP P42212
A	159	MET	-	LINKER	UNP P42212
A	160	VAL	-	LINKER	UNP P42212
A	223	LEU	PHE	ENGINEERED MUTATION	UNP P42212
A	225	CRO	SER	CHROMOPHORE	UNP P42212
A	225	CRO	TYR	CHROMOPHORE	UNP P42212
A	225	CRO	GLY	CHROMOPHORE	UNP P42212
A	252	ILE	VAL	ENGINEERED MUTATION	UNP P42212
A	304	THR	-	LINKER	UNP P62161
A	305	ARG	-	LINKER	UNP P62161


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total 4	Ca 4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	272	Total 272	O 272	0	0

- Molecule 1: myosin light chain kinase, green fluorescent protein, calmodulin chimera



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	128.98 Å 47.49 Å 68.67 Å 90.00° 99.72° 90.00°	Depositor
Resolution (Å)	27.74 – 1.70 27.36 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.8 (27.74-1.70) 93.9 (27.36-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 1.70 Å)	Xtriage
Refinement program	REFMAC 5.6.0111	Depositor
R, R_{free}	0.203 , 0.249 0.208 , 0.249	Depositor DCC
R_{free} test set	2147 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 53.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	0 of 42585 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3358	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	7/3127 (0.2%)	1.02	10/4210 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	44	TRP	CD2-CE2	5.88	1.48	1.41
1	A	371	GLU	CB-CG	5.66	1.62	1.52
1	A	371	GLU	CG-CD	5.48	1.60	1.51
1	A	301	GLU	CB-CG	-5.30	1.42	1.52
1	A	251	TYR	CD1-CE1	5.27	1.47	1.39
1	A	116	SER	CB-OG	-5.25	1.35	1.42
1	A	335	GLU	CG-CD	5.12	1.59	1.51

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	268	ARG	NE-CZ-NH2	-9.07	115.76	120.30
1	A	371	GLU	OE1-CD-OE2	-7.75	114.00	123.30
1	A	268	ARG	NE-CZ-NH1	7.63	124.11	120.30
1	A	255	ARG	NE-CZ-NH1	7.01	123.81	120.30
1	A	326	ASP	CB-CG-OD1	6.58	124.22	118.30
1	A	241	ASP	CB-CG-OD1	6.35	124.02	118.30
1	A	178	ASP	CB-CG-OD2	6.19	123.87	118.30
1	A	116	SER	CB-CA-C	5.23	120.04	110.10
1	A	394	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	A	256	THR	CA-CB-CG2	-5.09	105.28	112.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3082	0	3001	38	1
2	A	4	0	0	1	0
3	A	272	0	0	12	1
All	All	3358	0	3001	39	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:449:MET:HE2	3:A:670:HOH:O	1.74	0.86
1:A:309:THR:HG23	1:A:312:GLN:HB2	1.56	0.85
1:A:309:THR:HG21	1:A:311:GLU:OE1	1.76	0.85
1:A:67[B]:MET:HE3	1:A:76:LYS:NZ	1.93	0.84
2:A:455:CA:CA	3:A:567:HOH:O	1.57	0.81
1:A:448:MET:HG3	3:A:670:HOH:O	1.82	0.80
1:A:413:MET:HE2	1:A:420:LEU:HD11	1.68	0.75
1:A:67[B]:MET:HE3	1:A:76:LYS:HZ3	1.52	0.74
1:A:70:LYS:HG2	3:A:639:HOH:O	1.88	0.73
1:A:449:MET:CE	3:A:670:HOH:O	2.33	0.70
1:A:67[B]:MET:CE	1:A:76:LYS:HZ3	2.05	0.69
1:A:100[B]:THR:HG23	3:A:615:HOH:O	1.93	0.69
1:A:309:THR:CG2	1:A:311:GLU:OE1	2.41	0.67
1:A:67[B]:MET:CE	1:A:76:LYS:NZ	2.59	0.65
1:A:120:LYS:HD2	3:A:493:HOH:O	1.97	0.64
1:A:390:ARG:HG2	1:A:390:ARG:HH11	1.63	0.62
1:A:309:THR:HG22	1:A:312:GLN:OE1	2.02	0.60
1:A:410:ARG:HG3	1:A:425:VAL:HG21	1.83	0.60
1:A:86:GLU:OE2	3:A:711:HOH:O	2.17	0.59
1:A:413:MET:CE	1:A:420:LEU:HD11	2.34	0.58
1:A:390:ARG:HH11	1:A:390:ARG:CG	2.19	0.56
1:A:100[A]:THR:HG22	1:A:101:PRO:HD2	1.87	0.55
1:A:448:MET:CG	3:A:670:HOH:O	2.46	0.55
1:A:309:THR:CG2	1:A:312:GLN:HB2	2.36	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67[B]:MET:HE3	1:A:76:LYS:HZ1	1.72	0.51
1:A:390:ARG:HG2	1:A:394:ARG:NH2	2.27	0.50
1:A:351:GLU:O	1:A:355:MET:HG3	2.13	0.48
1:A:212:LEU:HD22	1:A:216:TRP:CE2	2.49	0.47
1:A:72:LYS:O	1:A:100[A]:THR:HG23	2.14	0.47
1:A:413:MET:HE1	1:A:418:GLU:HG3	1.97	0.46
1:A:285:LYS:HE3	3:A:583:HOH:O	2.15	0.46
1:A:100[B]:THR:CG2	3:A:615:HOH:O	2.58	0.45
1:A:56:ARG:O	1:A:60:LEU:HD12	2.15	0.45
1:A:310:GLU:HG2	3:A:602:HOH:O	2.17	0.44
1:A:162:LYS:O	1:A:165:GLU:HG3	2.17	0.44
1:A:59:SER:HA	1:A:82:ARG:NH1	2.33	0.44
1:A:135:LEU:HD13	1:A:202:THR:HG22	2.00	0.43
1:A:413:MET:HE2	1:A:420:LEU:CD1	2.42	0.42
1:A:367:ILE:HA	1:A:371:GLU:OE1	2.21	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:ASN:OD1	1:A:162:LYS:NZ[1_565]	2.03	0.17
3:A:549:HOH:O	3:A:643:HOH:O[2_555]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	378/450 (84%)	372 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	334/382 (87%)	324 (97%)	10 (3%)	48 26

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	LYS
1	A	116	SER
1	A	120	LYS
1	A	145	LEU
1	A	165	GLU
1	A	309	THR
1	A	311	GLU
1	A	334	LYS
1	A	390	ARG
1	A	418	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	A	225	1	23,23,24	3.24	5 (21%)	29,32,34	1.84	9 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	225	1	-	0/12/31/32	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	225	CRO	CA2-C2	-5.39	1.42	1.48
1	A	225	CRO	C2-N3	-4.08	1.31	1.39
1	A	225	CRO	O2-C2	3.10	1.29	1.23
1	A	225	CRO	C1-N2	4.08	1.38	1.32
1	A	225	CRO	CB2-CA2	12.35	1.46	1.35

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	225	CRO	O2-C2-CA2	-4.20	128.68	130.95
1	A	225	CRO	CG2-CB2-CA2	-3.54	125.61	130.22
1	A	225	CRO	N3-C1-N2	-3.34	108.97	111.56
1	A	225	CRO	CG1-CB1-CA1	-2.41	108.98	112.53
1	A	225	CRO	C1-CA1-N1	-2.22	104.30	108.91
1	A	225	CRO	CD2-CG2-CB2	-2.00	114.38	121.23
1	A	225	CRO	C3-CA3-N3	2.07	117.53	113.00
1	A	225	CRO	CA1-C1-N2	2.36	127.07	123.83
1	A	225	CRO	CA2-C2-N3	3.55	105.18	103.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/450 (85%)	0.71	44 (11%) 6 7	16, 28, 50, 75	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	377	ALA	8.6
1	A	307	GLN	5.1
1	A	60	LEU	4.9
1	A	145	LEU	4.5
1	A	42	ARG	4.4
1	A	51	VAL	4.4
1	A	374	THR	4.3
1	A	41	ARG	4.3
1	A	162	LYS	4.3
1	A	311	GLU	4.2
1	A	309	THR	4.1
1	A	419	LYS	4.0
1	A	227	VAL	4.0
1	A	126	ASN	3.6
1	A	450	THR	3.5
1	A	87	ASP	3.5
1	A	312	GLN	3.4
1	A	378	ARG	3.3
1	A	220	VAL	3.3
1	A	313	ILE	3.1
1	A	292	ASP	3.1
1	A	54	ILE	3.1
1	A	386	GLU	3.0
1	A	310	GLU	3.0
1	A	390	ARG	2.9
1	A	59	SER	2.7
1	A	308	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	48	GLY	2.6
1	A	396	PHE	2.6
1	A	353	GLN	2.6
1	A	346	ASN	2.5
1	A	229	CYS	2.5
1	A	349	GLU	2.5
1	A	50	ALA	2.4
1	A	223	LEU	2.4
1	A	125	PRO	2.4
1	A	334	LYS	2.3
1	A	442	TYR	2.3
1	A	385	SER	2.3
1	A	219	LEU	2.2
1	A	222	THR	2.1
1	A	61	ILE	2.1
1	A	257	ILE	2.1
1	A	165	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CRO	A	225	22/23	0.96	0.16	-	15,17,21,23	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CA	A	456	1/1	0.96	0.09	-1.84	34,34,34,34	0
2	CA	A	455	1/1	0.95	0.04	-2.28	31,31,31,31	0
2	CA	A	454	1/1	0.96	0.06	-2.62	30,30,30,30	0
2	CA	A	453	1/1	1.00	0.04	-2.82	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.