



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:10 PM GMT

PDB ID : 1SH6  
Title : Crystal structure of actin-binding domain of mouse plectin  
Authors : Sevcik, J.; Urbanikova, L.  
Deposited on : 2004-02-25  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3948 atoms, of which 2006 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Plectin 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	230	Total	C	H	N	O	S	136	0	0
			3774	1188	1890	341	348	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP Q9QXS1
A	2	SER	-	CLONING ARTIFACT	UNP Q9QXS1
A	3	HIS	-	CLONING ARTIFACT	UNP Q9QXS1
A	4	MET	-	CLONING ARTIFACT	UNP Q9QXS1
A	5	GLU	-	CLONING ARTIFACT	UNP Q9QXS1
A	6	PHE	-	CLONING ARTIFACT	UNP Q9QXS1
A	244	GLU	-	CLONING ARTIFACT	UNP Q9QXS1
A	245	PHE	-	CLONING ARTIFACT	UNP Q9QXS1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	58	Total	H	O	116	0
			174	116	58		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### ● Molecule 1: Plectin 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	32.52Å 51.23Å 144.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 48.29 – 1.99	Depositor EDS
% Data completeness (in resolution range)	97.8 (50.00-2.00) 95.6 (48.29-1.99)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.62 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.198 , 0.300 0.207 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	33.6	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 46.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 16849 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3948	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.62	19/1921 (1.0%)	1.44	27/2597 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	201	GLU	CD-OE1	10.90	1.37	1.25
1	A	12	VAL	CB-CG2	8.97	1.71	1.52
1	A	19	LYS	CB-CG	-8.90	1.28	1.52
1	A	56	VAL	CB-CG1	-7.44	1.37	1.52
1	A	73	LYS	CE-NZ	7.36	1.67	1.49
1	A	229	VAL	CB-CG2	-6.98	1.38	1.52
1	A	182	ASN	CB-CG	-6.82	1.35	1.51
1	A	71	PHE	CB-CG	6.69	1.62	1.51
1	A	73	LYS	CD-CE	6.59	1.67	1.51
1	A	42	GLU	CD-OE1	6.34	1.32	1.25
1	A	97	ASP	CB-CG	-6.26	1.38	1.51
1	A	157	PHE	CD1-CE1	-6.03	1.27	1.39
1	A	29	TRP	CZ3-CH2	-5.60	1.31	1.40
1	A	162	ARG	CB-CG	-5.57	1.37	1.52
1	A	27	LYS	CD-CE	5.46	1.64	1.51
1	A	153	ARG	CB-CG	5.43	1.67	1.52
1	A	27	LYS	CE-NZ	5.33	1.62	1.49
1	A	74	LEU	CG-CD1	5.19	1.71	1.51
1	A	17	PHE	CE1-CZ	-5.17	1.27	1.37

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	39	ASP	CB-CG-OD1	9.47	126.83	118.30
1	A	84	LEU	CB-CG-CD1	8.05	124.69	111.00
1	A	43	ASP	CB-CG-OD2	8.02	125.52	118.30
1	A	98	ASP	CB-CG-OD2	7.94	125.44	118.30
1	A	27	LYS	CD-CE-NZ	7.81	129.67	111.70
1	A	82	ASP	CB-CG-OD2	7.77	125.29	118.30
1	A	221	ASP	CB-CG-OD2	7.74	125.27	118.30
1	A	26	ILE	CG1-CB-CG2	7.61	128.14	111.40
1	A	60	ASP	CB-CG-OD2	7.00	124.60	118.30
1	A	131	ASP	CB-CG-OD2	6.95	124.56	118.30
1	A	40	LEU	CA-CB-CG	6.76	130.84	115.30
1	A	45	ARG	NE-CZ-NH1	6.74	123.67	120.30
1	A	165	ARG	NE-CZ-NH2	-6.54	117.03	120.30
1	A	40	LEU	CB-CG-CD2	6.49	122.03	111.00
1	A	45	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	A	10	ASP	CB-CG-OD2	6.28	123.95	118.30
1	A	87	ARG	NE-CZ-NH1	-6.08	117.26	120.30
1	A	203	ASP	CB-CG-OD2	5.97	123.67	118.30
1	A	101	ASP	CB-CG-OD2	5.92	123.63	118.30
1	A	234	ASP	CB-CG-OD1	5.83	123.54	118.30
1	A	64	ARG	NE-CZ-NH1	5.75	123.17	120.30
1	A	153	ARG	NE-CZ-NH2	5.55	123.08	120.30
1	A	70	ARG	NE-CZ-NH1	5.49	123.05	120.30
1	A	144	ARG	CG-CD-NE	5.40	123.14	111.80
1	A	26	ILE	CB-CA-C	-5.32	100.97	111.60
1	A	153	ARG	NE-CZ-NH1	-5.25	117.67	120.30
1	A	173	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	131	ASP	Peptide
1	A	173	ARG	Peptide
1	A	91	LEU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1884	1890	1875	76	0
2	A	58	116	0	5	0
All	All	1942	2006	1875	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:LYS:CE	1:A:73:LYS:NZ	1.67	1.58
1:A:78:GLN:NE2	1:A:82:ASP:OD1	1.78	1.17
1:A:71:PHE:HD1	2:A:297:HOH:O	1.41	1.03
1:A:128:GLN:HB2	1:A:132:MET:HE1	1.43	0.97
1:A:149:TYR:HB2	1:A:152:LEU:HD11	1.48	0.94
1:A:128:GLN:HB2	1:A:132:MET:CE	2.05	0.85
1:A:133:THR:OG1	1:A:136:GLU:HG3	1.78	0.83
1:A:78:GLN:NE2	1:A:82:ASP:CG	2.32	0.82
1:A:174:HIS:CE1	1:A:236:MET:HB2	2.18	0.78
1:A:174:HIS:HE1	1:A:236:MET:HB2	1.49	0.77
1:A:122:ASP:N	1:A:122:ASP:OD2	2.19	0.75
1:A:157:PHE:CG	1:A:226:ILE:CD1	2.72	0.72
1:A:78:GLN:HE21	1:A:82:ASP:CG	1.93	0.69
1:A:132:MET:O	1:A:137:LYS:NZ	2.24	0.69
1:A:157:PHE:CG	1:A:226:ILE:HD11	2.28	0.68
1:A:120:ILE:O	1:A:123:ILE:HD12	1.94	0.68
1:A:188:THR:OG1	1:A:191:GLU:HG3	1.95	0.67
1:A:48:HIS:HE1	2:A:270:HOH:O	1.77	0.66
1:A:42:GLU:HA	1:A:45:ARG:HD3	1.78	0.64
1:A:95:ARG:HB3	2:A:287:HOH:O	1.98	0.64
1:A:174:HIS:CE1	1:A:236:MET:CB	2.80	0.64
1:A:71:PHE:CD1	2:A:297:HOH:O	2.29	0.63
1:A:8:GLU:HG3	1:A:9:ARG:N	2.15	0.61
1:A:28:HIS:CD2	1:A:33:ALA:CB	2.84	0.60
1:A:157:PHE:CD2	1:A:226:ILE:HD11	2.37	0.59
1:A:157:PHE:CB	1:A:226:ILE:HD11	2.31	0.59
1:A:87:ARG:HH11	1:A:117:HIS:HE1	1.48	0.59
1:A:28:HIS:HD2	1:A:33:ALA:HB1	1.68	0.59
1:A:149:TYR:OH	1:A:172:HIS:O	2.21	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:VAL:H	1:A:128:GLN:NE2	2.02	0.57
1:A:157:PHE:CD2	1:A:226:ILE:CD1	2.89	0.56
1:A:179:ILE:HD11	1:A:181:MET:SD	2.46	0.56
1:A:170:ILE:HG22	1:A:236:MET:CE	2.36	0.56
1:A:213:GLU:CD	1:A:213:GLU:H	2.09	0.55
1:A:135:LYS:HD2	1:A:135:LYS:O	2.07	0.55
1:A:32:GLU:H	1:A:32:GLU:CD	2.11	0.54
1:A:172:HIS:HB2	1:A:179:ILE:CG1	2.38	0.53
1:A:24:HIS:HD2	1:A:27:LYS:CE	2.21	0.53
1:A:71:PHE:CD2	1:A:74:LEU:HD12	2.44	0.52
1:A:125:VAL:HG11	1:A:141:TRP:HB2	1.90	0.52
1:A:222:GLU:O	1:A:226:ILE:HG12	2.10	0.52
1:A:91:LEU:O	1:A:92:VAL:C	2.48	0.52
1:A:36:HIS:HD2	2:A:252:HOH:O	1.93	0.52
1:A:175:LYS:N	1:A:176:PRO:HD3	2.25	0.51
1:A:28:HIS:CD2	1:A:33:ALA:HB3	2.46	0.50
1:A:175:LYS:N	1:A:176:PRO:CD	2.75	0.50
1:A:175:LYS:HB3	1:A:178:LEU:HD22	1.93	0.49
1:A:174:HIS:CA	1:A:176:PRO:HD3	2.42	0.49
1:A:133:THR:HG1	1:A:136:GLU:HG3	1.76	0.48
1:A:28:HIS:HD2	1:A:33:ALA:CB	2.25	0.48
1:A:128:GLN:CB	1:A:132:MET:HE1	2.30	0.48
1:A:40:LEU:HD22	1:A:44:LEU:HD11	1.97	0.47
1:A:62:LEU:HD21	1:A:83:TYR:CD1	2.49	0.47
1:A:87:ARG:NH1	1:A:117:HIS:HE1	2.13	0.46
1:A:62:LEU:HD23	1:A:62:LEU:HA	1.87	0.45
1:A:174:HIS:CD2	1:A:174:HIS:N	2.82	0.45
1:A:93:ASN:ND2	1:A:93:ASN:O	2.50	0.45
1:A:91:LEU:HA	1:A:91:LEU:HD12	1.73	0.44
1:A:225:ILE:O	1:A:229:VAL:HG13	2.16	0.44
1:A:14:LYS:HE3	1:A:41:TYR:CD2	2.53	0.44
1:A:132:MET:HA	1:A:136:GLU:OE1	2.18	0.43
1:A:236:MET:O	1:A:236:MET:CG	2.65	0.43
1:A:159:THR:O	1:A:162:ARG:HB2	2.18	0.43
1:A:103:ASN:HD22	1:A:106:LEU:HB2	1.84	0.43
1:A:174:HIS:C	1:A:176:PRO:HD3	2.39	0.43
1:A:183:LYS:HD2	1:A:187:GLN:NE2	2.35	0.42
1:A:24:HIS:HD2	1:A:27:LYS:HE3	1.83	0.41
1:A:28:HIS:ND1	1:A:29:TRP:N	2.69	0.41
1:A:175:LYS:HB3	1:A:178:LEU:CD2	2.51	0.41
1:A:8:GLU:HG3	1:A:9:ARG:H	1.84	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:LYS:HD2	1:A:135:LYS:C	2.41	0.41
1:A:32:GLU:O	1:A:35:ARG:NH1	2.53	0.41
1:A:24:HIS:HA	1:A:27:LYS:HE3	2.02	0.40
1:A:40:LEU:HD22	1:A:44:LEU:CD1	2.52	0.40
1:A:132:MET:HE2	1:A:132:MET:HB2	1.95	0.40
1:A:8:GLU:CG	1:A:9:ARG:N	2.80	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	228/245 (93%)	214 (94%)	11 (5%)	3 (1%)	15 7

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	69	MET
1	A	172	HIS
1	A	67	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	208/223 (93%)	178 (86%)	30 (14%)	4 2

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	GLU
1	A	15	LYS
1	A	26	ILE
1	A	30	ARG
1	A	40	LEU
1	A	56	VAL
1	A	66	LYS
1	A	69	MET
1	A	91	LEU
1	A	93	ASN
1	A	119	GLN
1	A	122	ASP
1	A	125	VAL
1	A	126	SER
1	A	131	ASP
1	A	135	LYS
1	A	140	LEU
1	A	144	ARG
1	A	152	LEU
1	A	162	ARG
1	A	174	HIS
1	A	177	MET
1	A	178	LEU
1	A	179	ILE
1	A	182	ASN
1	A	183	LYS
1	A	198	SER
1	A	208	ARG
1	A	219	GLN
1	A	231	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	HIS
1	A	34	GLN
1	A	36	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	48	HIS
1	A	72	HIS
1	A	78	GLN
1	A	93	ASN
1	A	96	ASN
1	A	103	ASN
1	A	117	HIS
1	A	128	GLN
1	A	187	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	230/245 (93%)	0.27	8 (3%)	48 49	21, 35, 60, 68	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	131	ASP	3.3
1	A	152	LEU	3.2
1	A	92	VAL	2.9
1	A	174	HIS	2.8
1	A	179	ILE	2.7
1	A	153	ARG	2.7
1	A	148	GLY	2.5
1	A	90	LYS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.